

Durham Research Online

Deposited in DRO:

20 November 2013

Version of attached file:

Published Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Le Delliou, M. and Lacey, Cedric G. and Baugh, C. M. and Guiderdoni, B. and Bacon, R. and Courtois, H. and Sousbie, T. and Morris, S. L. (2005) 'The abundance of Ly emitters in hierarchical models.', *Monthly Notices of the Royal Astronomical Society.*, 357 (1). L11-L15.

Further information on publisher's website:

<http://dx.doi.org/10.1111/j.1745-3933.2005.00007.x>

Publisher's copyright statement:

This article has been published in the *Monthly Notices of the Royal Astronomical Society* © 2005 The Authors. Published by Oxford University Press on behalf of The Royal Astronomical Society. All rights reserved.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

The abundance of Ly α emitters in hierarchical models

M. Le Delliou,^{1,2*} C. Lacey,^{1,3} C. M. Baugh,³ B. Guiderdoni,⁴ R. Bacon,¹
H. Courtois,¹ T. Sousbie¹ and S. L. Morris³

¹Centre de Rech. Astron. de Lyon (CRAL), 9 avenue Charles André, 69561 Saint Genis Laval Cedex, France

²GAM – UMR 5139 – CNRS/in2p3, CC 85, U. M. II, Place Eugène Bataillon, 34095 Montpellier Cedex 5, France

³Institute for Computational Cosmology, University of Durham, Science Laboratories, South Road, Durham DH1 3LE

⁴Institut d’Astrophysique de Paris, CNRS, 98 bis Boulevard Arago, 75014 Paris, France

Accepted 2004 November 5. Received 2004 November 4; in original form 2004 July 29

ABSTRACT

We present predictions for the abundance of Ly α emitters in hierarchical structure formation models. We use the GALFORM semi-analytical model of galaxy formation to explore the impact on the predicted counts of varying assumptions about the escape fraction of Ly α photons, the redshift at which the Universe reionized and the cosmological density parameter. A model with a fixed escape fraction gives a remarkably good match to the observed counts over a wide redshift interval. The counts at bright fluxes are dominated by ongoing starbursts. We present predictions for the expected counts in a typical observation with the Multi-Unit Spectroscopic Explorer (MUSE) instrument proposed for the Very Large Telescope.

Key words: galaxies: formation – galaxies: high-redshift – cosmology: miscellaneous.

1 INTRODUCTION

Dedicated narrow-band searches for Ly α emitters have proven to be very efficient at detecting high-redshift galaxies (e.g. Hu & McMahon 1996; Cowie & Hu 1998; Rhoads et al. 2000; Ouchi et al. 2003). Objects found by this technique have to be confirmed spectroscopically, to rule out possible low-redshift interlopers that may arise due to emission lines other than Ly α falling within the targeted wavelength interval. Nevertheless, a significant fraction of the detections appear to be bona fide Ly α emitters, and the number of objects accumulated to date by this technique in the redshift interval $2.4 < z < 6.6$ is quite impressive. The Ly α emission line is also found in a significant fraction of Lyman break galaxies (e.g. Steidel et al. 1996), which are selected on the basis of their continuum emission.

The ubiquity of the Ly α line is at face value surprising, given that it is resonantly scattered by atomic hydrogen, and so is easily absorbed by even a small amount of dust in a neutral gaseous medium. It is suspected that most Ly α emitters have galactic winds (as is the case with Lyman break galaxies) which allow Ly α photons to escape from the galaxy after only a limited number of resonant scatterings (Kunth et al. 1998; Pettini et al. 2001). The Ly α line typically shows an asymmetric profile characteristic of such a process (e.g. Ahn 2004). The physics of this phenomenon is complicated, however, and remains poorly understood.

We present here the first predictions for the abundance of Ly α emitters at different redshifts made using a model which follows the formation and evolution of galaxies in a hierarchical universe. In previous work, simple, ad hoc prescriptions have been used to

assign star formation rates to dark matter haloes (Haiman & Spaans 1999; Santos et al. 2004). In this Letter, we use a semi-analytical model to make an *ab initio* calculation of the distribution of galaxy masses and star formation rates at different redshifts (e.g. Cole et al. 1994; Kauffmann, Guiderdoni & White 1994; Baugh et al. 1998; Somerville & Primack 1999; Hatton et al. 2003). The model we use is able to reproduce the observed properties of galaxies both locally and at high redshift (Cole et al. 2000; Baugh et al. 2005). The abundance of Ly α emitters is sensitive to the adopted cosmological model and to astrophysical phenomena, such as the fraction of Ly α photons escaping from galaxies and the distribution of galactic dust. Semi-analytical models are ideally suited to the exploration of such a parameter space.

The semi-analytical model is described in Section 2. In Section 3, we first present a compilation of the available observational data on the abundance of Ly α emitters at different redshifts, and then compare these data with the predictions from our models. Finally, we present our conclusions in Section 4.

2 THE MODEL

We use the semi-analytical model of galaxy formation, GALFORM, to make predictions for the abundance of Ly α emitters as a function of Ly α flux and redshift. The GALFORM model is described in full in Cole et al. (2000) and Benson et al. (2003); further details of the model used in this Letter are given in Baugh et al. (2005). The GALFORM model computes the star formation histories for the whole galaxy population. The following steps are taken to compute the Ly α emission from a model galaxy. (i) The number of Lyman continuum photons is computed from the star formation history in the model galaxy and the stellar initial mass function (IMF). In the

*E-mail: Morgan.LeDelliou@gamum2.in2p3.fr

Table 1. The parameters of the semi-analytical models for which the abundances of Ly α emitters are predicted. The first column gives the model label. The next three columns give the basic cosmological parameters: the density parameter, Ω , the cosmological constant, Λ , and the amplitude of density perturbations, as specified by σ_8 (σ_8 values are taken from Eke, Cole & Frenk 1996). In each case the baryon density is $\Omega_b = 0.04$ and the Hubble constant is $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$. Column five gives the redshift z_{reion} at which the universe is assumed to reionize. Column six gives the fraction f_{esc} of Ly α photons that escape from the model galaxies. In the case of model C, the escape fraction is computed from a dust extinction model, as described in the text. The final column gives the number counts of Ly α emitters for a reference MUSE observation. This is the number of emitters in 1 square arcmin in the redshift interval $2.8 < z < 6.7$ brighter than $3.9 \times 10^{-19} \text{ erg s}^{-1} \text{ cm}^{-2}$.

Model	Ω	Λ	σ_8	z_{reion}	f_{esc}	MUSE counts
A	0.3	0.7	0.93	10	0.02	70
B	0.3	0.7	0.93	10	0.1	248
C	0.3	0.7	0.93	10	dust	366
D	0.3	0.7	0.93	6	0.02	163
E	0.3	0.7	0.93	20	0.02	77
F	0.2	0.8	1.15	10	0.02	145
G	1	0	0.52	10	0.02	59

Table 2. Data compilation. The data are divided into unit redshift intervals: the following symbols are used to denote data from each redshift interval in the figures (filled square: [$z < 3$], downward filled triangle: [$3 < z < 4$], upward filled triangle: [$4 < z < 5$], open circle: [$z = 5.1$], filled circle: [$5 < z < 6$], cross: [$6 < z < 7$]). Column 1: redshift; column 2: redshift interval; column 3: Ly α flux (in $10^{-17} \text{ erg cm}^{-2} \text{ s}^{-1}$); column 4: cumulative counts per unit solid angle per unit redshift (in arcmin^{-2}); column 5: Poisson error on counts (in arcmin^{-2}); column 6: number of Ly α emitters; column 7: area of survey (in arcmin^2); column 8: factor applied to correct for contamination by low- z interlopers; column 9: method (NBF = narrow band filter, LS = long-slit spectroscopy); column 10: method used to reject or correct for low- z interlopers (EW = equivalent width, spec on N = follow-up spectroscopy of N objects); column 11: reference (D01: Dawson et al. 2001; F03: Fujita et al. 2003; H98: Hu, Cowie & McMahon 1998; H02: Hu et al. 2002; K03: Kodaira et al. 2003; K00: Kudritzki et al. 2000; O03: Ouchi et al. 2003; R00: Rhoads et al. 2000; R03: Rhoads et al. 2003; Sa04: Santos et al. 2004; S03: Shimasaku et al. 2003; S04: Shimasaku et al. 2004; Ste00: Steidel et al. 2000; Sti01: Stiavelli et al. 2001).

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
z	Δz	f	$\frac{d^2 N}{dz d\Omega} (> f)$	$\Delta(\frac{d^2 N}{dz d\Omega})$	Nobj	Area	Fcorr	Method	Confirmation	Ref.
2.42	0.14	20	0.33	0.04	58	1200	0.65	NBF	EW/colour	Sti01
3.09	0.07	2	2.3 ^a	0.3	12 ^a	78	0.94	NBF	EW/colour	Ste00
3.13	0.04	2	3.8	1.3	8	49	0.7	NBF	spec on 10	K00
3.43	0.06	1.5	3.5	0.9	16	75	0.87	NBF	spec on 15	H98
3.72	0.23	6.4	0.26	0.09	8	130	0.35	NBF	colours	F03
4.39	0.07	2.6	0.97	0.11	75	1100	0.33	NBF	spec on 3	R00
4.54	0.06	1.5	1.3	0.9	2	24	0.67	NBF	spec on 3	H98
4.79	0.08	0.5	0.46	0.07	41	1100	0.8	NBF	–	S04
4.86	0.06	0.5	0.52	0.09	34	1100	0.8	NBF	spec on 5	S03
4.86	0.06	0.3	1.6	0.2	52	540	0.6	NBF	colours	O03
5.1	1.0	0.012 ^b	48 ^b	48	1	0.02 ^b	–	LS	–	Sa04
“	“	0.037 ^b	30 ^b	15	4	0.14 ^b	–	“	–	“
“	“	0.12 ^b	4.0 ^b	2.3	3	0.75 ^b	–	“	–	“
“	“	0.37 ^b	0.89 ^b	0.51	3	3.4 ^b	–	“	–	“
“	“	1.2 ^b	0.14 ^b	0.14	1	7.5 ^b	–	“	–	“
5.3	1.0	2?	2.3	1.0	5	2.2	–	LS	–	D01
5.7	0.13	1.5	0.14	0.04	13	710	0.75	NBF	spec on 4	R03
6.56	0.10	0.6 ^b	20 ^b	20	1	0.46 ^b	1	NBF	spec on 1	H02
6.56	0.11	0.9	0.18	0.05	16	810	0.22	NBF	spec on 9	K03

Notes: ^acorrected for factor 6 overdensity; ^bcorrected for gravitational lensing.

Baugh et al. (2005) model, quiescent star formation in galactic discs produces stars with a Kennicutt (1998) IMF, whereas bursts of star formation triggered by galaxy mergers produce a flat (‘top-heavy’) IMF. (ii) The luminosity of the Ly α line is computed assuming that all Lyman continuum photons are absorbed in H II regions and produce Ly α photons according to case B recombination (Osterbrock 1989). (iii) The observed Ly α line emission depends on how many Ly α photons escape from the galaxy. We have taken two approaches to estimating the escape fraction. In the first, we simply assume that a fixed fraction, f_{esc} , of Ly α photons escape from the galaxy. Physically, this might arise if a fraction of the Ly α photons escape through holes in the galactic gas and dust distribution. In the second approach, we calculate the absorption of Ly α photons by a diffuse dust medium having the same spatial distribution as the stars, ignoring resonant scattering by neutral hydrogen. This may mimic what occurs if resonant scattering is suppressed by a galactic wind. We make a self-consistent calculation of the dust optical depth of the model galaxies, using the predicted galaxy scalelength, gas mass and metallicity. Full details of this dust extinction model can be found in Cole et al. (2000). We plan a more detailed calculation of the escape fraction of Ly α photons, including the effects of resonant scattering and gas outflows, in a future paper.

We explore the impact on the abundances of Ly α emitters of varying the redshift at which the Universe was reionized, z_{reion} , which is still poorly constrained.

We examine the consequences of making three choices: (i) $z_{\text{reion}} = 10$, (ii) $z_{\text{reion}} = 20$ and (iii) $z_{\text{reion}} = 6$. The first two values are consistent with the optical depth to last scattering suggested by the *Wilkinson Microwave Anisotropy Probe* (WMAP) measurement of the correlation between microwave background temperature and polarization (Kogut et al. 2003). The latter value is suggested by the detection of a Gunn–Peterson trough in the spectrum of a $z = 6.28$ quasar by Becker et al. (2001). In the model, gas is prevented from cooling in haloes with circular velocities below 60 km s^{-1} for redshifts $z < z_{\text{reion}}$. The models that we consider are listed in Table 1. Models A–E reproduce the observed luminosity function for the local galaxy population (e.g. Baugh et al. 2005). Finally, we also consider the effect on the counts of Ly α emitters of varying the cosmological density parameter, Ω (models F and G). Note that we have not attempted to vary any other parameters of the GALFORM model in these two cases to force the model to reproduce the local galaxy luminosity function.

3 THE OBSERVATIONAL DATA AND MODEL PREDICTIONS

We list in Table 2 a compilation of published observational data on number counts of Ly α emitters at different observed line fluxes and redshifts from ‘blank field’ surveys that we will compare against our model predictions. Note that some of the quantities listed in Table 2 are derived from the information given in the original sources, so we present this compilation in order to facilitate future comparisons between data and model predictions by other authors. We do not include data from surveys targeted around high- z objects such as quasars (e.g. Fynbo, Moller & Thomsen 2001), as in this case the number density of Ly α emitters may be biased by an unknown factor. A few of the surveys used spectroscopy to search for Ly α emitters directly. However, most used narrow-band imaging to identify objects having a strong emission line at the wavelength corresponding to Ly α at a particular redshift. Samples of objects obtained in this way are generally contaminated by lower redshift galaxies for which some other emission line (e.g. [O II]3737, [O III]5007, $H\alpha$) happens to fall at the same wavelength. This contamination is typically estimated and removed using the equivalent width of the emission line and/or broad-band colours, or from follow-up spectroscopy on a subsample of the objects. The methods used in the different surveys are indicated in the table.

The observational data are plotted in Fig. 1, with different symbols indicating data from a given unit redshift range. The predictions of our fiducial model, A from Table 1, are shown by the solid lines in Fig. 1. The escape fraction f_{esc} was set to give a reasonable match to the observed number counts at $z \sim 3$. This simple model, in which the escape fraction is independent of redshift and galaxy properties, does a surprisingly good job of matching the observed counts at different redshifts.

The separate contributions to the counts of Ly α emitters from galaxies that are forming stars quiescently and from ongoing starbursts (which in our model are triggered by galaxy mergers) are shown in Fig. 2 for model A over the redshift interval $z = 3$ –4. Quiescently star-forming galaxies dominate at fainter fluxes, whereas starbursts account for the brighter Ly α sources. This is largely due to the flat IMF assumed in starbursts, which typically yields 10 times the number of Lyman continuum photons for a given amount of star




Figure 1. The predicted number of Ly α emitters in model A, compared to the observational data compiled in Table 2. The observational data are divided into different redshift ranges, as indicated by the lower key, and we have plotted only datapoints based on more than one galaxy. Model predictions are shown by lines, as indicated by the upper key. The thickness of the lines increases with redshift. The vertical and horizontal dashed lines indicate the sensitivity limits in flux and number density for a reference MUSE observation (see text).




Figure 2. The predicted number counts of Ly α emitters in model A, in the redshift range $z = 3$ –4. The contributions from quiescent star forming galaxies and ongoing starbursts are plotted separately, as indicated in the key. The vertical and horizontal dashed lines are the same as in Fig. 1.

formation, compared with the Kennicutt (1998) IMF that we adopt for quiescent star formation.

In Fig. 3, we present the predicted counts of Ly α emitters as a function of redshift for the different GALFORM models listed in Table 1. The results for model A, our fiducial model, are reproduced for reference in each panel. In Fig. 3(b) we show the effect of increasing the escape fraction f_{esc} by a factor of 5. Fig. 3(b) also shows




Figure 3. The predicted counts of Ly α emitting galaxies, plotted in different redshift intervals, as indicated by the key. (a) Shows the predictions for model A; a subset of these are reproduced in each panel for reference. The data from Fig. 1 are also plotted here using the same symbols as before. The remaining panels show the impact on the predictions of changing different aspects of the model: (b) Varying the escape fraction or using a dust model. (c) Varying the redshift of reionization. (d) Varying the cosmological density parameter.

that assuming a fixed escape fraction $f_{\text{esc}} = 0.1$ produces similar number counts to a model in which the Ly α photons are absorbed by diffuse dust *without* resonant scattering, as might occur in a galactic wind. Fig. 3(c) shows the impact on the predicted counts of varying the redshift at which the Universe is reionized. For model D, with $z_{\text{reion}} = 6$, the number of faint Ly α sources changes substantially for $5 < z < 6$ and $4 < z < 5$, compared with the cases where $z_{\text{reion}} \geq 10$. This is because in model D gas is still able to cool in low-mass haloes (i.e. with circular velocities below 60 km s^{-1}) up to $z = 6$, and is available to form stars and thus generate Lyman continuum photons. Finally, Fig. 3(d) illustrates the effect of changing the cosmological density parameter whilst retaining a flat universe. The changes in the model predictions in this case primarily reflect the change in the normalization of density fluctuations, as specified by σ_8 in Table 1, which is adjusted to reproduce the local abundance of rich clusters when the density parameter is varied.

4 DISCUSSION AND CONCLUSIONS

The conclusion of this first study to look at the predictions of hierarchical models for the number of Ly α emitters is that simple models do remarkably well at reproducing the observed counts. Assuming that typically just 2 per cent of the Ly α photons escape from high- z galaxies, a value chosen to match the observed counts at $z \sim 3$ is

sufficient to give a reasonable match to the observed counts at faint fluxes over the redshift interval $2 < z < 6$.

This study demonstrates the capability of semi-analytical modelling to make predictions that can serve as an input into the design of new instruments. The Multi-Unit Spectroscopic Explorer (MUSE) (Henault et al. 2004) has been proposed to the European Southern Observatory (ESO) as a second-generation instrument for the Very Large Telescope. MUSE will be able to identify Ly α emitters over a redshift interval $2.8 < z < 6.7$ over a field of view of 1 arcmin^2 . An exposure of 80 h will reach a 5σ sensitivity of $3.9 \times 10^{-19} \text{ erg s}^{-1} \text{ cm}^{-2}$. We predict that MUSE will be able to detect a large number of such objects at this flux limit: around 70–400 per arcmin^2 (see the final column of Table 1). Observations with MUSE will be able to exclude some of the models we have considered, and therefore remove some of the uncertainties in our modelling of Ly α emission. More importantly, such observations will provide a critical test of our ideas about star formation in objects at high redshifts.

ACKNOWLEDGMENTS

CGL and CMB thank their GALFORM collaborators Andrew Benson, Shaun Cole and Carlos Frenk for allowing them to use the model in this paper. We acknowledge Bianca Poggianti for valuable discussions about modelling emission lines. We also thank the

referee for providing a speedy and helpful report that allowed us to improve the clarity of this Letter. MLeD would like to thank the CRAL (Observatoire de Lyon) for hospitality and financial support during the completion of this work. CMB and MLeD acknowledge support from the Royal Society. CGL is funded by PPARC.

REFERENCES

- Ahn S.-H., 2004, ApJ, 601, L25
 Baugh C. M., Cole S., Frenk C. S., Lacey C. G., 1998, ApJ, 498, 504
 Baugh C. M., Lacey C. G., Frenk C. S., Granato G. L., Silva L., Bressan A., Benson A. J., Cole S., 2005, MNRAS, 356, 1191
 Becker R. H. et al., 2001, AJ, 122, 2850
 Benson A. J., Frenk C. S., Baugh C. M., Cole S., Lacey C. G., 2003, ApJ, 599, 38
 Cole S., Aragon-Salamanca A., Frenk C. S., Navarro J. F., Zepf S., 1994, MNRAS, 271, 781
 Cole S., Lacey C. G., Baugh C. M., Frenk C. S., 2000, MNRAS, 319, 168
 Cowie L. L., Hu E. M., 1998, AJ, 115, 1319
 Dawson S., Stern D., Bunker A. J., Spinrad H., Dey A., 2001, AJ, 122, 598
 Eke V. R., Cole S., Frenk C. S., 1996, MNRAS, 282, 263
 Fujita S. S. et al., 2003, AJ, 125, 13
 Fynbo J. U., Moller P., Thomsen B., 2001, A&A, 374, 443
 Haiman Z., Spaans M., 1999, ApJ, 518, 138
 Hatton S., Devriendt J. E. G., Ninin S., Bouchet F. R., Guiderdoni B., Vibert D., 2003, MNRAS, 343, 75
 Henault F., Bacon R., Content R., Lantz B., Laurent F., Lemonnier J.-P., Morris S. L., 2004, in Mazuray L., Rogers P. J., Wartmann R., eds, Proc. SPIE Vol. 5249, Optical Design and Engineering. SPIE, Bellingham WA, p. 134
 Hu E. M., McMahon R. G., 1996, Nat, 382, 231
 Hu E. M., Cowie L. L., McMahon R. G., 1998, ApJ, 502, L99
 Hu E. M., Cowie L. L., McMahon R. G., Capak P., Iwamuro F., Kneib J.-P., Maihara T., Motohara K., 2002, ApJ, 568, L75
 Kauffmann G., Guiderdoni B., White S. D. M., 1994, MNRAS, 267, 981
 Kennicutt R. C., 1998, ApJ, 498, 541
 Kodaira K. et al., 2003, PASJ, 55, L17
 Kogut A. et al., 2003, ApJS, 148, 161
 Kudritzki R.-P. et al., 2000, ApJ, 536, 19
 Kunth, D., Mas-Hesse, J. M., Terlevich, E., Terlevich, R., Lequeux, J., Fall, S. M., 1998, A&A, 334, 11
 Osterbrock D. E., 1989, Astrophysics of Gaseous Nebulae and Active Galactic Nuclei. Univ. Science Book, Mill Valley, CA
 Ouchi M. et al., 2003, ApJ, 582, 60
 Pettini M., Steidel C. C., Adelberger K. L., Dickinson M., Giavalisco M., 2001, ApJ, 528, 96
 Rhoads J. E., Malhotra S., Dey A., Stern D., Spinrad H., Jannuzi B. T., 2000, ApJ, 545, L85
 Rhoads J. E. et al., 2003, AJ, 125, 1006
 Santos M. R., Ellis R. S., Kneib J. P., Richard J., Kuijken K., 2004, ApJ, 606, 683
 Shimasaku K. et al., 2003, ApJ, 586, L111
 Shimasaku K. et al., 2004, ApJ, 605, L93
 Somerville R. S., Primack J. R., 1999, MNRAS, 310, 1087
 Steidel C. C., Giavalisco M., Dickinson M., Adelberger K. L., 1996, AJ, 112, 352.
 Steidel C. C., Adelberger K. L., Shapley A. E., Pettini M., Dickinson M., Giavalisco M., 2000, ApJ, 532, 170
 Stiavelli M., Scarlata C., Panagia N., Treu T., Bertin G., Bertola F. 2001, ApJ, 561, L37

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.