Durham Research Online

Deposited in DRO:

13 March 2018

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Schoonmaker, R. and Lancaster, T. and Clark, S. J. (2018) 'Quantum mechanical tunneling in the automerization of cyclobutadiene.', *Journal of chemical physics.*, 148 (10). p. 104109.

Further information on publisher's website:

<https://doi.org/10.1063/1.5019254>

Publisher's copyright statement:

© 2018 American Institute of Physics. This article may be downloaded for personal use only. Any other use requires prior permission of the author and the American Institute of Physics. The following article appeared in Schoonmaker, R., Lancaster, T. Clark, S. J. (2018). Quantum mechanical tunneling in the automerization of cyclobutadiene. *The Journal of Chemical Physics* 148(10): 104109 and may be found at <https://doi.org/10.1063/1.5019254>

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

R. Schoonmaker, T. Lancaster, and S. J. Clark
Durham University, Department of Physics, South Road, Durham, DH1 3LE, UK
 (Dated: February 21, 2018)

Cyclobutadiene has a four-membered carbon ring with two double bonds, but this highly strained molecular configuration is almost square and, via a coordinated motion, the nuclei quantum mechanically tunnel through the high-energy square state to a configuration equivalent to the initial configuration under a 90° rotation. This results in a square ground state comprised of a superposition of two molecular configurations, that is driven by quantum tunnelling. Using a quantum mechanical model, and an effective nuclear potential from density functional theory, we calculate the vibrational energy spectrum and the accompanying wavefunctions. We use the wavefunctions to identify motions of the molecule, and detail how different motions can enhance or suppress the tunnelling rate. This is relevant for kinematics of tunnelling-driven reactions, and we discuss these implications. We are also able to provide a qualitative account of how the molecule will respond to an external perturbation, and how this may enhance or suppress IR-active vibrational transitions.

INTRODUCTION

The motions and locations of nuclei underpin how we define molecular structure, structural transitions, and chemical reactions. Cyclobutadiene as a free molecule exhibits nuclear delocalization that determines the point group symmetry, for although the lowest-energy electronic configuration suggests that the molecule is rectangular with two double and two single bonds, the molecule automerizes via nuclear tunnelling which results in an overall square symmetry [1–3]. While this automerization is similar to a reaction, since it involves the motion of nuclei to a different structural conformation, it is distinct since the reactant is chemically identical to the product, and the transition does not require thermal or collision processes [4]. These distinctive qualities therefore present an opportunity to study how structure changes when nuclei tunnel, without additional complications such as dynamic molecular collisions. Here we calculate the energy eigenstates of the nuclear motion that correspond to active infra-red (IR) excitations, and their corresponding nuclear wavefunctions. With these wavefunctions we are able to describe and explain how the different motions of the molecule can enhance or suppress the rate of tunnelling.

There are two electronic configurations at low energies in cyclobutadiene, a singlet state and a triplet state [5]. Historically it was unclear which state was most important in the low temperature behaviour of cyclobutadiene [6]. While it was found that a D_{2h} singlet state has the lowest energy [4, 5], the electronic structure of the triplet state remains of considerable interest due to conflicting accounts of the aromatic stabilization or antiaromatic destabilization, and the role of ring strain in the electronic structure [7–10]. At low temperatures the molecule tunnels between two D_{2h} singlet configurations [1, 11], and theoretical studies of this automerization suggest that a D_{4h} triplet state may determine the effective height of the reaction barrier [12–14]. However the precise structural configuration where the energies of the singlet and triplet state are equivalent is not precisely

known [15], if it exists at all [16, 17]. Additionally in the free molecule the strength of the spin-orbit interaction, which permits a transition between the singlet and triplet electronic states, is small by comparison to the kinetic energy of the nuclei [18], and this will suppress the transition between the singlet and triplet states. In this work, we therefore assume that singlet-triplet transitions are symmetry-forbidden, so only singlet states will be considered for tunnelling through the intermediate D_{4h} structural configuration.

Some theoretical studies of tunnelling in the singlet ground state focus on the energetic surface and barrier height of this nuclear tunnelling process, as these are required to calculate the tunnelling rate [13, 16, 19, 20]. In chemical systems, the most straightforward approaches to calculate tunnelling rates use WKB(J) formalism [14, 21, 22], however this formalism requires that reverse-tunnelling processes are negligible. There is therefore reason to doubt its applicability to the case of cyclobutadiene where the two configurations are symmetry-equivalent and reverse tunnelling can readily occur. Another common method of calculating rates uses instantons [23], which can be used to map the system onto a one-dimensional tunnelling problem [24, 25]. However as the nuclear potential is anharmonic in cyclobutadiene it is likely that an instanton-based model will not be able to accurately reproduce the tunnelling rate and other properties, such as the motion of the hydrogen nuclei, which is dependent on wavefunction configurations different from those on the instanton path.

Other theoretical work focuses on the determination of the Raman spectrum of the molecule, that is important to recognize and classify the experimental Raman response of the molecule [26, 27]. Due to time evolution's dependency on the Hamiltonian, the calculations of the Raman spectrum and the tunnelling rate in this single-molecule system are equivalent, and it is possible to study tunnelling under a Hamiltonian framework [2, 26].

The behaviour of interest to this work is tunnelling between cyclobutadiene's rectangular configurations. This tunnelling results in a small correction where each of the

FIG. 1. The (rectangular) a_g motions permitted in cyclobutadiene: C-C and C-H bond stretches, and the C-H bond waggle. The σ_x, σ_y reflection symmetries are also shown.

known Raman peaks that have been previously classified under a rectangular symmetry divide into pairs. A previous study by Čársky et al.[26] used a three-dimensional Taylor expansion on a potential calculated under the generalized valence bond method, to approximate the wavefunction and to calculate the energy separation of these pairs. Unfortunately the energy separation that they predicted was not experimentally observed when cyclobutadiene was bound in a solid matrix, possibly due to environmental breaking of the square symmetry[3, 28].

In this paper we propose a general method for calculating the energy eigenstates and wavefunctions of nuclei bound in molecules based on an effective potential energy surface determined by the electrons and Coulombic forces. We use this to calculate the Raman spectrum and molecular wavefunctions using the symmetry-conserving motions permitted under the rectangular symmetry. We are able to determine how the nuclear motions for different Raman resonance states affect the rate of tunnelling. We show how the calculation and interpretation of the wavefunction reveals the sensitivity of the system to environmental effects, and how the inclusion of hydrogenic motion affects the carbon motion and the strength of the tunnelling interaction.

METHODS

Assuming adiabatic separation between the electrons and nuclei, we consider the four in-plane (rectangular, D_{2h}) a_g symmetry-preserving motions of the molecule, as shown in Fig. 1. This assumes the effective poten-

FIG. 2. The calculation basis $\mathbf{r}'_{CH} = (d_{CH}, l_{CH})$, $\mathbf{r}'_C = (d_C, l_C)$ used here, shown on a single quadrant of the molecule. This new coordinate set is related to the absolute positions by $\mathbf{r}'_{CH} = \mathbf{r}_H - \mathbf{r}_C$, and $\mathbf{r}'_C = \mathbf{r}_C$.

tial, which is a function of the nuclear positions, can be expressed as a sum of potentials. Each of these potentials is related to a particular symmetry operator, and is a function of only those nuclear coordinates that break that particular symmetry operator[29, 30]. This effectively assumes that the motion in each of these symmetry directions is independent, permits us to decouple the symmetry-breaking and symmetry-conserving motions. The Hamiltonian for this system is therefore

$$\mathcal{H} = \frac{\mathbf{p}_C^\dagger \cdot \mathbf{p}_C}{2m_C} + \frac{\mathbf{p}_H^\dagger \cdot \mathbf{p}_H}{2m_H} + V_{\text{eff}}(\mathbf{r}_H, \mathbf{r}_C), \quad (1)$$

where $\mathbf{r}_C$ and $\mathbf{r}_H$ are the displacements of the carbon and hydrogen atoms from the centre of mass, m_C and m_H are the effective masses of the carbon and hydrogen that are determined by the concerted motion of the nuclei, and both being four times the value of the natural masses of the nuclei. Here V_{eff} is the effective potential determined by the adiabatically separated electrons, and the nuclear momenta operators defined in the usual way [21] with the canonical commutation relations $[\mathbf{p}_H \cdot \mathbf{e}_i, \mathbf{r}_H \cdot \mathbf{e}_j] = -i\hbar\delta_{ij}$, $[\mathbf{p}_C \cdot \mathbf{e}_i, \mathbf{r}_C \cdot \mathbf{e}_j] = -i\hbar\delta_{ij}$, where $\mathbf{e}_0$ and $\mathbf{e}_1$ are a pair of two-dimensional perpendicular unit vectors. However calculation in this basis is inconvenient because there are large regions of $\mathbf{r}_H$ which can be effectively ignored. Additionally a good basis choice is convenient for the interpretation of results, and so we choose a new basis for the calculation, shown in Fig. 2. This coordinate transformation results in a transformed set of canonical momenta that are related by $\mathbf{p}_C = \mathbf{p}'_C - \mathbf{p}'_{CH}$, and $\mathbf{p}_H = \mathbf{p}'_{CH}$. The Hamiltonian becomes

$$\mathcal{H} = \frac{\mathbf{p}'_C{}^\dagger \cdot \mathbf{p}'_C}{2m_C} - \frac{\mathbf{p}'_C{}^\dagger \cdot \mathbf{p}'_{CH} + \mathbf{p}'_{CH}{}^\dagger \cdot \mathbf{p}'_C}{2m_C} + \frac{m_C + m_H}{m_C m_H} \mathbf{p}'_{CH}{}^\dagger \cdot \mathbf{p}'_{CH} + V_{\text{eff}}. \quad (2)$$

This basis choice reflects the structure of the molecule, which is determined by V_{eff} . The potential is four dimensional, however because of the structure we are in practice able to subdivide the potential into three terms

that reflect the potential energy of the C-C ring or the C-H bond, and the cross terms between them. We use

$$V_{\text{eff}} = V_1(\mathbf{r}'_C) + V_2(\mathbf{r}'_{CH}) + V_3(\mathbf{r}'_C, \mathbf{r}'_{CH}), \quad (3)$$

$$V_1(\mathbf{r}'_C) = V_{\text{eff}}(\mathbf{r}'_C, \mathbf{R}_{CH}), \quad (4)$$

$$V_2(\mathbf{r}'_{CH}) = V_{\text{eff}}(\mathbf{R}_C, \mathbf{r}'_{CH}), \quad (5)$$

where the coordinates $\mathbf{R}_{CH}$ and $\mathbf{R}_C$ are the chosen such that $V_{\text{eff}}(\mathbf{R}_C, \mathbf{R}_{CH})$ is the global minimum of the potential. The potential $V_3(\mathbf{r}'_C, \mathbf{r}'_{CH})$ is the four-dimensional correction term that accounts for the coupled nature of the bonding. If we have chosen the basis well then $V_3(\mathbf{r}'_C, \mathbf{r}'_{CH})$ is small everywhere, and we can qualitatively characterize the molecule as a simple combination of C-C and C-H bond motions, in a form analogous to the harmonic approximation [29, 30].

The choice of this basis also enables us to impose appropriate boundary conditions on the system. A pair of potential cutoffs V_1^{max} , $V_{2,3}^{\text{max}}$ were chosen, so that in the calculation only regions where $V_1(\mathbf{r}'_C) < V_1^{\text{max}}$ and $V_2(\mathbf{r}'_{CH}) + V_3(\mathbf{r}'_C, \mathbf{r}'_{CH}) < V_{2,3}^{\text{max}}$ are included. A potential cut-off is appropriate as regions outside the boundary have too high a potential energy and so have a negligible amplitude contribution to the wavefunction and energy. We make two different choices of potential cutoff since the carbon and hydrogen nuclei have very different masses and different amounts of potential energy.

To perform the calculation, the coordinate space was divided into a Cartesian mesh. Values of the mesh spacing in each dimension, and the potential cutoffs V_1^{max} , $V_{2,3}^{\text{max}}$ were set so that the energy of the 2nd excitation from the ground state, which is the lowest energy IR vibration, was converged to one part in 30. This corresponded to 38000 points. This corresponded to a convergence of tunnelling separation energies to one part in 4, and due to the coarseness of the grid around the barrier, and the rapid change of the wavefunction in this region they are likely 20% smaller. The ratios between tunnelling separation energies were converged to one part in 40, so although the total separation energies are not well converged, their relative sizes are. This allows us to compare the strength of tunnelling between different vibration states. The eigenvalues were calculated using the dqds (differential quotient difference with shifts) algorithm [31], and the eigenvectors were calculated using Relatively Robust Representations (RRR) [32], as implemented in the LAPACK library [33].

The potential $V_{\text{eff}}(\mathbf{r}'_C, \mathbf{r}'_{CH})$ was calculated with density functional theory (DFT). This was performed using the plane-wave code CASTEP [34]. As plane-wave code relies on a periodic basis set, the size of the unit cell and the cutoff energy of the plane waves was converged, to where the error in the barrier height energy was less than 0.9 meV. The exchange correlation correction was calculated using the LDA [35].

FIG. 3. The potential $V_1(\mathbf{r}'_C)$, shown for a constant l_C chosen such that the potential energy at $\mathbf{r}'_C = (0, l_C)$ is minimized. Potentials calculated using LDA here ($l_C = 2.02$), and reduced multireference coupled-cluster method with singles and doubles (RMRCCSD) calculations ($l_C = 2.07$) taken from reference [27], are compared.

RESULTS

DFT calculations found that when the potential is separated into V_1 , V_2 and V_3 terms, the range of V_3 never exceeded 10 percent of the energetic variation from the global potential minimum, and there was no discernible change in the position of the minimum of $V_2(\mathbf{r}'_{CH}) + V_3(\mathbf{r}'_C, \mathbf{r}'_{CH})$. This will permit us to qualitatively interpret the results as linearly linked C-H and C-C bond motions. We find the V_1 potential, for which a cross-section is shown in Fig. 3, to be very similar to the standard 1-D double well system in which the first few energy levels are well localized. As their energy is less than the barrier height, their wavefunctions tunnel through the central barrier. The characteristic features of localized states in the double well, is that they are found in closely energetically spaced symmetric/antisymmetric pairs, and that the tunnelling rate across the barrier is proportional to this energy separation of the pairs. The more localized these states are, the smaller the energy separation between these pairs is, and correspondingly the smaller the rate of tunnelling. We performed computations on the two-dimensional potentials as well as the four-dimensional potential, to compare how motion of the hydrogen nucleus affects the tunnelling of the carbon ring.

Restricted 2D calculations

In order to understand how the multidimensional nature of the system affects the tunnelling rate, we performed calculations on two 2D subsystems in addition to the full 4D case. The 2D calculations were performed where either $V_2 + V_3 \equiv 0$ (called ring-only), or $V_1 + V_3 \equiv 0$ (called C-H-only). The results for the C-H-only calculation are straightforward as unlike V_1 there is only one potential well that is predominantly harmonic. The V_2 potential and wavefunctions are shown in Fig.(4). There

FIG. 4. a) The governing potential $V_2(\mathbf{r}_{CH})$. b), c), d) Wavefunctions of the principal excited states with fixed $\mathbf{r}_C$ so that V_1 is at a minimum. ψ is the unitless finite (and real) wavefunction and defined so that the sum over the points (located at line intersections) $\sum \psi^2 = 1$.

FIG. 5. The governing potential $V_1(\mathbf{r}'_C)$; the energies of the potential are in eV. The jagged edge at 1.4 eV chosen as the boundary of DFT calculation, above the 1.13 eV cutoff.

Ring Only			
state	D_{2h}	energy (cm^{-1})	TSE (cm^{-1})
0 & 1	ground	0	0.0008
2 & 3	$1A_g$	1067	0.112
4 & 5	$2A_g$	1611	0.112

C-H only		
state	D_{2h}	energy (cm^{-1})
0		0
1	$1A_g$	1248
2	$1A_g^2$	
3	$1A_g^3$	
4	$2A_g$	3139

TABLE I. Energies of eigenstates calculated under ring-only and C-H only constraints. States are labelled by energy hierarchy and rectangular symmetry considerations (D_{2h}). For the ring-only case the energy separation between symmetric and antisymmetric states is also shown, labelled as Tun. Sep.

FIG. 6. a)-f) Real wavefunctions of the finite 2-dimensional system with fixed $\mathbf{r}_{CH}$ so that $V_2 = 0.0$, with the first three (rectangular) states and their symmetric/antisymmetric pair. Values of ψ are normalized over the sum of their squares, all coordinates l_C , d_C are in Angstroms. a), b) correspond to the ground state under rectangular considerations, c), d) to the long-bond excitation, and e), f) to the short-bond excitation.

are two principal excitations, which we define as those states in which there is only one node in the wavefunction, as these correspond to the experimentally observable Raman excitations from the ground state. These states will dominate the spectrum since the wavelength of infrared radiation is much longer than the size of the molecule, so the gradient of a resonant electric field is roughly constant, and states with multiple wavefunction nodes will generate a smaller response. These principal excitations are fairly conventional with both the waggle mode where the wavefunction node is aligned along l_{CH} shown in Fig.(4c), and the stretch mode in with the wavefunction mode aligned along d_{CH} shown in Fig. (4b). The energies in table I show that the waggle mode has a lower energy than the stretch mode, due to the shape of the potential as the C-H bond is stiffer with respect to length changes than to lateral changes.

The V_1 potential used for the ring only calculation is shown in Fig.(5). The shape of the potential is analogous to a two dimensional version of the double well potential as shown in the cross-section shown in Fig.(3). Compared to the coupled-cluster calculations, the LDA potential used here has a similar barrier shape and height, but has wider wells [27]. Comparatively this will lead to a reduced localization of the wavefunction and weaker

FIG. 7. Pairwise average of the wavefunction densities of the first three principal rectangular excitations projected onto a)-c) the ring coordinates, d)-f) hydrogenic coordinates, and the fourth principal rectangular excitation projected onto g) ring coordinates, h) hydrogenic coordinates. Coordinate distances are in Å, and the densities normalized over the grid.

tunnelling than if a coupled-cluster potential was used. The contours around the two wells of the V_1 potential in Fig.5 resemble a pair of egg-shaped ovals, and the effects of the single and double electronic bonds are visible. The long direction of each oval is aligned with the single bond, and the short width of the oval is aligned with the double bond. The wavefunctions for the lowest six energy states shown in Fig.(6) are the principal excitation states. These need to be defined slightly differently to the C-H bond case since there are no calculations in which there is a single continuous wavefunction node, as it continues past the potential cutoff in the calculation. We define the principal excitation states here as the states in which there is only one wavefunction node in each potential well. Due to the anharmonicity of the potential the existence of multiple nodes does not suppress the Raman response. This is because in anharmonic systems under the per-

turbation of a linear spatial potential the ground state can transition to a multiple noded wavefunction, as the sum of the raising and lowering operators $\hat{a}^\dagger + \hat{a}$ is not proportional to the position operator $\hat{x}$. The wavefunctions show that the single and double bonds give rise to these principal excitations, with those corresponding to the length-excitation with the wavefunction node across the short axis of the oval (Fig. 6c,d)), and those corresponding to the width excitation with the wavefunction node along the long axis (Fig.6 e),f)).

The six lowest-energy states are shown in Table I. They are found in symmetric/antisymmetric pairs and are separated by a small energy which we call the tunnelling separation energy (labelled TSE). Within each pair the wavefunction density, the square of the wavefunction, is very similar and differs significantly only in the tunnelling region between the two wells. By contrast in this re-

States	D_{2h} label	energy (cm^{-1})	TSE (cm^{-1})
0&1	ground	0	0.025
2&3*	$1A_g$	839	0.074
4&5*	$2A_g$	1005	0.046
6&7*	$3A_g$	1481	0.26
8&9	$1A_g^2$	1661	0.13
10&11	$1A_g \times 2A_{1g}$	1811	0.14
12&13	$2A_g^2$	1980	0.057
...			
...			
28&29*	$4A_g(+)$	3073	0.13

TABLE II. Energies and the tunnelling separation energy between symmetric/antisymmetric pairs (TSE), of eigenstates ranked by energy and classified with D_{2h} symmetry considerations in the full 4-dimensional calculation. Principal excitations that we expect to dominate the Raman spectrum are marked with a star. A large number of states are not presented. Since they are not principal excitations they are unconverged. Due to this the state numbers of the final two excitations is likely incorrect.

FIG. 8. A_g excitations in cyclobutadiene

gion the wavefunction phase either stays the same (symmetric states) or changes sign (antisymmetric states). In each pair the symmetric state is of lower energy than the antisymmetric state, in the low energy states that we calculated. By analogy to the double well, the differences in energy between each state in the pair gives the rate of tunnelling for those states. These energy differences, shown in table I, determine the tunnelling rate. The greater the energy difference between the pair, the higher the tunnelling rate. This is because a localized state constructed from a superposition of the symmet-

ric/antisymmetric states will oscillate between the wells at a frequency $f = \Delta E/\hbar$, which increases as energy separation ΔE . As expected the pair splitting increases with respect to the total state energy, but this relationship is not linear; the length and width excitations both have around the same tunnelling separation energies despite the difference in the total energies. This is because the motion and the momentum in the short-bond is more tangentially aligned to the barrier, and so the wavefunction does not penetrate so far through it.

To summarise, in the two-dimensional calculations we find that in the C-H-only calculations the bond motion is very similar to the archetypal C-H bond, with a high energy stretch mode of energy around 3100 cm^{-1} , and a low energy waggle mode[30]. In the ring-only calculations we find that the potential reflects the long and short bonds determined by the electrons, and these two bonds are responsible for the two different vibrational excitation energies. The system behaves similarly to a 1-D double well as states are found in symmetric/antisymmetric pairs, but the energy of a vibrational state is not enough to determine the tunnelling, as the distribution of the momentum is also important.

4D calculations

When we consider the hydrogenic and carbon-ring motions together, there are four principal excitations possible from D_{2h} symmetry considerations, and the ground state [36]. The most important effect of coupling the hydrogen and ring systems together is a mixing of the motions. As a result the hydrogen and carbon motions become correlated and illustrative sketches of these correlated motions are shown in Fig. (8).

Of these the ring motions and the C-H waggle mix the most strongly, and form three different states. The C-H stretch motion only weakly couples with the ring motion, and as a result the energy is very similar to the C-H stretch calculated in the two-dimensional case. This can be seen from the projected densities, defined as $\rho_{\text{CH}}(\mathbf{r}_{\text{CH}}) = \sum_{\mathbf{r}_{\text{C}}} \psi(\mathbf{r}_{\text{C}}, \mathbf{r}_{\text{CH}})^2$, and $\rho_{\text{C}}(\mathbf{r}_{\text{C}}) = \sum_{\mathbf{r}_{\text{CH}}} \psi(\mathbf{r}_{\text{C}}, \mathbf{r}_{\text{CH}})^2$, shown in Fig.(7); since the wavefunctions' nodes do not pass through the four-dimensional space perpendicular to either the hydrogenic coordinates or the ring coordinates, there are no nodes where the density falls to zero in any of the density projections. Most importantly the density is still split into two regions on either side of the central barrier and this corresponds to a small energy separation between the symmetric and antisymmetric pair. This means that along with the correlations between hydrogenic and carbon-ring motion, the excited states are well localized into the rectangular states.

The first two excited states, $1A_g$ and $2A_g$ shown in Figs. (7 b,e) and c),f)), are the result of strong mixing between the C-H waggle and the long-bond excitation in the ring. By comparison, the $1A_g$ excitation is broader and mixes more strongly. This can be observed since

FIG. 9. The expectation function $\langle d_{CH}(\mathbf{r}_C) \rangle$, which shows how the hydrogen and carbon motions are correlated. The values of all axes are in Angstroms.

while in both excitations there is a saddle point between the two density maxima, which are present in both H and ring motion. In the $1A_g$ excitation the density at the saddle point is more similar to the density at the maxima maxima than in the $2A_g$ excitation. This is likely because the carbon and hydrogen nuclear movements are mixed, and must share the limited energy available in the state. This means they are constrained to remain near the minima in the potential. The $3A_g$ excitation is mostly comprised of the short bond excitation in the ring, and is accompanied by a much smaller amount of hydrogen motion than for the other excitations, as the projected density for the hydrogen is very similar to that of the ground state.

However the projected densities do not provide important information on correlations of the nuclear motion in the excited states. While they show how much the hydrogenic and carbon motions have combined, they do not show how the motions are correlated. This information is required to match the states with schematic motions in Fig. (8). It is given by the expectation function

$$\langle d_{CH}(\mathbf{r}_C) \rangle = \frac{\sum_{\mathbf{r}_{CH}} d_{CH} \psi(\mathbf{r}_{CH}, \mathbf{r}_C)^2}{\sum_{\mathbf{r}_{CH}} \psi(\mathbf{r}_{CH}, \mathbf{r}_C)^2}, \quad (6)$$

which calculates the expected position of the hydrogen for a choice of ring coordinates. This is shown in Fig. (9). Using this we are able to obtain the correlated motions of the atoms. In the $1A_g$ excitation, when the long bond is stretched, the hydrogen waggles in the same direction. The two are positively related, and so the hydrogen waggle and long-bond stretch are in phase. The $2A_g$ is the reverse situation, and so the motions are instead out of phase. In the $3A_g$ excitation, there is less movement overall in the C-H waggle, but the short-bond excitation and the waggle motion are also out of phase. In the $2A_g$ excitation there is a sharp change in the hydrogen displacement at $d_C = 0$ accompanied by a sign change in $\langle d_{CH} \rangle$, this is possible without the energetic penalty as-

sociated with a rapid change in the wavefunction because the wavefunction amplitude here is so low.

These calculations are able to calculate the energies and wavefunctions of nuclear energy states, which can be used to calculate all the properties of the excited states. However to perform this calculation effectively, and in order to interpret the results, the coordinate system used needed to reflect the structure of the potential energy surface. Since the potential energy surface is determined by the electrons we are able to use the bond lengths and angles (or the inter-atomic distances) as this basis. It is also important to restrict the space of the calculation, and this can be done by an appeal to the structure of the potential energy surface. It is important to ensure that the curvature of the wavefunctions, determined by the width of the well and the effective mass of the system in the chosen coordinates, is captured effectively. While this is done by choosing appropriate potential energy cutoffs and a choice of mesh grid spacing, the unequal curvature of the wavefunctions in Figs.(946) suggests that improvements could be made which would further reduce the size and complexity of the computation.

In the ring-only vibrations (detailed in Table I), the inclusion of the hydrogenic motion (detailed in Table II) suppresses tunnelling motion in long-bond excitation states, but enhances it in the short-bond excitation states. This is in contrast to the two-dimensional results in which the tunnelling rate is equal for both single and double-bond stretches. There are two causes for difference in the tunnelling rate as more degrees of freedom are considered: in the $1A_g$ excitation the combination of hydrogen motion with the long bond stretch lowers the energy of the vibration from 1067 cm^{-1} to 839 cm^{-1} , this lower energy results in a reduction of the nuclear momentum and the tunnelling energy difference from 0.112 cm^{-1} to 0.074 cm^{-1} . The second is caused by correlations between the carbon and hydrogen nuclei that suppress the tunnelling rate. For the $3A_g$ out-of-phase waggle state

shown in Fig. 9 b), as the ring conformation approaches $d_C = 0$, $\langle d_{CH} \rangle$ remains high, and at $d_C = 0$ there is a sharp change in $\langle d_{CH} \rangle$. This means that when the ring is in a conformation that is conducive to tunnelling, the hydrogen is out of place, and so the tunnelling rate is suppressed for this vibrational mode. Based on this analysis, the deuterated molecule would due to an increased mass have smaller transition frequencies and lower tunnelling rates for all the states, and we find this to be the case in related calculations.

SYMMETRY AND RAMAN EXCITATIONS

Point group symmetries are used to classify and assign the eigenstates of nuclear motion [29, 30], however in cyclobutadiene the symmetry of the molecule is ambiguous. This is because there are two equivalent minima in the potential energy surface, at nuclear configurations that correspond to a D_{2h} symmetry, but tunnelling means that the eigenstates of the system are superpositions of these two configurations with a D_{4h} symmetry. Additionally the typical energies of the tunnelling separation are small enough that they may be distorted by an external perturbation, so we give an account for how this can occur as well.

In harmonic systems only excitations from the ground state to eigenstates with single nodes (principal transitions) are easily observable. Point group symmetries are useful for such systems as they allow us to predict the number of, and symmetry of, these principle transitions [36]. When we treat cyclobutadiene as a molecule with D_{2h} symmetry, this theory predicts that our calculations will accommodate four principle transitions of A_g symmetry. When we treat it with D_{4h} symmetry, it is predicted that there are two principle transitions with a A_{1g} symmetry and two with B_{2g} symmetry. This conflicts with our results, which show ten eigenstates arranged in pairs of similar energy, corresponding to up to nine principle transitions. The discrepancy between these two models is due to the anharmonicity of the potential. While D_{4h} is a higher symmetry, the lowest energy nuclear configuration of D_{4h} symmetry is in fact a saddle point rather than a minimum so the requirement for this type of treatment, that the potential is harmonic, is not satisfied.

Since this anharmonicity is most significant in the V_2 potential, it is instructive to re-examine the 2-D wavefunctions of Fig. 6. All of these states satisfy D_{4h} symmetry: symmetric states correspond to A_{1g} motions, and the antisymmetric states to B_{2g} motions. However most of the wavefunction is localised around the energetic minima located at configurations of D_{2h} symmetry, and around which the potential is almost harmonic. From the energies in table I the effective width of the potential barrier at the D_{4h} symmetry point is high enough that the tunnelling separation energies are much less than the energies of the lowest energy eigenstates. This means that for en-

ergetic purposes superpositions of configurations around the two minima in the potential are well separated and only interact weakly, and the coarse structure of the spectrum (in which the symmetric/antisymmetric pairs are treated as single eigenstates) can be readily interpreted using a D_{2h} symmetry approach, that relies on harmonicity [26, 27]. The overall D_{4h} symmetry is therefore only relevant for an experiment sensitive to the tunnelling separation energy, or for high-energy excitations that can tunnel through the barrier more easily. This is similar to the inversion doubling of spectral peaks, for example in ammonia [37, 38].

Anharmonicity and symmetry is of further relevance to calculations of cyclobutadiene. In particular we made the assumption that the different symmetry breaking and conserving motions could be decoupled. However recent calculations [39, 40] found metastable configurations of similar energy to the ground state, including a puckered configuration with an energy higher than the ground state by 302cm^{-1} , and a distorted planar ring configuration like that observed in the related tetrasilyl-substituted molecule with out-of-plane C-H bonds, that was higher in energy by 533cm^{-1} . While these metastable configurations are different enough from the D_{2h} and D_{4h} configurations that they are unlikely to contribute to tunnelling, they show that the full potential energy surface of cyclobutadiene is complex and is not harmonic with respect to deviations that break the same symmetry. This suggests that motions which break different symmetries are not completely separable.

A comparison of the vibrational frequencies between theory and experiment supports this hypothesis. As is to be expected from Fig. 3, the DFT potential well is broader and shallower than more accurate potentials used elsewhere, and this leads to an underprediction of vibrational frequencies for all excitations from the experimental values[28]. Those other theoretical studies on cyclobutadiene that focus on more accurate calculations of the potential function, also separate the different symmetry motions, and show only a partial improvement over the work performed here[15, 16, 27]. This suggests that most of the remaining improvements to be made in the **determination** of vibrational frequencies will require us to lose the assumption that motions which break different symmetries are separable, as discussed above.

Raman excitations

Due to the tunnelling and the anharmonicity, it is not immediately clear how the intensities of infra-red transitions in cyclobutadiene are to be calculated. As such we will examine how Raman scattering can be used to probe transitions between energy levels. Transitions between nuclear vibrational energy levels will correspond directly to Raman resonances[29, 30], due to the symmetry in cyclobutadiene. The amplitude of an optical-induced transition is dependent on a transition due to the polar-

FIG. 10. a) the eight regions ($L_t^\pm, S_s^\pm$), that can be used to determine Raman excitations to the principal excited states, and maps of the sign of the phase in b) the symmetric $1A_g$ excitation and c) the antisymmetric $1A_g$ excitation

izability operator[41]

$$T_{a \rightarrow b} = \langle \psi_b | \hat{P} | \psi_a \rangle \quad (7)$$

where the transition is from nuclear vibration state a to state b , ψ_a is the wavefunction of state a , and the polarizability operator for the chosen field direction ϵ is given by $\hat{P}_\epsilon = \int d\mathbf{x}_n |\mathbf{x}_n\rangle \langle \mathbf{x}_n| P_\epsilon(\mathbf{x}_n)$, where the sum is over the permitted nuclear configurations $\mathbf{x}_n$ and $P_\epsilon(\mathbf{x}_n)$ is the polarizability of the electrons for the nuclear configuration $\mathbf{x}_n$. The polarizability operator is further defined as a projection of the polarizability tensor into the field direction ϵ .

The most straightforward application of this theory to simple hydrocarbons makes use of an approximation, that the polarizability tensor changes linearly with respect to changes in the molecular configuration [42, 43]. This means that in harmonic systems the transition amplitude is dependent only on the first derivatives of the polarizability tensor. Moreover because of the linear response, the overall polarizability of the molecule can be expressed as a sum of polarizability contributions from each individual bond. However since the polarizability is already a second-order perturbative property, even for harmonic systems its derivative can in many cases not be approximated as constant. Due to this and the anharmonicity, this method therefore needs to be modified in cyclobutadiene to calculate the amplitudes of Raman scattering.

We will compare the transitions from the ground state (state 0) to the two $1A_g$ states (states 2 and 3), to investigate the interaction of the Raman spectrum with the tunnelling. First we must set up the basis that we will study. Motivated by previous observations that most of the wavefunction is localised near the two minima with D_{2h} symmetry, we separate the configuration space into two, these are shown by the red ovals in Fig.10. It is necessary to use the wavefunctions of the excited vibrational states, and the only significant difference between these

two wavefunctions is in their phase dependence on $\mathbf{r}_C$. The phases for the excited states are shown for our two calculation regions in Fig.10 b) and c). For the electric field, we take its orientation to be along the direction of one of the C-C bonds, as this breaks the D_{4h} symmetry in a way that permits transitions from symmetric A_{1g} to antisymmetric B_{2g} states. Finally we need the polarizability responses of the bonds to this electric field. We label the responses for C-C bonds using $T_t^\pm$, where T has a value of either L for long bonds or S for short bonds, t describes the orientation of the electric field ϵ to the bond (either parallel or perpendicular), and $\pm$ describes the configuration of the bond which is either stretched (+) or compressed (-). With this labelling we are able to distinguish eight distinct sub-regions that contribute to the polarizability calculation, these are shown in Fig.10 a).

With this basis set we are able to calculate the transition amplitude in terms of these sub-regions of the configuration space. If we define the sub-region polarizability as

$$P(L_t^\pm, S_s^\pm) = \int_{(L_t^\pm, S_s^\pm)} d\mathbf{r}_C \int d\mathbf{r}_{CH} \langle \psi_3 | \mathbf{r}_C, \mathbf{r}_{CH} \rangle \hat{P}(\mathbf{r}_C, \mathbf{r}_{CH}) \langle \mathbf{r}_C, \mathbf{r}_{CH} | \psi_0 \rangle, \quad (8)$$

where the integral in $\mathbf{r}_C$ is over the limits of the sub region ($L_t^\pm, S_s^\pm$). We are then able to write the transition amplitude as a sum of the contributions of each sub-region

$$T_{0 \rightarrow 3} \approx \left(P(L_{\parallel}^+, S_{\perp}^-) + P(L_{\parallel}^+, S_{\perp}^+) \right) - \left(P(L_{\parallel}^-, S_{\perp}^-) + P(L_{\parallel}^-, S_{\perp}^+) \right) - \left(P(L_{\perp}^+, S_{\parallel}^-) + P(L_{\perp}^+, S_{\parallel}^+) \right) + \left(P(L_{\perp}^-, S_{\parallel}^-) + P(L_{\perp}^-, S_{\parallel}^+) \right). \quad (9)$$

This sum is approximate as we have not included the contribution from areas around the anharmonic central barrier, where the wavefunction amplitude is very small. Previous work calculated $T_{0 \rightarrow 3}$ to be approximately zero [3], but that $T_{0 \rightarrow 2}$ was non-zero. From this we can infer that $P(L_t^+, S_s^\pm) - P(L_t^-, S_s^\pm)$ is non-zero, as the $0 \rightarrow 2$ transition is non-zero, so for $T_{0 \rightarrow 3}$ the contributions from each side of the barrier cancel out. Simply put, the derivative of the polarizability is symmetric with respect to a reflection about $d_C = 0$. This is potentially unexpected, as it means that the polarizability response when the long bonds are aligned parallel to the electric field, is the same as when the short bonds are aligned parallel. What can be interpreted from this is that the long and short bonds are not separable because the polarizability must be dependent on the conformation of the whole ring. This puts cyclobutadiene in the same category as long alkene chains, in which the polarizability is dependent on the interactions between bonds, and cannot be isolated in individual bonds or sub-systems[44].

The energy differences between the symmetric and antisymmetric states are about 5 meV, as a result we expect these properties of the system to be sensitive with respect to interactions with the surrounding environment. To analyse how thermal and symmetry-breaking external potentials have an effect on the system, we divide the possible effects on the Raman spectrum of a symmetry-breaking external potential into three different classes when there is a i) negligible, ii) weak, or iii) large external potential, compared to the energy splitting caused by tunnelling.

When there is no external potential, there will be no localization into rectangular states. Optical excitations from symmetric to antisymmetric wavefunctions cannot be observed, and this means that the energy splitting caused by tunnelling is not observable. However if the temperature is high enough ($k_B T \approx$ tunnelling separation energy), then there is a thermal occupation of the antisymmetric pair of the ground state, and the Raman permitted transitions between antisymmetric pairs will be observable as excitations from the antisymmetric state to higher energy antisymmetric states are possible.

When the external potential is slightly larger than the lowest symmetric/antisymmetric pair energy difference, only those lowest energy states will be localized into a specific rectangular state, as these have the lowest nuclear momentum and the weakest tunnelling, and a small energetic perturbation will localize them. To first order the wavefunctions of these low energy states will be approximately, in terms of the free-state wavefunctions $\psi_{\pm} = \psi_0 \pm \psi_1$. The higher energy states will retain most of their tunnelling characteristics. The ground state will completely lose its square symmetry. This localisation means that the transition amplitude $T_{0 \rightarrow 3}$ changes, and instead is

$$T_{0 \rightarrow 3} \approx \sqrt{2} \left(P(L_{\parallel}^{+}, S_{\perp}^{-}) + P(L_{\parallel}^{+}, S_{\perp}^{+}) \right) - \sqrt{2} \left(P(L_{\parallel}^{-}, S_{\perp}^{-}) + P(L_{\parallel}^{-}, S_{\perp}^{+}) \right) \quad (10)$$

and excitations to all the high energy states will be observable regardless of the temperature. This enhancement is reminiscent of catalysis, in which a metastable binding to another medium enables transitions to new states without the need for an increase in temperature.

When the external potential is larger than the highest symmetric/antisymmetric pair energy difference, all the energy states accessible from a ground-state transition will be localized. There will be very little tunnelling across the barrier in any of the states. Due to the localization each state has a rectangular configuration and since all the states will be strongly localised, optical excitations between the two rectangular configurations will be strongly suppressed. Additionally at temperatures high enough to cause transitions between the two low energy rectangular states, tunnelling effects will not be found since tunnelling is suppressed for all states. Instead, depending on the exact form of the external potential, there

may be vibrational differences between states in which either the long or short bonds are aligned parallel to the gradient of the potential.

CONCLUSION

In general, tunnelling is important in many chemical reactions [14, 45, 46] and structural phenomena. While tunnelling in one dimension is easy to calculate, the extension to a multi-dimensional system like a coupled molecule represents a significant challenge [47]. A method of reducing this complexity is by choosing a suitable tunnelling pathway [45, 48]. In cyclobutadiene, as in ammonia [38], the tunnelling of particular states is sensitive to the distribution of momentum. The effect of this is that particular vibration states tunnel through the barrier along different paths, and so the calculation space must include these different tunnelling routes. The rates of tunnelling reactions that proceed via an adiabatic pathway are therefore dependent on eigenstates of the nuclear motion and their momentum distribution. We have shown that in some cases these can be understood as anharmonic extensions of the well understood harmonic resonance approximation, that is used to classify the resonances [29, 30]. This means that the nuclear motions under the well known harmonic approximation can be used to qualitatively understand the rates and pathways of tunnelling reactions, despite the multidimensional complexity of the problem.

Calculations performed here show that the square symmetry of cyclobutadiene is due to quantum tunnelling, comprised of correlated nuclear motion through an energetic barrier. This tunnelling leads to a small correction in the energies of the vibrational frequencies of the molecule calculated under an assumed rectangular symmetry. Each energy level separates into a symmetric and an antisymmetric pair, and by analogy to the 1D double well potential, the size of the energy separation between these symmetric and antisymmetric pairs is related to the tunnelling rate across the barrier. From a comparison of the different vibrational states we find that the tunnelling rate across the barrier is sensitive to the distribution of momentum in the molecule, and the correlations between the motion of the nuclei. Our expectations about the motion of the molecule, for example tunnelling suppression in the out-of-phase waggle state, is able to give an accurate account for the strength of the tunnelling process. We also find that despite the overall square symmetry a combination of square and rectangular labels is most appropriate to classify the dominant system excitations and the Raman spectrum. This is because the square potential is so anharmonic that the usual assumption, that at the symmetry point there is an energetic minimum, fails. Instead the global energetic minima are at points that have rectangular symmetry. The tunnelling effects are sensitive to an external potential, and a small external interaction can enable transitions to states that would otherwise be forbidden without a thermal excitation, while

a large external interaction will again suppress that transition. This process is reminiscent of catalysis activation, and over-binding where a catalyst suppresses a reaction by trapping the reactants in a bound state. We have demonstrated that this method of calculating wavefunctions and energies can incorporate and enable the analysis of many complex quantum mechanical phenomena under a single approach. Perhaps more importantly, it can successfully inform our qualitative understanding and intuition about molecular motion as it generates quantitative results.

- [1] B. K. Carpenter, Journal of the American Chemical Society **105**, 1700 (1983).
- [2] R. L. Redington, The Journal of Chemical Physics **109**, 10781 (1998).
- [3] P. Čársky and J. Michl, Theoretica chimica acta **84**, 125 (1992).
- [4] D. W. Whitman and B. K. Carpenter, Journal of the American Chemical Society **104**, 6473 (1982).
- [5] F. Fantuzzi, T. M. Cardozo, and M. A. C. Nascimento, ChemPhysChem **17**, 288 (2016).
- [6] G. Maier, H.-G. Hartan, and T. Sayrac, Angewandte Chemie International Edition in English **15**, 226 (1976).
- [7] R. Breslow, R. Grubbs, and S. Murahashi, Journal of the American Chemical Society **92**, 4139 (1970).
- [8] V. Gogonea, P. v. R. Schleyer, and P. R. Schreiner, Angewandte Chemie International Edition **37**, 1945 (1998).
- [9] T. Bally, Angewandte Chemie International Edition **45**, 6616 (2006).
- [10] A. Fattahi, L. Lis, Z. Tian, and S. R. Kass, Angewandte Chemie International Edition **45**, 4984 (2006).
- [11] A. M. Orendt *et al.*, Journal of the American Chemical Society **110**, 2648 (1988).
- [12] B. Kovacevic, D. Bari, Z. B. Maksi, and T. Miller, The Journal of Physical Chemistry A **108**, 9126 (2004).
- [13] T. Saito *et al.*, Chemical Physics Letters **498**, 253 (2010).
- [14] D. Ley, D. Gerbig, and P. R. Schreiner, Org. Biomol. Chem. **10**, 3781 (2012).
- [15] A. Balkov and R. J. Bartlett, The Journal of Chemical Physics **101**, 8972 (1994).
- [16] M. Eckert-Maksi, M. Vazdar, M. Barbatti, H. Lischka, and Z. B. Maksi, The Journal of Chemical Physics **125**, 064310 (2006).
- [17] A. Kostenko *et al.*, Angewandte Chemie International Edition **56**, 10183 (2017).
- [18] Y. Shiota, M. Kondo, and K. Yoshizawa, The Journal of Chemical Physics **115**, 9243 (2001), <https://doi.org/10.1063/1.1412250>.
- [19] P. Čársky, R. J. Bartlett, G. Fitzgerald, J. Noga, and V. Špirko, The Journal of Chemical Physics **89**, 3008 (1988).
- [20] R. Lefebvre and N. Moiseyev, Journal of the American Chemical Society **112**, 5052 (1990).
- [21] J. Binney and D. Skinner, *The Physics of Quantum Mechanics: An Introduction* (Cappella Archive, 2010).
- [22] W. T. Borden, Wiley Interdisciplinary Reviews: Computational Molecular Science **6**, 20 (2016).
- [23] S. Coleman, *Aspects of Symmetry: Selected Erice Lectures* (Cambridge University Press, 1985).
- [24] S. R. McConnell, A. Lhle, and J. Kstner, The Journal of Chemical Physics **146**, 074105 (2017).
- [25] M. T. Cvitas and S. C. Althorpe, Journal of Chemical Theory and Computation **12**, 787 (2016).
- [26] P. Čársky, V. Špirko, B. A. H. Jr., and L. J. Schaad, The Journal of Chemical Physics **92**, 6069 (1990).
- [27] X. Li and J. Paldus, The Journal of Chemical Physics **131**, 114103 (2009).
- [28] B. R. Arnold, J. G. Radziszewski, A. Campion, S. S. Perry, and J. Michl, Journal of the American Chemical Society **113**, 692 (1991).
- [29] E. Wilson, J. Decius, and P. Cross, *Molecular Vibrations: The Theory of Infrared and Raman Vibrational Spectra* (Dover Books on Chemistry Series (Dover Publications, 1955)).
- [30] G. Herzberg, *Molecular spectra and molecular structure. Vol.2: Infrared and Raman spectra of polyatomic molecules* (D. Van Nostrand Company Inc., 1945).
- [31] K. V. Fernando and B. N. Parlett, Numerische Mathematik **67**, 191 (1994).
- [32] B. N. Parlett and I. S. Dhillon, Linear Algebra and its Applications **309**, 121 (2000).
- [33] E. Anderson *et al.*, *LAPACK Users' Guide*, Third ed. (Society for Industrial and Applied Mathematics, Philadelphia, PA, 1999).
- [34] S. J. Clark *et al.*, Z. Kristall. **220**, 567 (2005).
- [35] W. Kohn and L. J. Sham, Phys. Rev. **140**, A1133 (1965).
- [36] E. Silberman and H. W. Morgan, *Use of group theory in the interpretation of infrared and Raman spectra* (Oak Ridge National Laboratory, TN, 1977).
- [37] C.-K. Lin, H.-C. Chang, and S. H. Lin, The Journal of Physical Chemistry A **111**, 9347 (2007), PMID: 17725334.
- [38] A. G. Csaszar and T. Furtenbacher, Phys. Chem. Chem. Phys. **18**, 1092 (2016).
- [39] H. Xu, S. Saebo, and C. U. Pittman, Structural Chemistry **25**, 635 (2014).
- [40] Y. Apeloig, Science **331**, 1277 (2011).
- [41] M. Born and K. Huang, *Dynamical theory of crystal lattices* (Oxford: Clarendon Press, Oxford, 1954).
- [42] L. D. A., Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences **217**, 203 (1953).
- [43] K. Gough, The Journal of chemical physics **91**, 2424 (1989).
- [44] H. S. Smal, P.-O. strand, and L. Jensen, The Journal of Chemical Physics **131**, 044101 (2009).
- [45] W. T. Borden, Wiley Interdisciplinary Reviews: Computational Molecular Science **6**, 20 (2016).
- [46] P. R. Schreiner, Journal of the American Chemical Society **139**, 15276 (2017).
- [47] V. Benderskii, D. Makarov, and C. Wight, *Chemical Dynamics at Low Temperatures* (Advances in Chemical Physics (Wiley, 1994)).
- [48] R. P. Bell, *Experimental evidence for tunnelling in chemical reactions* (Springer US, Boston, MA, 1980), pp. 106–144.

y x

a) r_{CH} potential (in eV)

b) Ground state

c) 1st excited state

d) 4th excited state

a) Symmetric ground state

b) Antisymmetric ground state

c) Symmetric 1st state

d) Antisymmetric 1st state

e) Symmetric 2nd state

f) Antisymmetric 2nd state

a) $1A_g$ excitation

b) $2A_g$ excitation

c) $3A_g$ excitation

d) $4A_g$ excitation

a) $1A_g$ excitation b) $2A_g$ excitation c) $3A_g$ excitation

b)

c)