

Durham Research Online

Deposited in DRO:

22 February 2019

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Galby, E. and Lima, P.T. and Paulusma, D. and Ries, B. (2019) 'Classifying k-edge colouring for H-free graphs.', *Information processing letters.*, 146 . pp. 39-43.

Further information on publisher's website:

<https://doi.org/10.1016/j.ipl.2019.02.006>

Publisher's copyright statement:

© 2019 This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Classifying k -Edge Colouring for H -free Graphs^{*}

Esther Galby¹, Paloma T. Lima², Daniël Paulusma³, and Bernard Ries¹

¹ Department of Informatics, University of Fribourg, Switzerland, {esther.galby,bernard.ries}@unifr.ch

² Department of Informatics, University of Bergen, Norway, paloma.lima@uib.no

³ Department of Computer Science, Durham University, UK, daniel.paulusma@durham.ac.uk

Abstract. A graph is H -free if it does not contain an induced subgraph isomorphic to H . For every integer k and every graph H , we determine the computational complexity of k -EDGE COLOURING for H -free graphs.

1 Introduction

A graph $G = (V, E)$ is k -edge colourable for some integer k if there exists a mapping $c : E \rightarrow \{1, \dots, k\}$ such that $c(e) \neq c(f)$ for any two edges e and f of G that have a common end-vertex. The *chromatic index* of G is the smallest integer k such that G is k -edge colourable. Vizing proved the following classical result.

Theorem 1 ([27]). *The chromatic index of a graph G with maximum degree Δ is either Δ or $\Delta + 1$.*

The EDGE COLOURING problem is to decide if a given graph G is k -edge colourable for some given integer k . Note that (G, k) is a yes-instance if G has maximum degree at most $k - 1$ by Theorem 1 and that (G, k) is a no-instance if G has maximum degree at least $k + 1$. If k is *fixed*, that is, k is not part of the input, then we denote the problem by k -EDGE COLOURING. It is trivial to solve this problem for $k = 2$. However, the problem is NP-complete if $k \geq 3$, as shown by Holyer for $k = 3$ and by Leven and Galil for $k \geq 4$.

Theorem 2 ([14, 20]). *For $k \geq 3$, k -EDGE COLOURING is NP-complete even for k -regular graphs.*

Due to the above hardness results we may wish to restrict the input to some special graph class. A natural property of a graph class is to be closed under vertex deletion. Such graph classes are called *hereditary* and they form the focus of our paper. To give an example, bipartite graphs form a hereditary graph class, and it is well-known that they have chromatic index Δ . Hence, EDGE COLOURING is polynomial-time solvable for bipartite graphs, which are perfect and triangle-free. In contrast, Cai and Ellis [4] proved that for every $k \geq 3$, k -EDGE COLOURING is NP-complete for k -regular comparability graphs, which are also perfect. They also proved the following two results, the first one of which shows that EDGE COLOURING is NP-complete for triangle-free graphs (the graph C_s denotes the cycle on s vertices).

Theorem 3 ([4]). *Let $k \geq 3$ and $s \geq 3$. Then k -EDGE COLOURING is NP-complete for k -regular C_s -free graphs.*

Theorem 4 ([4]). *Let $k \geq 3$ be an odd integer. Then k -EDGE COLOURING is NP-complete for k -regular line graphs of bipartite graphs.*

It is also known that EDGE COLOURING is polynomial-time solvable for chordless graphs [22], series-parallel graphs [16], split-indifference graphs [26] and for graphs of treewidth at most k for any constant k [1].

It is not difficult to see that a graph class \mathcal{G} is hereditary if and only if it can be characterized by a set $\mathcal{F}_{\mathcal{G}}$ of forbidden induced subgraphs (see, for example, [17]). Malyshev determined the complexity of 3-EDGE COLOURING for every hereditary graph class \mathcal{G} , for which $\mathcal{F}_{\mathcal{G}}$ consists of graphs that each have at most five vertices, except perhaps two graphs that may contain six vertices [23]. Malyshev performed a

^{*} The second author was supported by the Research Council of Norway via the project CLASSIS. The third author was supported by the Leverhulme Trust (RPG-2016-258).

similar complexity study for EDGE COLOURING for graph classes defined by a family of forbidden (but not necessarily induced) graphs with at most seven vertices and at most six edges [24].

We focus on the case where \mathcal{F}_G consists of a single graph H . A graph G is H -free if G does not contain an induced subgraph isomorphic to H . We obtain the following dichotomy for H -free graphs.

Theorem 5. *Let $k \geq 3$ be an integer and H be a graph. If H is a linear forest, then k -EDGE COLOURING is polynomial-time solvable for H -free graphs. Otherwise k -EDGE COLOURING is NP-complete even for k -regular H -free graphs.*

We obtain Theorem 5 by combining Theorems 3 and 4 with two new results. In particular, we will prove a hardness result for k -regular claw-free graphs for even integers k (as Theorem 4 is only valid when k is odd).

2 Preliminaries

The graphs C_n , P_n and K_n denote the path, cycle and complete graph on n vertices, respectively. A set I is an *independent set* of a graph G if all vertices of I are pairwise nonadjacent in G . A graph G is *bipartite* if its vertex set can be partitioned into two independent sets A and B . If there exists an edge between every vertex of A and every vertex of B , then G is *complete bipartite*. The *claw* $K_{1,3}$ is the complete bipartite graph with $|A| = 1$ and $|B| = 3$.

Let G_1 and G_2 be two vertex-disjoint graphs. The *join* operation \times adds an edge between every vertex of G_1 and every vertex of G_2 . The *disjoint union* operation $+$ merges G_1 and G_2 into one graph without adding any new edges, that is, $G_1 + G_2 = (V(G_1) \cup V(G_2), E(G_1) \cup E(G_2))$. We write rG to denote the disjoint union of r copies of a graph G .

A *forest* is a graph with no cycles. A *linear forest* is a forest of maximum degree at most 2, or equivalently, a disjoint union of one or more paths. A graph G is a *cograph* if G can be generated from K_1 by a sequence of join and disjoint union operations. A graph is a cograph if and only if it is P_4 -free (see, for example, [3]). The following well-known lemma follows from this equivalence and the definition of a cograph.

Lemma 1. *Every connected P_4 -free graph on at least two vertices has a spanning complete bipartite subgraph.*

Let $G = (V, E)$ be a graph. For a subset $S \subseteq V$, the graph $G[S] = (S, \{uv \in E \mid u, v \in S\})$ denotes the subgraph of G induced by S . We say that S is *connected* if $G[S]$ is connected. Recall that a graph G is H -free for some graph H if G does not contain H as an induced subgraph. A subset $D \subset V(G)$ is *dominating* if every vertex of $V(G) \setminus D$ is adjacent to least one vertex of D . We will need the following result of Camby and Schaudt.

Theorem 6 ([5]). *Let $t \geq 4$ and G be a connected P_t -free graph. Let X be any minimum connected dominating set of G . Then $G[X]$ is either P_{t-2} -free or isomorphic to P_{t-2} .*

Let $G = (V, E)$ be some graph. The *degree* of a vertex $u \in V$ is equal to the size of its neighbourhood $N(u) = \{v \mid uv \in E\}$. The graph G is *r -regular* if every vertex of G has degree r . The *line graph* of G is the graph $L(G)$, which has vertex set E and an edge between two distinct vertices e and f if and only if e and f have a common end-vertex in G .

3 The Proof of Theorem 5

To prove our dichotomy, we first consider the case where the forbidden induced subgraph H is a claw. As line graphs are claw-free, we only need to deal with the case where the number of colours k is even due to Theorem 4. For proving this case we need another result of Cai and Ellis, which we will use as a lemma. Let c be a k -edge colouring of a graph $G = (V, E)$. Then a vertex $u \in V$ *misses* colour i if none of the edges incident to u is coloured i .

Lemma 2 ([4]). For even $k \geq 2$, the complete graph K_k has a k -edge colouring with the property that $V(K_k)$ can be partitioned into sets $\{u_i, u'_i\}$ ($1 \leq i \leq \frac{k}{2}$), such that for $i = 1, \dots, \frac{k}{2}$, vertices u_i and u'_i miss the same colour, which is not missed by any of the other vertices.

We use Lemma 2 to prove the following result, which solves the case where k is even and $H = K_{1,3}$.

Lemma 3. Let $k \geq 4$ be an even integer. Then k -EDGE COLOURING is NP-complete for k -regular claw-free graphs.

Proof. Recall that k -EDGE COLOURING for k -regular graphs is NP-complete for every integer $k \geq 4$ due to Theorem 2. Consider an instance (G, k) of k -EDGE COLOURING, where G is k -regular for some even integer $k = 2\ell \geq 4$. From G we construct a graph G' as follows. First we replace every vertex v in G by the gadget $H(v)$ shown in Figure 1. Next we connect the different gadgets in the following way. Every gadget $H(v)$ has exactly k pendant edges, which are incident with vertices $v_1, \dots, v_\ell, v_{\ell+1}, \dots, v_{2\ell}$, respectively. As G is k -regular, every vertex has k neighbours in G . Hence, we can identify each edge uv of G with a unique edge $u_h v_i$ in G' , which is a pendant edge of both $H(u)$ and $H(v)$. It is readily seen that G' is k -regular and claw-free.




Fig. 1. The gadget $H(v)$ where $K_i(v)$ is a complete graph of size 2ℓ for $i = 1, 2$. Note that edges inside $K_1(v)$ and $K_2(v)$ are not drawn.

First suppose that G is k -edge colourable. Let c be a k -edge colouring of G . Consider a vertex $v \in V(G)$. For every neighbour u of v in G , we colour the pendant edge in $H(v)$ corresponding to the edge uv with colour $c(uv)$. As c assigned different colours to the edges incident to v , the 2ℓ pendant edges of $H(v)$ will receive pairwise distinct colours, which we denote by $x_1, \dots, x_\ell, y_1, \dots, y_\ell$. By Lemma 2, we can colour the edges of $K_1(v)$ in such a way that for $i = 1, \dots, \ell$, v_i and v'_i miss colour x_i . For $i = 1, \dots, \ell$, we can therefore assign colour x_i to edge $v'_i w$. Similarly, we may assume that for $i = 1, \dots, \ell$, $v_{\ell+i}$ and $v'_{\ell+i}$ miss colour y_i . For $i = 1, \dots, \ell$, we can therefore assign colour y_i to edge $v'_{\ell+i} w$. Recall that the colours $x_1, \dots, x_\ell, y_1, \dots, y_\ell$ are all different. Hence, doing this procedure for each vertex of G yields a k -edge colouring c' of G' .

Now suppose that G' is k -edge colourable. Let c' be a k -edge colouring of G' . Consider some $v \in V(G)$. Denote the pendant edges of $H(v)$ by e_i for $i = 1, \dots, 2\ell$, where e_i is incident to v_i (and to some vertex u_h in a gadget $H(u)$ for each neighbour u of v in G). Suppose that c' gave colour x to an edge wv'_i for some $1 \leq i \leq \ell$, say to wv'_1 , but not to any edge e_i for $i = 1, \dots, \ell$. Note that wv'_2, \dots, wv'_ℓ cannot be coloured x . As every vertex of G' has degree $k = 2\ell$, every v_i with $1 \leq i \leq \ell$ and every v'_j with $2 \leq j \leq \ell$ is incident to some edge coloured x . As x is neither the colour of e_1, \dots, e_ℓ nor the colour of wv'_2, \dots, wv'_ℓ , the complete graph $K_1(v) - v'_1$ contains a perfect matching all of whose edges have colour x . However, $K_1(v) - v'_1$ has odd size $2\ell - 1$. Hence, this is not possible. We conclude that each of the (pairwise distinct) colours of wv'_1, \dots, wv'_ℓ , which we denote by x_1, \dots, x_ℓ , is the colour of an edge e_i for some $1 \leq i \leq \ell$.

Let y_1, \dots, y_ℓ be the (pairwise distinct) colours of $wv'_{\ell+1}, \dots, wv'_{2\ell}$, respectively. By the same arguments as above, we find that each of those colours is also the colour of a pendant edge of $H(v)$ that is incident to a vertex $v_{\ell+i}$ for some $1 \leq i \leq \ell$. Note that $x_1, \dots, x_\ell, y_1, \dots, y_\ell$ are 2ℓ pairwise distinct colours, as they are colours of edges incident to the same vertex, namely vertex w . Hence, we can define a k -colouring c of G by setting $c(uv) = c'(u_h v_i)$ for every edge $uv \in E(G)$ with corresponding edge $u_h v_i \in E(G')$. \square

We note that the graph G' in the proof of Lemma 3 is not a line graph, as the gadget $H(v)$ is not a line graph: the vertices v'_1, v'_2, v_1, w form a diamond and by adding the pendant edge incident to v_1 and the edge $wv'_{\ell+1}$ we obtain an induced subgraph of $H(v)$ that is not a line graph.

To handle the case where the forbidden induced subgraph H is a path, we make the following observation.

Observation 1 *If a graph G of maximum degree k has a dominating set of size at most p , then G has at most $p(k+1)$ vertices.*

We use Observation 1 to prove the following lemma.

Lemma 4. *Let $k \geq 0$ and $t \geq 1$. Every connected P_t -free graph of maximum degree k has at most $f(k, t)$ vertices for some function f that only depends on k and t .*

Proof. Let G be a connected P_t -free graph of maximum degree at most k . We use induction on t .

First suppose $t = 4$ (and observe that if the claim holds for $t = 4$, it also holds for $t \leq 3$). As G is connected, G has a dominating set of size 2 due to Lemma 1. Hence, by Observation 1, we find that G has at most $f(k, 2) = 2(k+1)$ vertices.

Now suppose $t \geq 5$. Let X be an arbitrary minimum connected dominating set of G . By Theorem 6, $G[X]$ is either P_{t-2} -free or isomorphic to P_{t-2} . In the first case we use the induction hypothesis to conclude that $G[X]$ has at most $f(k, t-2)$ vertices. Hence, G has at most $f(k, t-2)(k+1)$ vertices by Observation 1. In the second case, we find that G has at most $(t-2)(k+1)$ vertices. We set $f(k, t) = \max\{f(k, t-2)(k+1), (t-2)(k+1)\}$. \square

We use Lemma 4 to prove our next lemma.

Lemma 5. *Let $k \geq 3$ and $t \geq 1$. Then k -EDGE COLOURING is linear-time solvable for P_t -free graphs.*

Proof. Let G be a P_t -free graph. We compute the set of connected components of G in linear time. For each connected component D of G we do as follows. We first compute in linear time the maximum degree Δ_D of D . If $\Delta_D \leq k-1$, then D is k -edge colourable by Theorem 1. If $\Delta_D \geq k+1$, then D is not k -edge colourable. Hence, we may assume that $\Delta_D = k$. By Lemma 4, D has at most $f(k, t)$ vertices for some function f that only depends on k and t . As we assume that k and t are constants, this means that we can now check in constant time if D is k -edge colourable. Note that G is k -edge colourable if and only if every connected component of G is k -edge colourable. Hence, by using the above procedure, deciding if G is k -edge colourable takes linear time. \square

We are now ready to prove Theorem 5, which we restate below.

Theorem 5. (restated) *Let $k \geq 3$ be an integer and H be a graph. If H is a linear forest, then k -EDGE COLOURING is linear-time solvable for H -free graphs. Otherwise k -EDGE COLOURING is NP-complete even for k -regular H -free graphs.*

Proof. First suppose that H contains a cycle C_s for some $s \geq 3$. Then the class of H -free graphs is a superclass of the class of C_s -free graphs. This means that we can apply Theorem 3. From now on assume that H contains no cycle, so H is a forest. Suppose that H contains a vertex of degree at least 3. Then the class of H -free graphs is a superclass of the class of $K_{1,3}$ -free graphs, which in turn forms a superclass of the class of line graphs. Hence, if k is odd, then we apply Theorem 4, and if k is even, then we apply Lemma 3. From now on assume that H contains no cycle and no vertex of degree at least 3. Then H is a linear forest, say with ℓ connected components. Let $t = \ell|V(H)|$. Then the class of H -free graphs is contained in the class of P_t -free graphs. Hence we may apply Lemma 5. This completes the proof of Theorem 5. \square

4 Conclusions

We gave a complete complexity classification of k -EDGE COLOURING for H -free graphs, showing a dichotomy between linear-time solvable cases and NP-complete cases. We saw that this depends on H being a linear forest or not. It would be interesting to prove a dichotomy result for EDGE COLOURING restricted to H -free graphs. Note that due to Theorem 5 we only need to consider the case where H is a linear forest. However, even determining the complexity for small linear forests H , such as the cases where $H = 2P_2$ and $H = P_4$, turns out to be a difficult problem. In fact, the computational complexity of EDGE COLOURING for split graphs, or equivalently, $(2P_2, C_4, C_5)$ -free graphs [10] and for P_4 -free graphs has yet to be settled, despite the efforts towards solving the problem for these graph classes [6, 8, 21].

On a side note, a graph is k -edge colourable if and only if its line graph is k -vertex colourable. In contrast to the situation for EDGE COLOURING, the computational complexity of VERTEX COLOURING has been fully classified for H -free graphs [19]. However, the computational complexity for k -VERTEX COLOURING restricted to H -free graphs has not been fully classified. It is known that for every $k \geq 3$, k -VERTEX COLOURING on H -free graphs is NP-complete if H contains a cycle [9] or an induced claw [14, 20], but the case where H is a linear forest has not been settled yet. The complexity status of k -VERTEX COLOURING is even still open for P_t -free graphs. More precisely, it is known that the cases $k \leq 2$, $t \geq 1$ (trivial), $k \geq 3$, $t \leq 5$ [13], $k = 3$, $6 \leq t \leq 7$ [2] and $k = 4$, $t = 6$ [7] are polynomial-time solvable and that the cases $k = 4$, $t \geq 7$ [15] and $k \geq 5$, $t \geq 6$ [15] are NP-complete. However, the remaining cases, that is, the cases where $k = 3$ and $t \geq 8$ are still open. We refer to the survey [11] or some recent papers [12, 18, 25] for further background information.

Acknowledgements

The present work was done when the second author was visiting the University of Fribourg funded by a scholarship of the University of Fribourg.

References

1. H. L. Bodlaender. Polynomial algorithms for graph isomorphism and chromatic index on partial k -trees. *Journal of Algorithms*, 11(4):631–643, 1990.
2. F. Bonomo, M. Chudnovsky, P. Maceli, O. Schaudt, M. Stein, and M. Zhong. Three-coloring and list three-coloring of graphs without induced paths on seven vertices. *Combinatorica*, 38:779–801, 2018.
3. A. Brandstädt, V. B. Le, and J. P. Spinrad. *Graph Classes: A Survey*. SIAM Monographs on Discrete Mathematics and Applications. Society for Industrial and Applied Mathematics (SIAM), 1999.
4. L. Cai and J. A. Ellis. NP-completeness of edge-colouring some restricted graphs. *Discrete Applied Mathematics*, 30(1):15–27, 1991.
5. E. Camby and O. Schaudt. A new characterization of P_k -free graphs. *Algorithmica*, 75(1), 2016.
6. B.-L. Chen, H.-L. Fu, and M. T. Ko. Total chromatic number and chromatic index of split graphs. *Journal of Combinatorial Mathematics and Combinatorial Computing*, 17:137–146, 1995.
7. M. Chudnovsky, S. Spirkl, and M. Zhong. Four-coloring P_6 -free graphs. *Proc. SODA 2019*, pages 1239–1256, 2019.
8. S. M. de Almeida, C. P. de Mello, and A. Morgana. Edge-coloring of split graphs. *Ars Combinatoria*, 119:363–375, 2015.
9. T. Emden-Weinert, S. Hougardy, and B. Kreuter. Uniquely colourable graphs and the hardness of colouring graphs of large girth. *Combinatorics, Probability and Computing*, 7(04):375–386, 1998.
10. S. Földes and P. L. Hammer. Split graphs. *Congressus Numerantium*, XIX:311–315, 1977.
11. P. A. Golovach, M. Johnson, D. Paulusma, and J. Song. A survey on the computational complexity of colouring graphs with forbidden subgraphs. *Journal of Graph Theory*, 84(4):331–363, 2017.
12. C. Groenland, K. Okrasa, P. Rzażewski, A. Scott, P. Seymour, and S. Spirkl. H -colouring P_t -free graphs in subexponential time. *CoRR*, 1803.05396, 2018.
13. C. T. Hoàng, M. Kamiński, V. V. Lozin, J. Sawada, and X. Shu. Deciding k -colorability of P_5 -free graphs in polynomial time. *Algorithmica*, 57(1):74–81, 2010.
14. I. Holyer. The NP-completeness of edge-coloring. *SIAM Journal on Computing*, 10(4):718–720, 1981.

15. S. Huang. Improved complexity results on k -coloring P_t -free graphs. *European Journal of Combinatorics*, 51:336–346, 2016.
16. D. S. Johnson. The NP-completeness column: An ongoing guide. *J. Algorithms*, 6(3):434–451, 1985.
17. S. Kitaev and V. V. Lozin. *Words and Graphs*. Monographs in Theoretical Computer Science. An EATCS Series. Springer, 2015.
18. T. Klímašová, J. Malík, T. Masařík, J. Novotná, D. Paulusma, and V. Slívová. Colouring $(P_r + P_s)$ -free graphs. *Proc. ISAAC 2018, LIPIcs*, 123:5:1–5:13, 2018.
19. D. Král', J. Kratochvíl, Zs. Tuza, and G. J. Woeginger. Complexity of coloring graphs without forbidden induced subgraphs. *Proc. WG 2001, LNCS*, 2204:254–262, 2001.
20. D. Leven and Z. Galil. NP completeness of finding the chromatic index of regular graphs. *Journal of Algorithms*, 4(1):35–44, 1983.
21. A. R. C. Lima, G. Garcia, L. Zatesko, and S. M. de Almeida. On the chromatic index of cographs and join graphs. *Electronic Notes in Discrete Mathematics*, 50:433–438, 2015.
22. R. C. S. Machado, C. M. H. de Figueiredo, and N. Trotignon. Edge-colouring and total-colouring chordless graphs. *Discrete Mathematics*, 313(14):1547–1552, 2013.
23. D. S. Malyshev. The complexity of the edge 3-colorability problem for graphs without two induced fragments each on at most six vertices. *Sib. elektr. matem. izv.*, 11:811–822, 2014.
24. D. S. Malyshev. Complexity classification of the edge coloring problem for a family of graph classes. *Discrete Mathematics and Applications*, 27:97–101, 2017.
25. D. S. Malyshev. The complexity of the vertex 3-colorability problem for some hereditary classes defined by 5-vertex forbidden induced subgraphs. *Graphs and Combinatorics*, 33(4):1009–1022, 2017.
26. C. Ortiz, N. Maculan, and J. L. Szwarcfiter. Characterizing and edge-colouring split-indifference graphs. *Discrete Applied Mathematics*, 82(1-3):209–217, 1998.
27. V. G. Vizing. On an estimate of the chromatic class of a p -graph. *Diskret. Analiz.*, 3:25–30, 1964.