

Durham Research Online

Deposited in DRO:

16 September 2021

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

McHugh, C. and Hind, K. and O'Halloran, A. and Davey, D. and Farrell, G. and Wilson, F. (2021) 'Body Mass and Body Composition Changes over 7 Years in a Male Professional Rugby Union Team.', *International Journal of Sports Medicine* .

Further information on publisher's website:

<https://doi.org/10.1055/a-1403-2906>

Publisher's copyright statement:

© 2021. Thieme. All rights reserved.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

1 **Accepted: IJSM, April 2021**

2 **Body mass and body composition changes over 7 years in male professional rugby union**
3 **team.**

4

5 Clíodhna McHugh, Karen Hind, Aoife O'Halloran, Daniel Davey, Gareth Farrell, Fiona Wilson

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

1 **Abstract**

2 The purpose of this study was to investigate longitudinal body mass and body composition
3 changes in rugby union players (n= 123) from one professional club, (i) according to position
4 [forwards (n= 58) versus backs (n= 65)], (ii) analysis of players with 6 consecutive seasons of
5 DXA scans (n= 21) and, (iii) to examine differences by playing status [academy and
6 international], over 7 years. Players [mean age: 26.8y, body mass index: 28.9kg.m²] received
7 DXA scans at four-time points within each year. A modest (but non-significant) increase in
8 mean total mass (0.8kg) for professional players was reflected by increased lean mass and
9 reduced body fat mass. At all-time points, forwards had a significantly greater total mass,
10 lean mass and body fat percentage compared to backs (p< 0.05). Academy players
11 demonstrated increased total and lean mass and decreased body fat percentage over the
12 first 3 years of senior rugby, although this was not significant. Senior and academy
13 international players had greater lean mass and lower body fat percentage (p< 0.05) than
14 non-international counterparts. Despite modest increases in total mass; reflected by
15 increased lean mass and reduced fat mass, no significant changes in body mass or body
16 composition, irrespective of playing position were apparent over 7 years.

17

18

19

20

21

- 1 **Keywords:** Body composition; Dual-energy X-ray absorptiometry; Athletes; Longitudinal
- 2 analysis.

1 Introduction

2 Rugby Union (herein referred to as 'rugby') is a physically demanding, high-intensity,
3 collision sport [1], with distinct differences in body mass required for playing positions [2].
4 Body composition in rugby is an important component to success, as power-to-body mass
5 ratio underlies many of the sporting movements [3]. Previous research provides insight into
6 compositional characterisation [4, 5], positional differences [1, 6], and inter-seasonal
7 changes [1, 4]. Jones et al was the first to investigate longitudinal body composition changes
8 in rugby league players over 6 years and, demonstrated the individuality of changes [7].
9 However, interpretation of findings is limited due to the small sample size (n= 12) and
10 absence of analysis based on positions [1]. It is also not clear if the findings can be
11 generalised to rugby union, as there are different demands between the two rugby codes.

12 Since 1955, the average rugby players' body mass has increased by approximately 25%,
13 from 85 to 105kg [8]. It remains unknown if players may reach a point where increasing
14 mass is not a consequence of lean mass (LM) but rather body fat [9, 10]. Research supports
15 that lower skinfolds are associated with greater game time [11], and physical performance
16 [12]. Therefore, excess fat mass (FM) is counterproductive to the power-to-weight ratio, as
17 well as acceleration and metabolic efficiency of players [13]. The body composition of
18 players are routinely assessed, predominately to monitor development of physiological
19 capacities (e.g. speed, aerobic fitness) [14], and injury prevention [4, 15]. Direct
20 measurements of body fat percentage (%BF), measured by dual-energy x-ray
21 absorptiometry (DXA); a superior tool for providing accurate and highly detailed body
22 composition assessment in athletic populations [16].

1 To date, there is a dearth of research investigating longitudinal body composition changes in
2 rugby players, particularly investigating if mass is continuing to increase. Therefore, the aim
3 of this study was to explore 7- year longitudinal DXA data from one professional rugby team
4 to identify body composition trends. Fuller et al, reported that the mean total mass (TM) of
5 players has increased since 2002, and significantly for forwards ($p < 0.01$)[17]. However, to
6 our knowledge, no study has analysed the compositional components responsible for the
7 increased TM. Secondly, we aimed to investigate the longitudinal body composition changes
8 of players with 6 years of DXA scan data. Furthermore, no study has compared body
9 composition profiles of players within the same club, based on their international playing
10 status. This study also aimed to explore body composition based on playing status and
11 during the transition from academy to senior rugby. A secondary aim was to explore %BF
12 classification, derived from DXA.

13

14 **Materials and methods**

15 *Study design*

16 The study assessed 7- year longitudinal changes in TM, LM, FM and %BF in professional
17 rugby players from one club between 2012 and 2019. Players were scanned at 4 time-points
18 each season; baseline, pre-season, mid-season and post-season (Figure 1).

19

20

Insert Figure 1

21

22 *Participants*

1 A total of 123 professional male rugby players from one European Rugby Championship Cup
2 team received DXA scans over 7 years. DXA scan data was available for 21 players for 6 of
3 the 7 consecutive years. Players were grouped by position. Forwards included props,
4 hookers, locks, back. Backs included centres, scrum-halves, fly-halves, wingers and fullbacks.
5 The number of players each season ranged from 19 to 63, with an even distribution
6 between positions (Table 1).

7 Sub-group analyses of senior international players (any senior club player playing
8 international rugby between 2015-2019) and non-international players (any senior club
9 player not playing international rugby between 2015-2019). Further sub-group analysis of
10 academy players aged 17-20 years (any player playing academy rugby prior to transition to
11 senior rugby), by playing position was included. Academy players' DXA scans were analysed
12 using the same time point for each player during their first 3- years as a senior rugby player.
13 Players diet were controlled by the lead nutritionist, specific to positional demands and
14 training days; aerobic, resistance and rest. This study was conducted in accordance with ethical
15 standards in sport and exercise science research [18]. Ethical approval was provided by the
16 Institution Research Ethics Committee. Additional approval was obtained to access pseudo-
17 anonymised data from the host club for the time period of the study.

18

19 *Body composition scan acquisition and analysis*

20 All players on the roster between 2012 and 2019 received total body DXA (EnCore version
21 15.0, GE Lunar Healthcare, Madison, WI) scans across the season. Standardised scanning
22 protocols were followed to ensure consistency in scan acquisition [16]. All scans were
23 conducted by a skilled technologist and analysed following the manufacturer's guidelines

1 [19]. Athletes were scanned early in the morning (7:00 - 9:00 am), prior to food or fluid
2 ingestion, in a euhydrated state with void bladder, and wearing minimal clothing. Height
3 (cm) and weight (kg) were measured prior to scans. Athletes lay in a supine position on the
4 DXA scanner bed and were positioned with hands in a fully pronated position with an
5 approximate 5cm gap between hands and thigh and the use of GE positioning straps at
6 lower leg to support consistent positioning. The mode used was automatically selected by
7 the software and was dependent on body thickness (standard mode $\leq 25\text{cm}$; thick mode: $>$
8 25cm). Athletes were instructed to remain in position until otherwise instructed. Scans were
9 analysed using GE Lunar Encore software (Version 15.0) and were overseen by a clinical
10 densitometrist certified by the International Society of Clinical Densitometry. The DXA
11 system was serviced annually by the manufacturer and a daily calibration protocol provided
12 quality assurance. No significant drift in calibration was reported during the study time
13 points. No deviation or software upgrades were reported.

14 The outcomes of interest derived from the DXA scan were to compare %BF values across the
15 7- years according to values published for male athletes (Gallagher et al., 2000) using DXA
16 scans: low ($<8.0\%$), normal (8.0–19.9%), above normal (20.0–24.9%), and high ($\geq 25.0\%$) [20,
17 21]. Body mass index (BMI) was calculated as kg.m^2 .

18

19 *Statistical analysis*

20 All analyses were carried out using 'R' version 3.6.1 (R Foundation for Statistical Computing,
21 Vienna, Austria) [22]. Descriptive statistics were calculated as mean [standard deviation
22 (SD)]. Data was found to be normally distributed. Standard linear models were used with
23 normality assumption to determine body composition differences for TM, LM, FM and %BF

1 by playing position; forwards v backs, international v non-international players and academy
2 v senior players. The Mann Kendall Trend tests were used to analyse data over 7 years
3 (2012/13 – 2018/19) for consistently increasing or decreasing trends (monotonic) in y-
4 values using baseline and post-season time points. Out of the full cohort, there were 21
5 players for whom a full, uninterrupted longitudinal data set (over 6 years) was available.
6 This data set was analysed separately using repeated measures ANOVA. Using post-season
7 scans, players were grouped by %BF classification [20, 21]. Distribution of %BF classifications
8 were analysed for all players and by position, international vs non-international and,
9 academy vs non-academy. A two-sample Kolmogorov Smirnov test was used to identify a
10 significant difference in distribution of %BF classification. Comparison of body composition
11 classification were determined using data from players last recorded scan. Changes to
12 academy players' body composition during the first 3 years of senior rugby were assessed
13 using data from the same time point for each player's first 3 seasons. Scan one was
14 compared to scan two and scan one was compared to scan three to investigate significant
15 changes. Significant differences are represented by $p < 0.05$.

16

17 **Results**

18 Table 1 presents the mean (SD) for TM, LM and %BF for baseline scans between 2012 and
19 2019. There were no significant changes in TM, LM or %BF for each year. Over the 7 years,
20 TM increased from 101.6 to 102.4kg with a reduction in variability (SD), LM increased from
21 80.1 to 81.9kg and, %BF decreased from 17.1 to 15.9%. The mean (SD) TM for players at
22 baseline were: year 1: 101.6kg (13.5), year 2: 102.3kg (12.7), year 3: 101.9kg (11.9), year 4:
23 102kg (12.7), year 5: 103kg (12.6), year 6: 101.3kg (12.6) and, year 7: 102.4kg (11.4).

1 Forwards demonstrated no significant change in TM, LM or %BF between the 7 years. Backs
2 demonstrated no significant change in TM, LM or %BF across the 7 years. Forwards had
3 significantly greater TM, LM and %BF than backs for each of the 7 years ($p < 0.05$).

4

5 ****Insert Table 1****

6

7 Using baseline and post-season data, no significant increase or decrease trends were found
8 for TM, LM or %BF for the team or by position between years ($p > 0.05$) (Table 2).

9

10 ****Insert Table 2****

11

12 DXA scans were performed on 21 players from 6 consecutive years. In this sub-group there
13 was a significant change in TM for all players over 6 years [$F(5, 100) = 32.4, p < 0.00$].
14 Significant differences in TM were identified between year 1 and 2 ($p < 0.01$), year 1 and year
15 3 ($p < 0.01$), year 1 and year 4 ($p < 0.01$) and year 3 and year 6 ($p < 0.05$). There was no
16 significant difference found for %BF, LM or FM. By position, forwards had significantly
17 greater mean TM, %BF, LM and FM for each of the 6 years. Forwards ($n = 14$) had a
18 significant change in TM [$F(5, 65) = 4.50, p < 0.01$], but no significant change in %BF, FM or
19 LM. Backs ($n = 7$) had no significant change in TM, %BF, LM or FM. No significant differences
20 in means for TM, %BF, LM or FM were identified.

1 Table 3 presents the %BF classification of players using post season scans for each year, by
2 position. There was a significant difference in the distribution of %BF classification between
3 forwards and backs across all 7 seasons ($p < 0.05$). A greater proportion of forwards were
4 categorised as having excess fat (above normal: 20-24.9% and high: >25%) compared to
5 backs for each year. In the 2018/19 season, no player had %BF in the high category,
6 although 6 forwards categorised as having %BF above normal. One back in 2018 and 2019
7 had a %BF value <8%. Backs were predominately categorised as normal (8-19.9%) and
8 forwards were predominately categorised as normal (8-19.9%) or above normal (20-24.9%).

9

10 ****Insert Table 3****

11

12 Thirty-four academy players with complete data from their first 3 seasons of exposure to
13 senior rugby were analysed. There were no significant changes in body composition
14 between first and second DXA scans or between first and third DXA scans ($p > 0.05$) (Table
15 4). During the first 3 years of senior rugby, academy players demonstrated non-significant
16 increases in TM and LM and reduction in FM and %BF. Sub-group analysis of academy
17 players by position, showed comparable body composition changes.

18

19 ****Insert Table 4****

20

21 Table 5 presents the mean differences (SD) of players based on their professional status;
22 international players or non-international players over 4 years (2015-19). The number of

1 backs and forwards between groups was evenly distributed for each year. International
2 players between 2015 and 2019 had significantly greater LM (mean difference: 3.3 - 6.3kg)
3 and significantly lower %BF (mean difference: 2.3 - 3.2%) than non-international players for
4 all 4 years ($p < 0.05$). International forwards had a significantly lower %BF than non-
5 international forwards for all 4 years ($p < 0.05$). Differences in %BF between international
6 and non-international backs was not significantly different for all 4 years (Table 5).

7

8

Insert Table 5

9

10 **Discussion**

11 We report that TM increased modestly in rugby players monitored over 7 years, reflecting a
12 longitudinal increase in LM and reduction in %BF, most notably in forwards. At all-time
13 points, forwards had a significantly greater TM, LM and %BF compared to backs ($p < 0.05$).

14 Cross-sectional analysis of all players over each individual season showed forwards
15 consistently had greater reduction in %BF, whereas backs had a greater increase in TM. In
16 the sub-group analysis of academy players transition to senior rugby, academy players
17 demonstrated increases in TM, LM and a decrease in %BF, although not significant.

18 International players had a more favourable body composition. These findings enable
19 greater understanding of changes to players body composition over time and at different
20 stages in their career.

21 A key finding includes that despite no significant difference, the modest increase in TM
22 between the 7 years was reflected by an increase in LM and decrease in %BF. The average

1 mid-season body mass for forwards was $111.1 \pm 7.8\text{kg}$ and backs was $94.8 \pm 9\text{kg}$, in the
2 2018-19 season. The average body mass for forwards during the 2019 World cup was 114kg;
3 the lightest forward weighing 80kg and the heaviest weighing 153kg [23]. Despite reports of
4 an international trend of substantial increase in average body mass of rugby players [17,
5 24], data from our cohort does not support this. Although it is plausible that increases in
6 body mass occurred before 2012 in this cohort. Furthermore, it has been suggested that
7 increases in TM is possibly related to changes in injury severity seen in rugby since 2002
8 [25].

9 Our finding that rugby forwards have greater %BF than backs is consistent with previous
10 research [1, 2, 6, 26]. Using data from 2018-19, forwards in this study had higher levels of
11 %BF compared to findings reported in previous studies [1, 5, 6, 27]. The negative health
12 consequences of elevated %BF are well documented [28, 29]. FM acts as ballast in
13 biomechanical terms, but adipose tissue is a vital endocrine organ for general health [30].
14 There remain several rugby forwards with %BF above desired healthy ranges. Rugby
15 forwards are taller and heavier as they are predominately engaged in static play;
16 scrummaging and rucking [31-34]. The higher %BF in forwards may provide protective
17 effects against injuries due to the higher frequency of tackles and contact [26, 35]. However,
18 this has not been found to be consistent and therefore, do not outweigh the potential long-
19 term cardiometabolic risks associated with elevated %BF [36, 37].

20 Sub-group analysis of 21 players (forwards: 14; backs: 7) with 6 consecutive years of data
21 indicated a significant increase in TM for all players and forwards, however no significant
22 change in %BF, LM or FM. It has been well reported that increasing mass through LM is
23 more beneficial to performance and health than FM [35]. Increased %BF has been shown to

1 have a negative relationship with performance [38], and potentially lead to increased injury
2 risk [39, 40].

3 Our findings indicate non-significant increases in TM and LM and concomitant decreases in
4 FM during the first 3 years of senior rugby. Body composition changes of academy players is
5 comparable for forwards and backs. Possible justifications for body composition changes,
6 include growth, maturation [41, 42], and exposure to a professional training environment
7 [43]. Till et al reported significant differences in anthropometrics of academy players over 4
8 years, and significant positional differences [43]. The lack of significant changes in our
9 cohort is possibly due to the average age of first scans was 17 years, where the most
10 significant changes have been reported between 16-17 years [42, 43]. Academy forwards
11 had a significantly greater TM, LM and, %BF compared to backs. When compared to Till et
12 al., (2016), our cohort have greater TM and lower %BF [27]. Compared to senior players,
13 academy players have a greater %BF, despite having a lower TM and LM. Academy players
14 are likely to still be in the natural growing years [41, 42]; therefore, monitoring of body
15 composition should be regarded as an important component of player monitoring [44].

16 Despite similar distribution of players per playing position; backs and forwards, international
17 rugby players were found to have significantly lower %BF and greater LM compared to non-
18 international players at all time points over 4 years (2015-19). To our knowledge, this is the
19 first study to investigate the difference in body composition between international and non-
20 international players from the same club. International players represent a subgroup of
21 rugby players who are performing at the highest level of selection achievable.

22 Characterisation of this population in comparison to those who did not achieve this level of
23 selection is useful for practitioners and future players aspiring to reach this playing status.

1 This difference in body composition may represent a genetic factor or independent training
2 component. All players within this study are from the same club and thus exposed to the
3 same training and nutritional programs. Thus, within the context of this study, it is not
4 possible to quantify the cause.

5 While the current study addresses multiple gaps in the literature, some limitations exist.

6 When assessing trends, baseline and post-season time points were used due to having the
7 least amount of missing data. Furthermore, findings are reflective of body composition
8 trends within one professional rugby club and is not representative of individual changes.

9 Despite classifying players by position, a larger sample size would allow for further
10 classification [33]. Generalisation of longitudinal findings are limited due to a small sample
11 size (n= 21) with 6 consecutive seasons of DXA data. Sub-group analysis of academy players
12 and international players have small number of participants; therefore, limiting the
13 generalisability of findings. Goals pertaining to body composition changes can be highly
14 specific to individual athletes. Although it is not possible to account for individual goals and
15 baseline body mass, data is presented to reflect changes specific to position, playing status
16 and, academy players. Finally, no formal hydration tests were performed.

17

18 *Conclusions*

19 Although no significant body composition trends, irrespective of position were apparent for
20 rugby players over 7 years, findings from this study provides useful information for
21 practitioners supporting the physical conditioning of rugby players. Although the sample
22 size analysing longitudinal changes is small, findings provide some insight into an area
23 previously unanswered. Rugby practitioners need to consider the cost benefit of increasing

1 player's mass for performance benefits and the potential long-term health risks associated
2 with elevated mass. Although mass is an integral component to performance, players with
3 increased mass were found to have a greater propensity to have %BF above desired healthy
4 ranges. This is particularly prudent to academy and non-international players who have
5 comparable TM to senior and international players but significantly different compositional
6 profiles. Differences between academy and senior players are expected. Findings from this
7 study provide insight into the longitudinal changes of academy players body composition.
8 Rugby practitioners need to be conscious of these differences and use a longitudinal
9 approach to data measurement to assess player development.

Figure legends

Figure 1: Timeline of DXA scan assessments within each season.

Table 1: Baseline body composition values for 7- years; from 2012 to 2019.

Table 2: Trend analysis for changes in body composition from 2012/13 to 2018/19 season.

Table 3: Classification of body fat percentage classification by playing position.

Table 4: Body composition of players during transition from academy to professional rugby.

Table 5: Body composition comparison between International players and non-international players.

References

1. Jones B, Till K, Barlow M et al. Anthropometric and Three-Compartment Body Composition Differences between Super League and Championship Rugby League Players: Considerations for the 2015 Season and Beyond. *PloS one* 2015; 10: e0133188. doi:10.1371/journal.pone.0133188.
2. Lees MJ, Oldroyd B, Jones B et al. Three-Compartment Body Composition Changes in Professional Rugby Union Players Over One Competitive Season: A Team and Individualized Approach. *J Clin Densitom* 2017; 20: 50-57. doi:10.1016/j.jocd.2016.04.010
3. Cronin JB, Hansen KT. Strength and power predictors of sports speed. *J Strength Cond Res* 2005; 19: 349-357.
4. Georgeson EC, Weeks BK, McLellan C et al. Seasonal change in bone, muscle and fat in professional rugby league players and its relationship to injury: a cohort study. *BMJ open* 2012; 2 (6). doi:10.1136/bmjopen-2012-001400
5. Harley JA, Hind K, O'hara JP. Three-compartment body composition changes in elite rugby league players during a super league season, measured by dual-energy X-ray absorptiometry. *J Strength Cond Res* 2011; 25: 1024-1029. doi: 10.1519/JSC.0b013e3181cc21fb
6. Morehen JC, Routledge HE, Twist C et al. Position specific differences in the anthropometric characteristics of elite European Super League rugby players. *Eur J Sport Sci* 2015; 15: 523-529. doi:10.1080/17461391.2014.997802
7. Jones B, Till K, Roe G et al. Six-year body composition change in male elite senior rugby league players. *J Sport Sci* 2018; 36: 266-271. doi:10.1080/02640414.2017.1300313
8. Hill NE, Rilstone S, Stacey MJ et al. Changes in northern hemisphere male international rugby union players' body mass and height between 1955 and 2015. *BMJ Open Sport Exerc Med* 2018; 4: e000459. doi: 10.1136/bmjsem-2018-000459
9. Noel MB, Vanheest JL, Zaneteas P et al. Body composition in Division I football players. *J Strength Cond Res* 2003; 17: 228-237.
10. McHugh C, Hind K, Wyse J et al. Increases in DXA-derived visceral fat across one season in professional rugby union players: importance of visceral fat monitoring in athlete body composition assessment. *J Clin Densitom* 2020. doi: 10.1016/j.jocd.2020.09.001
11. Gabbett TJ, Jenkins DG, Abernethy B. Relationships between physiological, anthropometric, and skill qualities and playing performance in professional rugby league players. *J Sport Sci* 2011; 29: 1655-1664. doi: [10.1080/02640414.2011.610346](https://doi.org/10.1080/02640414.2011.610346)
12. Till K, Cobley S, O'Hara J et al. Using anthropometric and performance characteristics to predict selection in junior UK Rugby League players. *J Sci Med Sport* 2011; 14: 264-269. doi: 10.1016/j.jsams.2011.01.006
13. Withers R, Craig N, Bourdon P et al. Relative body fat and anthropometric prediction of body density of male athletes. *Eur J Appl Physiol Occup Physiol* 1987; 56: 191-200. doi: 10.1007/BF00640643
14. Baker DG, Newton RU. Comparison of lower body strength, power, acceleration, speed, agility, and sprint momentum to describe and compare playing rank among professional rugby league players. *J Strength Cond Res* 2008; 22: 153-158. doi: 10.1519/JSC.0b013e31815f9519

15. Gabbett TJ. Changes in physiological and anthropometric characteristics of rugby league players during a competitive season. *J Strength Cond Res* 2005; 19: 400. doi: [10.1519/00124278-200505000-00027](https://doi.org/10.1519/00124278-200505000-00027)
16. Nana A, Slater GJ, Stewart AD et al. Methodology review: using dual-energy X-ray absorptiometry (DXA) for the assessment of body composition in athletes and active people. *Int J Sport Nutr Exerc Metab* 2015; 25: 198-215. doi:[10.1123/ijsnem.2013-0228](https://doi.org/10.1123/ijsnem.2013-0228)
17. Fuller CW, Taylor, A.E., Brooks, J.H. and Kemp, S.P. Changes in the stature, body mass and age of English professional rugby players: a 10-year review. *J Sports Sci* 2013; 31: 795-802. doi:[10.1080/02640414.2012.753156](https://doi.org/10.1080/02640414.2012.753156)
18. Harriss D, MacSween A, Atkinson G. Ethical standards in sport and exercise science research: 2020 update. *International journal of sports medicine* 2019; 40: 813-817
19. Systems GM. Lunar enCORE: Safety and Specification Manual In. Madison, Wisconsin GE Medical Systems Lunar; 2009: DF-1.5.
20. Santos DA, Dawson JA, Matias CN et al. Reference values for body composition and anthropometric measurements in athletes. *PloS one* 2014; 9: e97846. doi:[10.1371/journal.pone.0097846](https://doi.org/10.1371/journal.pone.0097846)
21. Gallagher D, Heymsfield SB, Heo M et al. Healthy percentage body fat ranges: an approach for developing guidelines based on body mass index. *Am J Clin Nutr* 2000; 72: 694-701. doi: [10.1093/ajcn/72.3.694](https://doi.org/10.1093/ajcn/72.3.694)
22. Team R Core. A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2012.
23. RugbyPass. Heaviest players at 2019 Rugby World Cup revealed 2019. In Internet doi:<https://www.rugbypass.com/news/heaviest-players-at-2019-rugby-world-cup-revealed/>; accessed 10 August 2020.
24. Olds T. The evolution of physique in male rugby union players in the twentieth century. *J Sport Sci* 2001; 19: 253-262. doi: [10.1080/026404101750158312](https://doi.org/10.1080/026404101750158312)
25. West S, Starling, L., Kemp, S.P., Williams, S., Cross, M., Taylor, A., Brooks, J. and Stokes, K. Trends in match injury risk in professional male rugby union—a 16-season review of 10 851 match injuries in the English Premiership (2002-2019): The Professional Rugby Injury Surveillance Project. *Br J Sports Med* 2020. doi: [10.1136/bjsports-2020-102529](https://doi.org/10.1136/bjsports-2020-102529)
26. Bell W, Evans W, Cobner D et al. The regional placement of bone mineral mass, fat mass, and lean soft tissue mass in young adult Rugby Union players. *Ergonomics* 2005; 48: 1462-1472. doi: [10.1080/00140130500101007](https://doi.org/10.1080/00140130500101007)
27. Till K, Jones B, O'Hara J et al. Three-compartment body composition in academy and senior rugby league players. *J Sports Physiol Perform* 2016; 11: 191-196. doi: [10.1123/ijsp.2015-0048](https://doi.org/10.1123/ijsp.2015-0048)
28. Björntorp P. Metabolic implications of body fat distribution. *Diabetes care* 1991; 14: 1132-1143. doi: [10.2337/diacare.14.12.1132](https://doi.org/10.2337/diacare.14.12.1132)
29. Scherrer U, Randin D, Tappy L et al. Body fat and sympathetic nerve activity in healthy subjects. *Circulation* 1994; 89: 2634-2640. doi: [10.1161/01.CIR.89.6.2634](https://doi.org/10.1161/01.CIR.89.6.2634)
30. Ackland TR, Lohman TG, Sundgot-Borgen J et al. Current status of body composition assessment in sport. *Sports Med* 2012; 42: 227-249. doi: [10.2165/11597140-000000000-00000](https://doi.org/10.2165/11597140-000000000-00000)
31. Roberts SP, Trewartha G, Higgitt RJ et al. The physical demands of elite English rugby union. *J Sport Sci* 2008; 26: 825-833. doi: [10.1080/02640410801942122](https://doi.org/10.1080/02640410801942122)

32. Austin D, Gabbett T, Jenkins D. The physical demands of Super 14 rugby union. *J Sci Med Sport* 2011; 14: 259-263. doi: 10.1016/j.jsams.2011.01.003
33. Cahill N, Lamb K, Worsfold P et al. The movement characteristics of English Premiership rugby union players. *J Sport Sci* 2013; 31: 229-237. doi: [10.1080/02640414.2012.727456](https://doi.org/10.1080/02640414.2012.727456)
34. Quarrie KL, Hopkins WG, Anthony MJ et al. Positional demands of international rugby union: Evaluation of player actions and movements. *J Sci Med Sport* 2013; 16: 353-359. doi: [10.1016/j.jsams.2012.08.005](https://doi.org/10.1016/j.jsams.2012.08.005)
35. Gabbett TJ. Science of rugby league football: a review. *J Sport Sci* 2005; 23: 961-976. doi: 10.1080/02640410400023381
36. Lenz M, Richter T, Mühlhauser I. The morbidity and mortality associated with overweight and obesity in adulthood: a systematic review. *Dtsch Arztebl Int* 2009; 106: 641. doi: 10.3238%2Farztebl.2009.0641
37. Hu FB, Willett WC, Li T et al. Adiposity as compared with physical activity in predicting mortality among women. *N Eng J Med* 2004; 351: 2694-2703. doi:10.1056/NEJMoa042135
38. Saunders PU, Pyne DB, Telford RD et al. Factors affecting running economy in trained distance runners. *Sports Med* 2004; 34: 465-485. doi: 10.2165/00007256-200434070-00005
39. Duthie GM. A framework for the physical development of elite rugby union players. *Int J Sports Physiol Perform* 2006; 1: 2-13. doi: <https://doi.org/10.1123/ijsp.1.1.2>
40. Higham D, Pyne D, Anson J et al. Distribution of fat, non-osseous lean and bone mineral mass in international rugby union and rugby sevens players. *Int J Sports Med* 2014; 35: 575-582. doi: 10.1055/s-0033-1355419
41. Malina RM, Bouchard C, Bar-Or O. Growth, maturation, and physical activity: Human kinetics; 2004.
42. Vääntinen T, Blomqvist M, Nyman K et al. Changes in body composition, hormonal status, and physical fitness in 11-, 13-, and 15-year-old Finnish regional youth soccer players during a two-year follow-up. *J Strength Cond Res* 2011; 25: 3342-3351. doi: 10.1519/JSC.0b013e318236d0c2
43. Till K, Jones B, Darrall-Jones J et al. Longitudinal development of anthropometric and physical characteristics within academy rugby league players. *J Strength Cond Res* 2015; 29: 1713-1722. doi: 10.1519/JSC.0000000000000792
44. Malina RM. Body composition in athletes: assessment and estimated fatness. *Clin Sports Med* 2007; 26: 37-68. doi: 10.1016/j.csm.2006.11.004