

Durham Research Online

Deposited in DRO:

16 June 2014

Version of attached file:

Published Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Georgiou, Nicholas and Wade, Andrew R. (2014) 'Non-homogeneous random walks on a semi-infinite strip.', *Stochastic processes and their applications.*, 124 (10). pp. 3179-3205.

Further information on publisher's website:

<http://dx.doi.org/10.1016/j.spa.2014.05.005>

Publisher's copyright statement:

© 2014 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>).

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.



Non-homogeneous random walks on a semi-infinite strip

Nicholas Georgiou*, Andrew R. Wade

Department of Mathematical Sciences, Durham University, South Road, Durham, DH1 3LE, UK

Received 11 February 2014; received in revised form 15 May 2014; accepted 16 May 2014

Available online 21 May 2014

Abstract

We study the asymptotic behaviour of Markov chains (X_n, η_n) on $\mathbb{Z}_+ \times S$, where \mathbb{Z}_+ is the non-negative integers and S is a finite set. Neither coordinate is assumed to be Markov. We assume a moments bound on the jumps of X_n , and that, roughly speaking, η_n is close to being Markov when X_n is large. This departure from much of the literature, which assumes that η_n is itself a Markov chain, enables us to probe precisely the recurrence phase transitions by assuming asymptotically zero drift for X_n given η_n . We give a recurrence classification in terms of increment moment parameters for X_n and the stationary distribution for the large- X limit of η_n . In the null case we also provide a weak convergence result, which demonstrates a form of asymptotic independence between X_n (rescaled) and η_n . Our results can be seen as generalizations of Lamperti's results for non-homogeneous random walks on \mathbb{Z}_+ (the case where S is a singleton). Motivation arises from modulated queues or processes with hidden variables where η_n tracks an internal state of the system.

© 2014 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>).

MSC: primary 60J10; secondary 60F05; 60F15; 60K15; 60K25

Keywords: Non-homogeneous random walk; Recurrence classification; Weak limit theorem; Lamperti's problem; Modulated queues; Correlated random walk

* Corresponding author. Tel.: +44 1913344153.

E-mail addresses: nicholas.georgiou@durham.ac.uk (N. Georgiou), andrew.wade@durham.ac.uk (A.R. Wade).

1. Introduction

There are many applications that naturally give rise to Markov processes on a product state-space $\mathbb{X} \times S$ where S describes some operating regime or internal state of the system, which influences the motion of the process in the primary space \mathbb{X} . Important classes of examples include, among others,

- modulated queues, in which S may contain operating states of the servers or other auxiliary information such as the size of a retrial buffer, as arise in various applications such as those described by Neuts in [26];
- regime-switching processes in mathematical finance or ecology, where S may contain market or other environmental information;
- physical processes with internal degrees of freedom, where S may describe internal energy or momentum states of a particle, such as adopted by Sinai as a tool for studying the Lorentz gas (see e.g. [18]), or exemplified by the so-called correlated or persistent random walk.

In several of the key examples, the S -component of the process is ‘hidden’, and the main interest is in the asymptotic behaviour of the \mathbb{X} -component of the process.

In the most classical setting, the projection of the process onto S is itself Markovian. In this case, the queueing models become *Markov-modulated* [26,10], while other examples fit into the class of *Markov random walks* [14]. This case also includes processes that can be represented as *additive functionals of Markov chains* [27]. Such models pose a variety of mathematical questions, which have been studied rather deeply over several decades using various techniques that take advantage of the additional Markov structure, and much is now known.

Much less is known when the process projected onto S is *not* Markovian: the main focus of the present work is to replace the Markovian assumption by a weaker (asymptotic) condition that provides sufficient structure. This relaxation is necessary to probe more intimately the recurrence–transience phase transition for these models, since the natural setting (paralleling the classical work of Lamperti) is to suppose that the law of the process is *non-homogeneous* in \mathbb{X} , in particular, the mean drift of the \mathbb{X} -component of the process will be asymptotically zero. This non-homogeneity precludes, in general, the S -component of the process from being Markovian, but admits our weaker conditions.

To avoid technicalities, yet provide a setting rich enough to explore many interesting phenomena, we take \mathbb{X} to be the countable set $\mathbb{Z}_+ := \{0, 1, 2, \dots\}$ and take S to be finite. These models are already of interest for numerous applications, and there is an existing literature devoted to random walks on half strips ($\mathbb{Z}_+ \times \{0, 1, \dots, m\}$) or strips ($\mathbb{Z} \times \{0, 1, \dots, m\}$): see [9,24,7,8] and references therein.

As an example consider the following queueing model. A queue is served by a single server and experiences arrivals at rate λ ; the service rate is modulated via an internal state of the server η_n , as well as the length of the queue X_n (in discrete time, i.e., in terms of the jump process). Allowing the service rate to depend on the queue length distinguishes this model from the class of *semi-Markov* queues [26]. When $(X_n, \eta_n) = (x, i)$, $x \geq 1$, the service rate is $\rho_i(x) = \rho(1 - 2c_i/x)$, where c_i , $i \in S$ are parameters of model with $|c_i| < 1/2$. In the case where $c_i \equiv 0$ for all i , the internal states of the server are indistinguishable and the model is simply (the jump process of) an $M/M/1$ queue with arrival rate λ and service rate ρ ; the critical case from the point of view of recurrence and transience is $\rho = \lambda$, and so that is the most interesting setting to perturb with non-zero c_i . So we take $\rho = \lambda$ from now on. The specification of the model is completed by stipulating that whenever an arrival (departure) occurs the internal state of the server transitions according to the stochastic matrix (a_{ij}) ((b_{ij})). In other words,

given $(X_n, \eta_n) = (x, i), x \geq 1,$

$$(X_{n+1}, \eta_{n+1}) = \begin{cases} (x + 1, j) & \text{with probability } \frac{1}{2 \left(1 - \frac{c_i}{x}\right)} a_{ij}; \\ (x - 1, j) & \text{with probability } \frac{1 - \frac{2c_i}{x}}{2 \left(1 - \frac{c_i}{x}\right)} b_{ij}. \end{cases}$$

Given $(X_n, \eta_n) = (0, i), (X_{n+1}, \eta_{n+1}) = (1, j)$ with probability $a_{ij}.$

In general, (η_n) is not itself a Markov chain, so this model falls outside the usual Markov-modulated queue framework. However, for large queue lengths the probabilities of arrival and departure are approximately equal, and so the η_n process should be well approximated by the Markov chain on S with transition matrix $M_{ij} = (a_{ij} + b_{ij})/2.$ Under the condition that the matrix M be irreducible, our results determine conditions for transience and recurrence in terms of the stationary distribution of the chain with transition matrix M and the constants $c_i.$

2. Model and main results

We now describe precisely our model. Our state-space is the *half-strip* $\mathbb{Z}_+ \times S,$ where S is finite and nonempty; for $k \in S,$ we call the subset $\mathbb{Z}_+ \times \{k\}$ a *line.* We consider an irreducible Markov chain $(X_n, \eta_n) \in \mathbb{Z}_+ \times S,$ with transition probabilities

$$\mathbb{P}[(X_{n+1}, \eta_{n+1}) = (y, j) \mid (X_n, \eta_n) = (x, i)] = p(x, i, y, j), \tag{2.1}$$

and provide conditions for recurrence/transience of $(X_n),$ in a sense that we explain below. Throughout we use the notation $\mathcal{F}_n := \sigma(X_0, \eta_0, \dots, X_n, \eta_n)$ and $\mathbb{R}_+ := [0, \infty).$

The process (X_n) is typically not itself a Markov chain; under our standing assumptions, however, it does inherit the recurrence/transience dichotomy from $(X_n, \eta_n),$ as the following result shows.

Lemma 2.1. *Exactly one of the following holds:*

- (i) *If (X_n, η_n) is recurrent, then $\mathbb{P}[X_n = 0 \text{ i.o.}] = 1.$*
- (ii) *If (X_n, η_n) is transient, then $\mathbb{P}[X_n = 0 \text{ i.o.}] = 0,$ and $X_n \rightarrow \infty$ a.s.*

In the former case, we call (X_n) recurrent, and in the latter case, we call (X_n) transient.

Similarly, a natural distinction between positive- and null-recurrence holds.

Lemma 2.2. *There exists a (unique) measure ν on \mathbb{Z}_+ such that*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} \mathbf{1}\{X_k = x\} = \nu(x), \quad \text{a.s.}, \tag{2.2}$$

for any $x \in \mathbb{Z}_+.$ Exactly one of the following holds:

- (i) *If (X_n, η_n) is null, then $\nu(x) = 0$ for all $x \in \mathbb{Z}_+.$*
- (ii) *If (X_n, η_n) is positive-recurrent, then $\nu(x) > 0$ for all $x \in \mathbb{Z}_+$ and $\sum_{x \in \mathbb{Z}_+} \nu(x) = 1.$*

If (X_n) is recurrent, then we say that it is null-recurrent or positive-recurrent according to which of (i) or (ii) holds.

The proofs of [Lemmas 2.1](#) and [2.2](#) are standard and are omitted.

In the cases that we consider, we will assume that the displacement of the X -coordinate has bounded p -moments for some $p < \infty$:

(B_p) There exists a constant $C_p < \infty$ such that $\mathbb{E}[|X_{n+1} - X_n|^p \mid \mathcal{F}_n] \leq C_p$, a.s. $\forall n$.

In particular, (B_p) for some $p > 4$ will suffice for all of our results, while for some of our results $p > 1$ is sufficient.

Define $q_x(i, j) = \sum_{y \in \mathbb{Z}_+} p(x, i, y, j)$. We also assume:

(Q_∞) $\lim_{x \rightarrow \infty} q_x(i, j) = q(i, j)$ exists for all $i, j \in S$, and $(q(i, j))$ is an irreducible stochastic matrix.

Note that since $\sum_{j \in S} q_x(i, j) = 1$, the limit in (Q_∞) is necessarily stochastic; however the irreducibility of $(q(i, j))$ does not follow from the irreducibility of $(q_x(i, j))$ for all $x \in \mathbb{Z}_+$. For some of our results, it is necessary to assume a stronger condition than (Q_∞) that controls the rate of convergence of $q_x(i, j)$, namely:

(Q_∞^+) There exists $\delta_0 > 0$ such that $\max_{i, j \in S} |q_x(i, j) - q(i, j)| = O(x^{-\delta_0})$ as $x \rightarrow \infty$, and $(q(i, j))$ is an irreducible stochastic matrix.

Given (Q_∞), we define (η_n^*) to be a Markov chain on S with transition probabilities given by $q(i, j)$. Since (η_n^*) is irreducible and finite there exists a unique stationary distribution π on S with $\pi(j) > 0$ for all $j \in S$ and satisfying $\pi(j) = \sum_{i \in S} \pi(i)q(i, j)$.

Remark 2.3. A sufficient condition for (Q_∞^+) is that there exists $x_0 \in \mathbb{Z}_+$ such that

(H) $p(x, i, y, j) = r(y - x, i, j)$

for all $x \geq x_0$, i.e., for all x large enough, the transition probabilities depend on x and y only through $y - x$. Then, $q_x(i, j) = q(i, j) = \sum_{z \geq -x_0} r(z, i, j)$ for all $x \geq x_0$. The homogeneity condition (H) plays an important role in much of the existing literature, but is too restrictive for our purposes. We discuss (H) and some of its consequences, including the connection to the theory of additive functionals of Markov chains, in Section 3.1. For now, we remark that if (H) holds for all $x \geq x_0$, then necessarily $X_{n+1} - X_n$ is uniformly bounded below (by $-x_0$).

We denote the moments of the displacements in the X -coordinate by

$$\mu_k(x, i) := \mathbb{E}[(X_{n+1} - X_n)^k \mid X_n = x, \eta_n = i] = \sum_{j \in S} \sum_{y \in \mathbb{Z}_+} (y - x)^k p(x, i, y, j);$$

then μ_1 is well defined provided (B_p) holds for some $p \geq 1$, while μ_2 is finite if (B_p) holds for some $p \geq 2$. Our results will apply to the following two cases:

(M_C) There exist $d_i \in \mathbb{R}$ such that for all $i \in S$, as $x \rightarrow \infty$, $\mu_1(x, i) = d_i + o(1)$;

(M_L) There exist $c_i \in \mathbb{R}$ and $s_i^2 \in \mathbb{R}_+$, with at least one s_i^2 nonzero, such that for all $i \in S$, as $x \rightarrow \infty$, $\mu_1(x, i) = c_i/x + o(x^{-1})$ and $\mu_2(x, i) = s_i^2 + o(1)$.

Since S is finite, the implicit constants in the $x \rightarrow \infty$ error terms in these expressions (and similar ones later on) may be chosen uniformly over i . Just as above, some of our results will require a stronger assumption than (M_L) that controls the error terms as a function of x , namely:

(M_L^+) There exists $\delta_1 > 0$ such that, as $x \rightarrow \infty$,

$$\mu_1(x, i) = \frac{c_i}{x} + O(x^{-1-\delta_1}) \quad \text{and} \quad \mu_2(x, i) = s_i^2 + O(x^{-\delta_1}).$$

Next we state our main results. The first two are concerned with the classification of the process as transient, null-recurrent, or positive-recurrent. Of these, first we consider the case where

each line is associated with a drift that is asymptotically constant, and where at least one of these constants is nonzero.

Theorem 2.4. *Suppose that (B_p) holds for some $p > 1$, and conditions (M_C) and (Q_∞) hold. Then the following classification applies.*

- (i) *If $\sum_{i \in S} d_i \pi(i) > 0$, then X_n is transient.*
- (ii) *If $\sum_{i \in S} d_i \pi(i) < 0$, then X_n is positive-recurrent.*

In the special case of (Q_∞) in which $q_x \equiv q$ does not depend on x , **Theorem 2.4** is contained in **Theorem 3.1.2** of **Fayolle et al. [9]**, who imposed, in part, an assumption of a uniform lower bound on $X_{n+1} - X_n$. In the generality of (Q_∞) , part (ii) is contained in a paper of **Falin [7]**, who also stated a version of part (i) assuming that **(H)** holds for x large enough.

The next result deals with the case of drift conditions of Lamperti-type.

Theorem 2.5. *Suppose that (B_p) holds for some $p > 2$, and conditions (Q_∞) and (M_L) hold. The following sufficient conditions apply.*

- *If $\sum_{i \in S} (2c_i - s_i^2) \pi(i) > 0$, then X_n is transient.*
- *If $|\sum_{i \in S} 2c_i \pi(i)| < \sum_{i \in S} s_i^2 \pi(i)$, then X_n is null-recurrent.*
- *If $\sum_{i \in S} (2c_i + s_i^2) \pi(i) < 0$, then X_n is positive-recurrent.*

If, in addition, (Q_∞^+) and (M_L^+) hold, then the following condition also applies (yielding an exhaustive classification):

- *If $|\sum_{i \in S} 2c_i \pi(i)| = \sum_{i \in S} s_i^2 \pi(i)$, then X_n is null-recurrent.*

In the case where S is a singleton, **Theorem 2.5** reduces essentially to results of **Lamperti [19,21]**, and so our result can be seen as a generalization of **Lamperti’s**.

Our final main result concerns the weak convergence of (X_n, η_n) . The limit statement will involve the distribution function $F_{\alpha,\theta}$ defined for parameters $\alpha > 0$ and $\theta > 0$ as the (normalized) lower incomplete Gamma function,

$$F_{\alpha,\theta}(x) = \frac{\gamma(\alpha, x^2/\theta)}{\Gamma(\alpha)} = \frac{1}{\Gamma(\alpha)} \int_0^{x^2/\theta} z^{\alpha-1} e^{-z} dz, \quad (x \geq 0). \tag{2.3}$$

Note that, if $Z \sim \Gamma(\alpha, \theta)$ is a gamma random variable with shape parameter $\alpha > 0$ and scale parameter $\theta > 0$, then $\mathbb{P}[\sqrt{Z} \leq x] = F_{\alpha,\theta}(x)$. (In the special case with $\alpha = 1/2$ and $\theta = 2$, $F_{\alpha,\theta}$ is the distribution of the square-root of a χ^2 random variable with one degree of freedom, i.e., the absolute value of a standard normal random variable.)

Theorem 2.6. *Suppose that (B_p) holds for some $p > 4$, and conditions (Q_∞) and (M_L) hold. Suppose that the matrix q appearing in (Q_∞) is aperiodic. Suppose also that $\sum_{i \in S} (2c_i + s_i^2) \pi(i) > 0$. Then, for any $k \in S$ and $x \in \mathbb{R}_+$,*

$$\lim_{n \rightarrow \infty} \mathbb{P} \left[n^{-1/2} X_n \leq x, \eta_n = k \right] = \pi(k) F_{\alpha,\theta}(x),$$

where

$$\alpha = \frac{1}{2} + \frac{\sum_{i \in S} c_i \pi(i)}{\sum_{i \in S} s_i^2 \pi(i)}, \quad \text{and} \quad \theta = 2 \sum_{i \in S} s_i^2 \pi(i). \tag{2.4}$$

- Remarks 2.7.** (i) Under the hypothesis of [Theorem 2.6](#), [Theorem 2.5](#) shows that the process is null-recurrent or transient; [Theorem 2.6](#) demonstrates a form of asymptotic independence between X_n (rescaled) and η_n (which converges to π). By contrast, in the positive-recurrent aperiodic case, $\mathbb{P}[X_n \leq x, \eta_n = k]$ (with no scaling) possesses a limit, but that limit cannot be identified without additional assumptions (and the limit distribution of η_n need not even be π).
- (ii) The case of [Theorem 2.6](#) in which S is a singleton is essentially Lamperti’s weak convergence result from [20].
- (iii) If in addition (Q_∞^+) and (M_L^+) hold, then the boundary case $\sum_{i \in S} (2c_i + s_i^2)\pi(i) = 0$ is null-recurrent, by [Theorem 2.5](#). In this case the proof given in [Section 5.2](#) can be modified to show that $n^{-1/2}X_n \rightarrow 0$ in probability; this is consistent with the fact that the $\alpha \rightarrow 0$ limit of $F_{\alpha,\theta}$ corresponds to a point mass at 0.
- (iv) With some additional work, the arguments in [Section 5.2](#) should yield the process version of [Theorem 2.6](#): in the sense of finite dimensional distributions, as $n \rightarrow \infty$,

$$\left(n^{-1/2}X_{nt}, \eta_{nt} \right)_{t \in [0,1]} \longrightarrow (x_t, \omega_t)_{t \in [0,1]},$$

where $(2/\theta)^{1/2}x_t$ is a Bessel process with dimension 2α and ω_t is an S -valued white noise process whose finite-dimensional marginals are sequences of i.i.d. π -distributed variables.

The remainder of the paper is organized as follows. In [Section 3](#) we give some additional context to the present work by describing how our setting generalizes the literature on additive functionals of Markov chains, and by presenting some additional examples, including a variant of the correlated random walk. [Section 4](#) contains the bulk of our analysis, which proceeds via considering an embedded Markov chain. The proofs of the main theorems are then completed in [Section 5](#).

To simplify the presentation in the rest of the paper, we often write $\mathbb{P}_{x,i}[\cdot]$ for $\mathbb{P}[\cdot \mid X_0 = x, \eta_0 = i]$, corresponding to the law of the Markov chain with initial state $(x, i) \in \mathbb{Z}_+ \times S$; similarly for (expectation) $\mathbb{E}_{x,i}$.

We finish this section with some general remarks. Our method of proof is different from other approaches in the literature. Falin [7,8], while also making use of Foster–Lyapunov results, bases his computations on a delicate algebraic calculation. Rogers [27] uses an embedded Markov chain, as we do, but his analysis relies on the additive functional representation (see [Section 3.1](#)). Our approach to the excursion estimates for the embedded process, via the Doob decomposition, makes the emergence of the ‘pseudo-drift’ quantities particularly intuitive from a probabilistic perspective: see the discussion around [\(3.3\)](#).

The case where S is infinite can give rise to completely different phenomena from the finite setting, and we do not consider this here. Under suitable assumptions, however, such as uniform versions of our asymptotic conditions (Q_∞) , (M_C) or (M_L) , and sufficient moments for τ and the increments of X_n , the results of the present paper should extend to the infinite setting.

3. Examples and remarks on the literature

3.1. Homogeneity and additive functionals

As mentioned in [Remark 2.3](#), condition (H) is assumed in much of the literature. A special structure emerges when (H) is imposed for *all* x . Indeed, one then has that (η_n) itself is a Markov chain, since

$$\mathbb{P}[\eta_{n+1} = j \mid (X_n, \eta_n) = (x, i)] = \sum_{y \in \mathbb{Z}} r(y - x, i, j) = \sum_{z \in \mathbb{Z}} r(z, i, j) = q(i, j). \tag{3.1}$$

A similar argument shows that $(X_n - X_{n-1}, \eta_n)$ is a Markov chain on $\mathbb{Z} \times S$, with

$$\mathbb{P}[(X_{n+1} - X_n, \eta_{n+1}) = (z, j) \mid (X_n - X_{n-1}, \eta_n) = (y, i)] = r(z, i, j).$$

Then if $\psi : \mathbb{Z} \times S \rightarrow \mathbb{Z}$ is given by $\psi(z, i) = z$, we may write

$$X_n = X_0 + \sum_{k=0}^{n-1} \psi(X_{k+1} - X_k, \eta_{k+1}),$$

which represents X_n as an *additive functional of a Markov chain*.

However, on state-space $\mathbb{Z}_+ \times S$, assuming that (H) holds for all x is very restrictive, and implies that $X_{n+1} - X_n \geq 0$ a.s. (see Remark 2.3). So in the homogeneous setting, it makes sense to instead take the state space to be $\mathbb{Z} \times S$ so that (2.1) now holds with x and y in \mathbb{Z} . Assuming that (H) holds for all $x \in \mathbb{Z}$ now yields the additive functional structure above, without imposing additional restrictions on the magnitude of $X_{n+1} - X_n$.

In either case, we may note that

$$\mathbb{E}[X_{n+1} - X_n \mid (X_n, \eta_n) = (x, i)] = \sum_{z \in \mathbb{Z}} \sum_{j \in S} z r(z, i, j) =: \mu_1(i), \tag{3.2}$$

say, assuming that the mean increments are well defined; so there is a constant mean drift $\mu_1(i)$ for each $i \in S$.

Moreover, if π is the stationary distribution on S associated with the Markov chain (η_n) given by (3.1), then a calculation shows that the Markov chain $(X_n - X_{n-1}, \eta_n)$ has stationary distribution $\varpi(z, i)$ on $\mathbb{Z} \times S$ given by

$$\varpi(z, i) = \sum_{k \in S} \pi(k) r(z, k, i).$$

In this context, a result of Rogers [27] on additive functionals of Markov chains shows that recurrence classification of (X_n) depends on the sign of

$$\sum_{i \in S} \sum_{z \in \mathbb{Z}} \varpi(z, i) \psi(z, i) = \sum_{i \in S} \sum_{z \in \mathbb{Z}} z \sum_{k \in S} \pi(k) r(z, k, i) = \sum_{k \in S} \pi(k) \mu_1(k). \tag{3.3}$$

There are many similar results in the literature for additive functionals of Markov chains in more general spaces, and related results in ergodic theory concerning ‘co-cycles’ (see, e.g., [2]). However, the methods adapted to this additive functional structure seem to depend crucially on the homogeneity assumption (H).

The interpretation of the quantity of (3.3) is as a ‘pseudo-drift’ accumulated over i.i.d. excursions of the Markov chain: see Rogers [27]. We take this idea further, as the analogues of these excursions in our setting are not i.i.d., due to the additional non-homogeneity. However, our methods exploit the essential structure that remains.

3.2. Correlated random walk

In the one-dimensional *correlated random walk*, a particle performs a random walk on \mathbb{Z} with a short-term memory: the distribution of X_{n+1} depends not only on the current position X_n , but also on the ‘direction of travel’ $X_n - X_{n-1}$. Formally, $(X_n, X_n - X_{n-1})$ is a Markov chain on $\mathbb{Z} \times \{-1, +1\}$. Supposing also that (H) holds for all $x \in \mathbb{Z}$, this is a special case of the framework discussed in Section 3.1, with $\eta_n = X_n - X_{n-1}$.

One standard version of the model supposes that the nonzero transition probabilities are given by $p(x, i, x + j, j) = r(j, i, j) = q(i, j)$, where

$$q(i, j) = \begin{cases} \frac{1}{2} + \rho_i & \text{if } j = i \\ \frac{1}{2} - \rho_i & \text{if } j \neq i \end{cases}$$

is the transition matrix of the Markov chain (η_n) , and $\rho_i \in (-\frac{1}{2}, \frac{1}{2})$ are fixed parameters. For this random walk, the additive structure described in Section 3.1 is particularly simply expressed via $X_n = X_0 + \sum_{k=0}^{n-1} \eta_{k+1}$.

Corresponding to q is the stationary distribution $\pi(i) = \frac{(1/2) - \rho_{-i}}{1 - \rho_i - \rho_{-i}}$, and the mean drifts given by (3.2) are now $\mu_1(i) = \sum_{j \in S} jq(i, j) = 2i\rho_i$. Then we see that the ‘pseudo-drift’ (3.3) is zero if and only if $\rho_i = \rho$ is the same for each i ; the random walk is recurrent in exactly this case.

A positive ρ_i corresponds to *persistence* of the walker in direction i (the walker has an ‘inertia’); a negative ρ_i corresponds to a walker who *vacillates* in direction i , and has an increased propensity to turn around.

Such models have a long history, and have been studied under different names by many different researchers: as ‘persistent random walks’ by Fürth [11], ‘correlated random walks’ by Gillis [12], ‘random walks with restricted reversals’ by Domb and Fisher [5], and, recently, ‘Newtonian random walks’ by Lenci [22]. Under appropriate rescaling, the model leads to the *telegrapher’s equation* in the scaling limit, as discussed by Goldstein [13] and Kac [17]. There has been a large amount of recent work on correlated random walk and related models; a small selection is [1,28,3,15]. Motivation for studying these models arises from several sources, including physical Brownian motion [11] and models for molecular configurations [4]. We refer to [16] for some additional background and references.

As an application of our main results, consider the following variation on the one-dimensional correlated random walk, intended to probe more precisely the recurrence–transience phase transition. This time we take the state-space to be $\mathbb{Z}_+ \times \{-1, +1\}$ to fit into the setting of Section 2. We suppose that the nonzero transition probabilities are $p(x, i, x + j, j) = q_x(i, j)$, where

$$q_x(i, j) = \begin{cases} \frac{1}{2} + \frac{c_i}{2x} + O(x^{-1-\delta}) & \text{if } j = i \\ \frac{1}{2} - \frac{c_i}{2x} + O(x^{-1-\delta}) & \text{if } j \neq i \end{cases}$$

for some constants $\delta > 0$ and $c_i \in \mathbb{R}$. For $c_i > 0$, the walk is persistent in direction i , and for general c_i the symmetry between the two directions present in the recurrent case is broken: how does this affect the recurrence?

Under these assumptions, (Q_∞^+) holds with $q(i, j) = 1/2$ for all i, j , so that $\pi(i) = 1/2$ for $i = \pm 1$. Also,

$$\begin{aligned} \mu_1(x, i) &= \sum_{j \in S} jq_x(i, j) = \frac{ic_i}{x} + O(x^{-1-\delta}), \quad \text{and} \\ \mu_2(x, i) &= \sum_{j \in S} j^2q_x(i, j) = 1, \quad \text{for } x \geq 1, \end{aligned}$$

so that (M_L^+) holds (with $s_i^2 = 1$ for $i = \pm 1$). Applying Theorems 2.5 and 2.6 yields the following result (where we write c_+ for c_{+1} and c_- for c_{-1}).

Corollary 3.1. *If $c_+ - c_- < -1$, then the walk is positive-recurrent. If $c_+ - c_- > 1$, then the walk is transient. If $|c_+ - c_-| \leq 1$, then the walk is null-recurrent. Moreover, if $c_+ - c_- > -1$, then*

$$\lim_{n \rightarrow \infty} \mathbb{P}[n^{-1/2}X_n \leq x, \eta_n = i] = \frac{1}{2}F_{(c_+ - c_- + 1)/2, 2}(x).$$

3.3. Modulated queue

To finish this section we return to the queueing model as presented in the introduction. Recall that the critical case from the point of view of recurrence and transience is when $\rho = \lambda$, and we are interested in the behaviour of the model under perturbations of the constants c_i for $i \in S$. For this model we have $q_x(i, j) = (a_{ij} + b_{ij})/2 + O(x^{-1})$ so provided that the matrix $M_{ij} = (a_{ij} + b_{ij})/2$ is irreducible, condition (Q_∞^+) holds. We see that

$$\mu_1(x, i) = \frac{c_i}{x} + O(x^{-2}), \quad \text{and} \quad \mu_2(x, i) = 1,$$

so that (M_L^+) holds. Let π be the stationary distribution associated with transition matrix M , and set $\bar{c} = \sum_{i \in S} c_i \pi(i)$. Applying Theorems 2.5 and 2.6 yields the following result (cf. Corollary 3.1).

Corollary 3.2. *If $\bar{c} < -1/2$, then the Markov chain is positive-recurrent. If $\bar{c} > 1/2$, then the Markov chain is transient. If $|\bar{c}| \leq 1/2$, then the Markov chain is null-recurrent. Moreover, if $\bar{c} > -1/2$, then*

$$\lim_{n \rightarrow \infty} \mathbb{P}[n^{-1/2}X_n \leq x, \eta_n = i] = \frac{1}{2}F_{\bar{c} + (1/2), 2}(x).$$

4. Analysis via an embedded Markov chain

4.1. Overview

To analyse (X_n, η_n) we look at an embedded process (Y_n) , which records the X -coordinate of the chain when it returns to a given line. Formally, we label an arbitrary state $0 \in S$. Then set $\tau_0 = \min\{n \in \mathbb{Z}_+ : \eta_n = 0\}$, and for $m \geq 0$ set $\tau_{m+1} = \min\{n > \tau_m : \eta_n = 0\}$, where we adopt the usual convention that $\min \emptyset = \infty$. To ease exposition, we introduce a ‘coffin’ state ∂ and define the embedded process Y_n on $\mathbb{Z}_+ \cup \{\partial\}$ by

$$Y_n = \begin{cases} X_{\tau_n} & \text{if } \tau_n < \infty, \\ \partial & \text{if } \tau_n = \infty. \end{cases}$$

We also introduce $\tau = \min\{n > 0 : \eta_n = 0\}$ (so $\tau = \tau_0 \mathbf{1}\{\eta_0 \neq 0\} + \tau_1 \mathbf{1}\{\eta_0 = 0\}$).

For any $n \in \mathbb{Z}_+$, given $\tau_n < \infty$ and $X_{\tau_n} = x$, the strong Markov property for the time-homogeneous Markov chain (X_n, η_n) shows that $(X_{\tau_n+m}, \eta_{\tau_n+m})_{m \geq 0}$ is independent of $(X_0, \eta_0), \dots, (X_{\tau_n}, \eta_{\tau_n})$ and is distributed as a copy of $(X_m, \eta_m)_{m \geq 0}$ given $(X_0, \eta_0) = (x, 0)$. In particular, on $\tau_{n+1} < \infty$, the pair $(X_{\tau_{n+1}}, \tau_{n+1} - \tau_n)$ depends on $(X_0, \eta_0), \dots, (X_{\tau_n}, \eta_{\tau_n})$ only through X_{τ_n} . Hence Y_n is a Markov chain and, given $Y_n = x$, the random variable $\tau_{n+1} - \tau_n$ has the same distribution as τ conditional on $(X_0, \eta_0) = (x, 0)$.

We refer to $(X_m, \eta_m)_{\tau_n \leq m \leq \tau_{n+1}}$ as the n th excursion from the line 0. The basis for our analysis of the embedded Markov chain (Y_n) will be an analysis of a single excursion, depending on the

starting position. A key component of this analysis is a coupling result, which we present in the next subsection.

4.2. Coupling construction

Lemma 4.1. *Suppose that condition (B_p) holds for some $p > 0$ and condition (Q_∞) holds. Then there exists a Markov chain (X_n, η_n, η_n^*) on $\mathbb{Z}_+ \times S \times S$ such that*

- (X_n, η_n) is a Markov chain on $\mathbb{Z}_+ \times S$ with transition probabilities $p(x, i, y, j)$;
- (η_n^*) is a Markov chain on S with transition probabilities $q(i, j)$; and
- for all $n \in \mathbb{Z}_+$ and all $i \in S$,

$$\lim_{x \rightarrow \infty} \mathbb{P} \left[\bigcap_{0 \leq k \leq n} \{\eta_k = \eta_k^*\} \mid X_0 = x, \eta_0 = \eta_0^* = i \right] = 1. \tag{4.1}$$

Finally, suppose in addition that (Q_∞^+) holds. Then there exists $\delta > 0$ such that, for any $A < \infty$, for all $i \in S$, as $x \rightarrow \infty$,

$$1 - \mathbb{P} \left[\bigcap_{0 \leq k \leq A \log x} \{\eta_k = \eta_k^*\} \mid X_0 = x, \eta_0 = \eta_0^* = i \right] = O(x^{-\delta}). \tag{4.2}$$

The statements of Lemma 4.1 will follow from a coupling argument. Essentially, Eq. (4.1) is proved using a maximal coupling of η_n and η_n^* ; the condition (Q_∞) that $q_x(i, j)$ has a limit as $x \rightarrow \infty$ means that we can control the probability of decoupling, provided that X_n stays sufficiently large, and it is this dependence on X_n that introduces a (minor) complication to an otherwise standard argument. Eq. (4.2) is proved in a similar manner using the stronger condition (Q_∞^+) on $q_x(i, j)$; the full details of the proof can be found in the Appendix.

In the remainder of this subsection we explore some consequences of the coupling described in Lemma 4.1. First we introduce additional notation in the context of the joint probability space on which the coupled process (X_n, η_n, η_n^*) is constructed. We denote by τ^* the first return time to 0 of the Markov chain (η_n^*) , namely

$$\tau^* := \min\{n \geq 1 : \eta_n^* = 0\}.$$

Moreover, we write $\mathbb{P}_{x,i,j}$ for the probability measure conditional on $X_0 = x, \eta_0 = i, \eta_0^* = j$, and $\mathbb{E}_{x,i,j}$ for the corresponding expectation.

Irreducibility of the time-homogeneous Markov chain (X_n, η_n) and finiteness of S imply that for any x , there exist $m(x) < \infty$ and $\varphi(x) > 0$ such that

$$\mathbb{P}_{x,i}[\tau \leq m(x)] \geq \varphi(x) \quad \text{for all } i. \tag{4.3}$$

In the specific case that $q_x(i, j)$ is constant in x , the process (η_n) is distributed exactly as the finite irreducible Markov chain (η_n^*) , so the functions $m(x)$ and $\varphi(x)$ in (4.3) can be chosen to be uniform over x . Our first consequence of the above coupling is that (4.3) can be strengthened to such a uniform version under our weaker conditions: roughly speaking, assumption (Q_∞) implies that (η_n) is sufficiently close to (η_n^*) when the X -coordinate of (X_n, η_n) is sufficiently large, and irreducibility does the rest.

Lemma 4.2. *Suppose that condition (B_p) holds for some $p > 0$ and condition (Q_∞) holds. Then there exist $m < \infty$ and $\varphi > 0$ such that, for all i and all x ,*

$$\mathbb{P}_{x,i}[\tau \leq m] \geq \varphi. \tag{4.4}$$

In the proof of this result, and at several points later on, we consider the event

$$E_n := \bigcap_{0 \leq \ell \leq n} \{\eta_\ell = \eta_\ell^*\}. \tag{4.5}$$

Proof of Lemma 4.2. We work with the Markov chain (X_n, η_n, η_n^*) given in Lemma 4.1. Since η^* is a finite irreducible Markov chain, there exist $m < \infty$ and $\varphi > 0$ such that $\mathbb{P}_{x,i,i}[\tau^* \leq m] \geq 2\varphi$ for all i and all x . Conditional on η_n and η_n^* remaining coupled up to time m , we have $\tau \leq m$ if and only if $\tau^* \leq m$; hence

$$\mathbb{P}_{x,i,i}[\tau \leq m] \geq \mathbb{P}_{x,i,i}[E_m \cap \{\tau^* \leq m\}] \geq \mathbb{P}_{x,i,i}[\tau^* \leq m] - \mathbb{P}_{x,i,i}[E_m^c].$$

But by Lemma 4.1, there exists x_0 such that $\mathbb{P}_{x,i,i}[E_m^c] \leq \varphi$ for all $x \geq x_0$ and hence (4.4) holds for all i and all $x \geq x_0$.

But also, since $x_0 < \infty$, for any $m(x)$ and $\varphi(x)$ satisfying (4.3), we define $m_0 := \max_{x \leq x_0} m(x) < \infty$ and $\varphi_0 := \min_{x \leq x_0} \varphi(x) > 0$ so that, for any $x \leq x_0$,

$$\mathbb{P}_{x,i}[\tau \leq m_0] \geq \mathbb{P}_{x,i}[\tau \leq m(x)] \geq \varphi(x) \geq \varphi_0, \quad \text{for all } i.$$

So, redefining m and φ as necessary, (4.4) in fact holds for all i and all x . \square

4.3. Excursion durations and occupation estimates

Next we give an exponential tail bound for the duration of excursions, uniform in the initial location.

Lemma 4.3. *Suppose that condition (B_p) holds for some $p > 0$ and condition (Q_∞) holds. Then there exist constants $c > 0$ and $C < \infty$ such that, for all x, n , and r ,*

$$\mathbb{P}[\tau_{n+1} - \tau_n > r \mid X_{\tau_n} = x] \leq Ce^{-cr}.$$

Proof. Recall that since $\tau_{n+1} - \tau_n$ conditional on $Y_n = x$ has the same distribution as τ conditional on $X_0 = x, \eta_0 = 0$, it suffices to show that, for some constants $C, c > 0$,

$$\mathbb{P}_{x,i}[\tau > r] \leq Ce^{-cr}, \quad \text{for all } x \text{ and } i. \tag{4.6}$$

(We then get the claimed result for $\tau_{n+1} - \tau_n$ by setting $i = 0$.) Recall that, by Lemma 4.2, $\mathbb{P}_{x,i}[\tau \leq m] \geq \varphi$. Moreover, using the time-homogeneity of (X_n, η_n) , for all x and i ,

$$\begin{aligned} \mathbb{P}_{x,i}[\tau \leq km + m \mid \tau > km] &\geq \min_{y,j} \mathbb{P}[\tau \leq km + m \mid \tau > km, X_{km} = y, \eta_{km} = j] \\ &= \min_{y,j} \mathbb{P}[\tau \leq m \mid X_0 = y, \eta_0 = j] \geq \varphi, \end{aligned}$$

for all positive integers k . But this implies that, for all positive integers k ,

$$\mathbb{P}_{x,i}[\tau > km] = \prod_{j=1}^k \mathbb{P}_{x,i}[\tau > jm \mid \tau > (j-1)m] \leq (1-\varphi)^k.$$

Finally, for general $r \in \mathbb{Z}_+$, there exists an integer k such that $km \leq r < (k+1)m$, so

$$\mathbb{P}_{x,i}[\tau > r] \leq \mathbb{P}_{x,i}[\tau > km] \leq (1-\varphi)^k < (1-\varphi)^{r/m-1} \leq Ce^{-cr},$$

for constants $C, c > 0$ dependent only on φ and m , giving (4.6). \square

The next result shows that the mean occupation time of (X_n, η_n) on line i per excursion can be approximated by the mean occupation time of (η_n^*) in state i per excursion.

Lemma 4.4. *Suppose that condition (B_p) holds for some $p > 0$ and condition (Q_∞) holds. Then, for any $i \in S$,*

$$\lim_{x \rightarrow \infty} \mathbb{E}_{x,0} \sum_{k=0}^{\tau-1} \mathbf{1}\{\eta_k = i\} = \frac{\pi(i)}{\pi(0)}.$$

If, in addition, (Q_∞^+) holds, then there exists $\delta > 0$ such that, for any $i \in S$, as $x \rightarrow \infty$,

$$\left| \mathbb{E}_{x,0} \sum_{k=0}^{\tau-1} \mathbf{1}\{\eta_k = i\} - \frac{\pi(i)}{\pi(0)} \right| = O(x^{-\delta}).$$

Proof. Again we work with the Markov chain (X_n, η_n, η_n^*) whose existence is given in the statement of Lemma 4.1. Fix $i \in S$. For the duration of this proof, we write

$$W := \sum_{k=0}^{\tau-1} \mathbf{1}\{\eta_k = i\}, \quad \text{and} \quad W^* := \sum_{k=0}^{\tau^*-1} \mathbf{1}\{\eta_k^* = i\}.$$

Since (η_n^*) is a Markov chain on S with transition probabilities $q(i, j)$, standard Markov chain theory yields $\mathbb{E}_{x,0,0}[W^*] = \pi(i)/\pi(0)$, for any $x \in \mathbb{Z}_+$. The statements of the lemma will follow from suitable estimates for $\mathbb{E}_{x,0,0}[|W - W^*|]$.

Again define E_n by (4.5). Then, for any positive integer n ,

$$\begin{aligned} \mathbb{E}_{x,0,0}[|W - W^*|] &\leq \mathbb{E}_{x,0,0}[|W - W^*| \mathbf{1}(E_n) \mathbf{1}\{\tau \vee \tau^* \leq n\}] \\ &\quad + \mathbb{E}_{x,0,0}[|W - W^*| \mathbf{1}(E_n^c) \mathbf{1}\{\tau \vee \tau^* \leq n\}] \\ &\quad + \mathbb{E}_{x,0,0}[|W - W^*| \mathbf{1}\{\tau \vee \tau^* > n\}] \\ &\leq 0 + n\mathbb{P}_{x,0,0}[E_n^c] + \mathbb{E}_{x,0,0}[(\tau \vee \tau^*) \mathbf{1}\{\tau \vee \tau^* > n\}]. \end{aligned}$$

Moreover,

$$\mathbb{E}_{x,0,0}[(\tau \vee \tau^*) \mathbf{1}\{\tau \vee \tau^* > n\}] \leq \mathbb{E}_{x,0,0}[\tau \mathbf{1}\{\tau > n\}] + \mathbb{E}_{x,0,0}[\tau^* \mathbf{1}\{\tau^* > n\}]. \tag{4.7}$$

Here, by Cauchy–Schwarz and the tail estimates in Lemma 4.3,

$$\mathbb{E}_{x,0,0}[\tau \mathbf{1}\{\tau > n\}] \leq (\mathbb{E}_{x,0,0}[\tau^2])^{1/2} (\mathbb{P}_{x,0,0}[\tau > n])^{1/2} \leq C e^{-cn}, \tag{4.8}$$

for some constants $C < \infty$ and $c > 0$, not depending on x , and similarly for the term involving τ^* . For the first statement in the lemma, it suffices to show that

$$\lim_{x \rightarrow \infty} \mathbb{E}_{x,0,0}[|W - W^*|] = 0. \tag{4.9}$$

Under assumption (Q_∞) , it follows from (4.8) and its analogue for τ^* that for any $\varepsilon > 0$ we may choose $n \geq n_0$ sufficiently large so that the right-hand side of (4.7) is less than ε , and then $\mathbb{E}_{x,0,0}[|W - W^*|] \leq n\mathbb{P}_{x,0,0}[E_n^c] + \varepsilon$. For fixed n , $\mathbb{P}_{x,0,0}[E_n^c] \rightarrow 0$ as $x \rightarrow \infty$ by (4.1), so that $\limsup_{x \rightarrow \infty} \mathbb{E}_{x,0,0}[|W - W^*|] \leq \varepsilon$. Since $\varepsilon > 0$ was arbitrary, (4.9) follows.

For the second statement in the lemma, under assumption (Q_∞^+) , we use a similar argument but with $n = n(x) = \lfloor A \log x \rfloor$. As before,

$$\mathbb{E}_{x,0,0}[|W - W^*|] \leq n(x)\mathbb{P}_{x,0,0}[E_{n(x)}^c] + \mathbb{E}_{x,0,0}[(\tau \vee \tau^*) \mathbf{1}\{\tau \vee \tau^* > n(x)\}].$$

For a sufficiently large choice of constant A , the exponential bound (4.8) shows that the right-hand side of (4.7) decays as a power of x , for $n = n(x)$. Finally, the term $n(x)\mathbb{P}_{x,0,0}[E_{n(x)}^c]$ also decays as a power of x , by (4.2), and so we see that $\mathbb{E}_{x,0,0}[|W - W^*|]$ decays as a power of x , as required. \square

4.4. Recurrence and transience relationships

In this subsection we demonstrate the equivalence of recurrence properties of the embedded process (Y_n) to those of the process (X_n) .

From this point of the paper onwards, we will be increasingly concerned with multiple excursions, and it is useful to introduce the notation $\sigma_0 := 0$ and, for $n \in \mathbb{Z}_+$,

$$\sigma_{n+1} := \tau_{n+1} - \tau_n$$

for the durations of the excursions. Recall the definition of Y_n from Section 4.1. Under our conditions (cf. Lemma 4.3), $\sigma_n < \infty$ a.s. for each n . Hence $Y_n \neq \partial$ a.s., and we can identify Y_n with X_{τ_n} for all n . For the remainder of the paper we employ this slight abuse of notation, and assume that the state space of (Y_n) is \mathbb{Z}_+ . The next result relates recurrence of (X_n) to recurrence of (Y_n) .

Lemma 4.5. *Suppose that condition (B_p) holds for some $p > 0$ and condition (Q_∞) holds. Then the process (Y_n) is an irreducible Markov chain (on \mathbb{Z}_+). Moreover*

- (i) (X_n) is recurrent if and only if (Y_n) is recurrent.
- (ii) (X_n) is positive-recurrent if and only if (Y_n) is positive-recurrent.

Proof. As explained in Section 4.1, the fact that (Y_n) is a Markov chain follows from the strong Markov property for (X_n, η_n) .

Irreducibility of (Y_n) follows from the irreducibility of (X_n, η_n) , as follows. For any $x, y \in \mathbb{Z}_+$, there exists a finite path in the state space $\mathbb{Z}_+ \times S$ from $(x, 0)$ to $(y, 0)$ that the chain (X_n, η_n) has a positive probability of following. But then the (finite) subpath consisting of the points that are on line 0 corresponds to a path in the state space \mathbb{Z}_+ that (Y_n) has a positive probability of following.

Now, for statement (i), the fact that $Y_n = 0$ exactly when $X_{\tau_n} = 0$ implies $\{Y_n = 0 \text{ i.o.}\}$ if and only if $\{(X_n, \eta_n) = (0, 0) \text{ i.o.}\}$, so (Y_n) is recurrent if and only if (X_n, η_n) is recurrent. Using Lemma 2.1, we have (Y_n) is recurrent if and only if (X_n) is recurrent.

Finally, we verify (ii). Let

$$\xi = \min\{n \geq 1 : Y_n = 0\}, \quad \text{and} \quad \zeta = \min\{n \geq 1 : (X_n, \eta_n) = (0, 0)\}.$$

Then (Y_n) is positive-recurrent if and only if $\mathbb{E}_{x,0}\xi < \infty$ for some (hence all) x , while (X_n, η_n) is positive-recurrent if and only if $\mathbb{E}_{x,0}\zeta < \infty$. However, ξ and ζ are related since, given $\eta_0 = 0$, it is the case that $\tau_0 = 0$ and $\zeta = \tau_\xi$, i.e.,

$$\zeta = \sum_{k=0}^{\xi-1} \sigma_{k+1} = \sum_{k=0}^{\infty} \sigma_{k+1} \mathbf{1}\{k < \xi\}. \tag{4.10}$$

In particular, (4.10) shows that $\zeta \geq \xi$, a.s., so $\mathbb{E}_{x,0}\zeta < \infty$ implies that $\mathbb{E}_{x,0}\xi < \infty$. For the implication in the other direction, take expectations in the final expression in (4.10) and use

linearity of expectations and Fubini’s Theorem to get

$$\begin{aligned} \mathbb{E}_{x,0}\zeta &= \mathbb{E}_{x,0} \sum_{k=0}^{\infty} \mathbb{E}[\sigma_{k+1} \mathbf{1}\{k < \xi\} \mid \mathcal{F}_{\tau_k}] \\ &= \mathbb{E}_{x,0} \sum_{k=0}^{\infty} \mathbf{1}\{k < \xi\} \mathbb{E}[\sigma_{k+1} \mid \mathcal{F}_{\tau_k}], \end{aligned}$$

since $\{k < \xi\} \in \mathcal{F}_{\tau_k}$. But, by Lemma 4.3, $\mathbb{E}[\sigma_{k+1} \mid \mathcal{F}_{\tau_k}]$ is uniformly bounded by a constant, C , say, so that

$$\mathbb{E}_{x,0}\zeta \leq C \mathbb{E}_{x,0} \sum_{k=0}^{\infty} \mathbf{1}\{k < \xi\} = C \mathbb{E}_{x,0}\xi.$$

Hence $\mathbb{E}_{x,0}\zeta < \infty$ if and only if $\mathbb{E}_{x,0}\xi < \infty$. Finally, (ii) follows from Lemma 2.2, which gives the equivalence of positive-recurrence for (X_n, η_n) and (X_n) . \square

4.5. Increment moment estimates

So far, we have studied the excursions of (X_n, η_n) away from the line $\eta_n = 0$ in terms of the η -coordinate. The next stage is to study the behaviour, over an excursion, of the X -coordinate. In particular, we estimate the moments of $Y_{n+1} - Y_n$, with a view to later applying a Lamperti condition to determine the recurrence/transience of (Y_n) . First, we need estimates on the maximum deviation of X_n during a single excursion:

$$D_n := \max_{\tau_n \leq m \leq \tau_{n+1}} |X_m - X_{\tau_n}|; \tag{4.11}$$

note that the distribution of D_n given $X_{\tau_n} = x$ depends only on x and not on n .

Lemma 4.6. *Suppose that condition (B_p) holds for some $p > 0$ and condition (Q_∞) holds. Then, for any $q \in (0, p)$,*

$$\sup_x \mathbb{P}[D_n \geq d \mid X_{\tau_n} = x] = O(d^{-q}), \quad \text{and} \quad \sup_x \mathbb{E}[D_n^q \mid X_{\tau_n} = x] < \infty.$$

Proof. We have

$$\begin{aligned} \mathbb{P}[D_n \geq d \mid X_{\tau_n} = x] &\leq \mathbb{P}[\sigma_{n+1} \geq y \mid X_{\tau_n} = x] + \mathbb{P}[D_n \geq d, \sigma_{n+1} < y \mid X_{\tau_n} = x] \\ &\leq C e^{-cy} + \mathbb{P}\left[\max_{\tau_n \leq m \leq \tau_{n+y}} |X_m - X_{\tau_n}| \geq d \mid X_{\tau_n} = x\right], \end{aligned}$$

for all $d \geq 0$ and $y > 0$, by Lemma 4.3. Here,

$$\begin{aligned} &\mathbb{P}\left[\max_{\tau_n \leq m \leq \tau_{n+y}} |X_m - X_{\tau_n}| \geq d \mid X_{\tau_n} = x\right] \\ &\leq \mathbb{P}\left[\max_{\tau_n \leq m \leq \tau_{n+y}} \sum_{\ell=\tau_n}^{m-1} |X_{\ell+1} - X_\ell| \geq d \mid X_{\tau_n} = x\right] \\ &\leq \mathbb{P}\left[\bigcup_{\tau_n \leq \ell \leq \tau_{n+y}-1} \left\{|X_{\ell+1} - X_\ell| \geq \frac{d}{y}\right\} \mid X_{\tau_n} = x\right] \\ &\leq y C_p \left(\frac{d}{y}\right)^{-p}, \end{aligned}$$

which follows from the inequalities of Boole and Markov and the fact that

$$\begin{aligned} &\mathbb{E}[|X_{\ell+1} - X_{\ell}|^p \mid X_{\tau_n} = x] \\ &= \sum_{z,i} \mathbb{E}[|X_{\ell+1} - X_{\ell}|^p \mid X_{\ell} = z, \eta_{\ell} = i] \mathbb{P}[X_{\ell} = z, \eta_{\ell} = i \mid X_{\tau_n} = x] \leq C_p, \end{aligned}$$

by assumption (B_p) . Then, taking $y = d^{(p-q)/(1+p)}$, where $q \in (0, p)$, we obtain $\mathbb{P}[D_n \geq d \mid X_{\tau_n} = x] = O(d^{-q})$, as claimed. The final claim follows from the fact that

$$\mathbb{E}[D_n^{\alpha} \mid X_{\tau_n} = x] = \sum_{d=1}^{\infty} \mathbb{P}[D_n^{\alpha} \geq d \mid X_{\tau_n} = x] \leq \int_0^{\infty} \mathbb{P}[D_n^{\alpha} \geq t \mid X_{\tau_n} = x] dt,$$

which is finite when $\alpha \in (0, q)$, where q can be arbitrarily close to p . \square

We are now in a position to calculate the moments of $Y_{n+1} - Y_n$. The first case to consider is when, for each i , $\mu(x, i)$ is asymptotically d_i .

Lemma 4.7. *Suppose that condition (B_p) holds for some $p > 1$, and conditions (Q_{∞}) and (M_C) hold. Then there exists $\varepsilon > 0$ such that*

$$\sup_x \mathbb{E}[|Y_{n+1} - Y_n|^{1+\varepsilon} \mid Y_n = x] < \infty. \tag{4.12}$$

Also, as $x \rightarrow \infty$,

$$\mathbb{E}[Y_{n+1} - Y_n \mid Y_n = x] = \frac{1}{\pi(0)} \sum_{i \in S} d_i \pi(i) + o(1). \tag{4.13}$$

Proof. First, note that $|Y_{n+1} - Y_n| = |X_{\tau_{n+1}} - X_{\tau_n}| \leq |D_n|$, a.s., where D_n is given by (4.11). Then the statement (4.12) follows from Lemma 4.6 with (B_p) for $p > 1$.

It remains to prove (4.13); by the time-homogeneity of (X_n, η_n) and since $Y_n = X_{\tau_n}$, it suffices to consider $\mathbb{E}_{x,0}[X_{\tau} - X_0]$. The Doob decomposition for X_n is

$$X_n - X_0 = M_n + \sum_{k=0}^{n-1} \mathbb{E}[X_{k+1} - X_k \mid X_k, \eta_k],$$

where M_n is a martingale with $M_0 = 0$. Hence, by definition of $\mu_1(x, i)$,

$$X_n - X_0 = M_n + \sum_{k=0}^{n-1} \mu_1(X_k, \eta_k) = M_n + \sum_{i \in S} \sum_{k=0}^{n-1} \mu_1(X_k, i) \mathbf{1}\{\eta_k = i\}.$$

Since $\mathbb{E}\tau < \infty$, and $\mathbb{E}[|M_{n+1} - M_n| \mid \mathcal{F}_n] \leq 2\mathbb{E}[|X_{n+1} - X_n| \mid \mathcal{F}_n] \leq 2C_1$, a.s., (by the $p = 1$ case of (B_p)), the Optional Stopping Theorem gives $\mathbb{E}M_{\tau} = M_0 = 0$. Therefore,

$$\mathbb{E}_{x,0}[X_{\tau} - X_0] = \sum_{i \in S} \mathbb{E}_{x,0} \left[\sum_{k=0}^{\tau-1} \mu_1(X_k, i) \mathbf{1}\{\eta_k = i\} \right]. \tag{4.14}$$

Now, let $D = \max_{0 \leq k \leq \tau} |X_k - X_0|$, and set $A_x = \{D < x^{\gamma}\}$, for some $\gamma \in (0, 1)$. Note that, conditional on $X_0 = x$ and $\eta_0 = 0$, the random variable D has the same distribution as the random variable D_n defined at (4.11) given $X_{\tau_n} = x$, so by Lemma 4.6 we have

$$\mathbb{P}_{x,0}[A_x^c] = \mathbb{P}_{x,0}[D \geq x^{\gamma}] = O(x^{-\gamma}). \tag{4.15}$$

Now, given $X_0 = x$ and A_x , we have for all $0 \leq k \leq \tau$ that $X_k \geq x - x^\gamma \geq x/2$, say, for all x sufficiently large. Thus, by (M_C) , for any $\theta > 0$, there exists $x_0 < \infty$ such that, given $X_0 = x \geq x_0$,

$$\max_{i \in S} \max_{0 \leq k \leq \tau} |\mu_1(X_k, i) - d_i| \mathbf{1}(A_x) \leq \theta, \quad \text{a.s.}$$

Since $\max_{x,i} |\mu_1(x, i)| < \infty$ and $\max_i |d_i| < \infty$, it follows that there exists a constant $C < \infty$ such that, given $X_0 = x \geq x_0$,

$$\max_{i \in S} \max_{0 \leq k \leq \tau} |\mu_1(X_k, i) - d_i| \leq \theta + C \mathbf{1}(A_x^c), \quad \text{a.s.}$$

Hence, given $X_0 = x \geq x_0$,

$$\left| \sum_{k=0}^{\tau-1} \mu_1(X_k, i) \mathbf{1}\{\eta_k = i\} - \sum_{k=0}^{\tau-1} d_i \mathbf{1}\{\eta_k = i\} \right| \leq \theta \tau + C \tau \mathbf{1}(A_x^c), \quad \text{a.s.}$$

Here, by the Cauchy–Schwarz inequality,

$$\mathbb{E}_{x,0}[\tau \mathbf{1}(A_x^c)] \leq (\mathbb{E}_{x,0}[\tau^2])^{1/2} (\mathbb{P}_{x,0}[A_x^c])^{1/2} = O(x^{-\gamma/2}),$$

using (4.15) and the fact that τ has all moments, by Lemma 4.3. So, for any $\delta > 0$, we can choose $x_1 < \infty$ sufficiently large so that, given $X_0 = x \geq x_1$,

$$\max_i \left| \mathbb{E}_{x,0} \left[\sum_{k=0}^{\tau-1} \mu_1(X_k, i) \mathbf{1}\{\eta_k = i\} \right] - \mathbb{E}_{x,0} \left[\sum_{k=0}^{\tau-1} d_i \mathbf{1}\{\eta_k = i\} \right] \right| \leq \delta.$$

Together with Lemma 4.4 and (4.14) this yields (4.13). \square

Lemma 4.8. *Suppose that condition (B_p) holds for some $p > 2$, and conditions (Q_∞) and (M_L) hold. Then there exists $\varepsilon > 0$ such that*

$$\sup_x \mathbb{E}[|Y_{n+1} - Y_n|^{2+\varepsilon} | Y_n = x] < \infty. \tag{4.16}$$

Also, as $x \rightarrow \infty$,

$$\mathbb{E}[Y_{n+1} - Y_n | Y_n = x] = \frac{1}{\pi(0)} \sum_{i \in S} \frac{c_i \pi(i)}{x} + o(x^{-1}); \tag{4.17}$$

$$\mathbb{E}[(Y_{n+1} - Y_n)^2 | Y_n = x] = \frac{1}{\pi(0)} \sum_{i \in S} s_i^2 \pi(i) + o(1). \tag{4.18}$$

If, in addition (Q_∞^+) and (M_L^+) hold, then there exists $\delta > 0$ such that

$$\mathbb{E}[Y_{n+1} - Y_n | Y_n = x] = \frac{1}{\pi(0)} \sum_{i \in S} \frac{c_i \pi(i)}{x} + O(x^{-1-\delta}); \tag{4.19}$$

$$\mathbb{E}[(Y_{n+1} - Y_n)^2 | Y_n = x] = \frac{1}{\pi(0)} \sum_{i \in S} s_i^2 \pi(i) + O(x^{-\delta}); \tag{4.20}$$

Proof. First, since $|Y_{n+1} - Y_n| \leq D_n$, with D_n as defined at (4.11), and because Lemma 4.6 implies that $\sup_x \mathbb{E}[(D_n)^{2+\varepsilon} | X_{\tau_n} = x] < \infty$, (4.16) follows.

The proof of (4.17) and (4.18) using (Q_∞) and (M_L) and the proof of (4.19) and (4.20) using (Q_∞^+) and (M_L^+) are essentially the same, the only difference being in the error terms associated to each expression. We present the proof of (4.19) and (4.20); it should be clear how to adapt the argument to prove (4.17) and (4.18).

We proceed as in the proof of Lemma 4.7. Indeed, we follow the reasoning from the second paragraph of that proof through to Eq. (4.14), giving

$$\mathbb{E}_{x,0}[X_\tau - X_0] = \sum_{i \in S} \mathbb{E}_{x,0} \left[\sum_{k=0}^{\tau-1} \mu_1(X_k, i) \mathbf{1}\{\eta_k = i\} \right],$$

and we let $D = \max_{0 \leq k \leq \tau} |X_k - X_0|$, and set $A_x = \{D < x^\gamma\}$ as before, but now we require $\gamma \in (1/2, 1)$. Note that, conditional on $X_0 = x$ and $\eta_0 = 0$, the random variable D has the same distribution as the random variable D_n defined at (4.11) given $X_{\tau_n} = x$, so by Lemma 4.6 we have that $\mathbb{P}_{x,0}[D \geq d] = O(d^{-p'})$ for some $p' > 2$ since τ has all moments and (B_p) holds for some $p > 2$.

Now, given $X_0 = x$ and A_x , we have $|X_k - x| \leq D < x^\gamma$ for $k \leq \tau$, so that, by (M_L^+) ,

$$\left| \mu_1(X_k, i) - \frac{c_i}{x} \right| \leq \left| \frac{c_i}{X_k} - \frac{c_i}{x} \right| + O((x - x^\gamma)^{-1-\delta_1}) = O(x^{\gamma-2}) + O(x^{-1-\delta_1}),$$

uniformly for $0 \leq k \leq \tau$. Therefore $\mu_1(X_k, i) \mathbf{1}(A_x) = (c_i/x + O(x^{\gamma-2}) + O(x^{-1-\delta_1})) \mathbf{1}(A_x)$, which means that $\mu_1(X_k, i) = c_i/x + O(x^{\gamma-2}) + O(x^{-1-\delta_1}) + O(1) \mathbf{1}(A_x^c)$. So,

$$\begin{aligned} & \mathbb{E}_{x,0} \left[\sum_{k=0}^{\tau-1} \mu_1(X_k, i) \mathbf{1}\{\eta_k = i\} \right] \\ &= \mathbb{E}_{x,0} \left[\left(\frac{c_i}{x} + O(x^{\gamma-2}) + O(x^{-1-\delta_1}) + O(1) \mathbf{1}(A_x^c) \right) \sum_{k=0}^{\tau-1} \mathbf{1}\{\eta_k = i\} \right], \end{aligned}$$

where the implicit constants are uniform in x and in i . By (Q_∞^+) and the second statement in Lemma 4.4, we have that

$$\mathbb{E}_{x,0} \left[\sum_{k=0}^{\tau-1} \mathbf{1}\{\eta_k = i\} \right] = \frac{\pi(i)}{\pi(0)} + O(x^{-\delta'}), \tag{4.21}$$

for some $\delta' > 0$, so

$$\begin{aligned} \mathbb{E}_{x,0}[X_\tau - X_0] &= \frac{1}{\pi(0)} \sum_{i \in S} \frac{c_i \pi(i)}{x} + O(x^{-1-\delta'}) + O(x^{\gamma-2}) + O(x^{-1-\delta_1}) \\ &\quad + O(1) \mathbb{E}_{x,0}[\tau \mathbf{1}(A_x^c)]. \end{aligned}$$

Here, by Hölder’s inequality, for all $r, s > 0$ with $r^{-1} + s^{-1} = 1$,

$$\mathbb{E}_{x,0}[\tau \mathbf{1}(A_x^c)] \leq (\mathbb{E}_{x,0}[\tau^r])^{1/r} (\mathbb{P}_{x,0}[A_x^c])^{1/s}.$$

Since τ has all moments, we can take $s = p'/2 > 1$, so that $\mathbb{E}_{x,0}[\tau \mathbf{1}(A_x^c)] = O(x^{-2\gamma})$. Then, since $\gamma \in (1/2, 1)$, $\delta_1 > 0$ and $\delta' > 0$ we have, for some $\delta'' > 0$,

$$\mathbb{E}[Y_{n+1} - Y_n \mid Y_n = x] = \frac{1}{\pi(0)} \sum_{i \in S} \frac{c_i \pi(i)}{x} + O(x^{-1-\delta''}).$$

To calculate the second moment of $X_\tau - X_0$, we will make repeated use of the algebraic identity $a^2 - b^2 = (a - b)^2 + 2b(a - b)$, which will help to simplify the calculations that follow. Taking the Doob decomposition for X_n^2 , we write

$$\begin{aligned} X_n^2 - X_0^2 &= M_n + \sum_{k=0}^{n-1} \mathbb{E}[X_{k+1}^2 - X_k^2 \mid X_k, \eta_k] \\ &= M_n + \sum_{k=0}^{n-1} \left(\mathbb{E}[(X_{k+1} - X_k)^2 \mid X_k, \eta_k] + 2X_k \mathbb{E}[X_{k+1} - X_k \mid X_k, \eta_k] \right) \\ &= M_n + \sum_{k=0}^{n-1} (\mu_2(X_k, \eta_k) + 2X_k \mu_1(X_k, \eta_k)) \\ &= M_n + \sum_{i \in S} \sum_{k=0}^{n-1} (s_i^2 + 2c_i + O(X_k^{-\delta_1})) \mathbf{1}\{\eta_k = i\}, \end{aligned}$$

by (M_L^+) , where M_n is a martingale satisfying $M_0 = 0$. Moreover, given $X_0 = x$,

$$\begin{aligned} |M_{n \wedge \tau}| &\leq |X_{n \wedge \tau}^2 - X_0^2| + C\tau \\ &= (X_{n \wedge \tau} - X_0)^2 + 2X_0|X_{n \wedge \tau} - X_0| + C\tau \\ &\leq D^2 + 2xD + C\tau, \end{aligned}$$

where $D = \max_{0 \leq k \leq \tau} |X_k - X_0|$ is as defined earlier, and $C < \infty$ is a constant. Thus, $M_{n \wedge \tau}$ is uniformly integrable (in n) and so by the Optional Stopping Theorem $\mathbb{E}M_\tau = M_0 = 0$. Therefore,

$$\mathbb{E}_{x,0}[X_\tau^2 - X_0^2] = \sum_{i \in S} (s_i^2 + 2c_i) \mathbb{E}_{x,0} \left[\sum_{k=0}^{\tau-1} \mathbf{1}\{\eta_k = i\} \right] + \mathbb{E}_{x,0} \left[\sum_{k=0}^{\tau-1} O(X_k^{-\delta_1}) \right].$$

As in the calculation of the first moment, we can bound the error term by bootstrapping on the event A_x : writing $O(X_k^{-\delta_1}) = O(x^{-\delta_1}) + O(1)\mathbf{1}(A_x^c)$, we get

$$\begin{aligned} \mathbb{E}_{x,0} \left[\sum_{k=0}^{\tau-1} O(X_k^{-\delta_1}) \right] &= O(x^{-\delta_1}) + O(1)\mathbb{E}_{x,0}[\tau \mathbf{1}(A_x^c)] \\ &= O(x^{-\delta_1}) + O(x^{-2\gamma}), \end{aligned}$$

as above, and therefore, by (4.21),

$$\mathbb{E}_{x,0}[X_\tau^2 - X_0^2] = \frac{1}{\pi(0)} \sum_{i \in S} (s_i^2 + 2c_i)\pi(i) + O(x^{-\delta'}) + O(x^{-\delta_1}) + O(x^{-2\gamma}).$$

Now we use $X_\tau^2 - X_0^2 = (X_\tau - X_0)^2 + 2X_0(X_\tau - X_0)$ to get

$$\mathbb{E}_{x,0}[(X_\tau - X_0)^2] = \frac{1}{\pi(0)} \sum_{i \in S} s_i^2 \pi(i) + O(x^{-\delta''}),$$

for some $\delta''' > 0$. Finally, taking $\delta = \min\{\delta'', \delta'''\}$ yields (4.19) and (4.20), as required. \square

5. Proofs of main results

5.1. Recurrence classification

To prove Theorems 2.4 and 2.5, we use the increment moment estimates from Section 4.5 together with some Foster–Lamperti conditions to classify the process (Y_n) , and then deduce the classification for (X_n) from the equivalence results in Section 4.4.

For Theorem 2.5, under Lamperti-type drift assumptions, we apply the following classification result.

Lemma 5.1 (Lamperti). *Let (Z_n) be an irreducible time-homogeneous Markov chain on \mathbb{Z}_+ . Suppose that there exists $\varepsilon > 0$ such that*

$$\sup_z \mathbb{E}[|Z_{n+1} - Z_n|^{2+\varepsilon} \mid Z_n = z] < \infty; \tag{5.1}$$

$$\liminf_{z \rightarrow \infty} \mathbb{E}[|Z_{n+1} - Z_n|^2 \mid Z_n = z] > 0. \tag{5.2}$$

Let $\mu_k(z) = \mathbb{E}[(Z_{n+1} - Z_n)^k \mid Z_n = z]$.

- If $\liminf_{z \rightarrow \infty} (2z\mu_1(z) - \mu_2(z)) > 0$, then Z_n is transient.
- If $|2z\mu_1(z)| \leq \mu_2(z) + O(z^{-\delta})$, for some $\delta > 0$, then Z_n is null-recurrent.
- If $\limsup_{z \rightarrow \infty} (2z\mu_1(z) + \mu_2(z)) < 0$, then Z_n is positive-recurrent.

Lemma 5.1 is essentially due to Lamperti [19,21], although the form given here is taken from Menshikov et al. [25, Theorem 3]. The conditions for recurrence and transience are contained in Theorem 3.2 of [19], and the condition for positive-recurrence is contained in Theorem 2.1 of [21]. The condition for null-recurrence here is slightly sharper than Lamperti’s original results [21].

Proof of Theorem 2.5. We apply Lemma 5.1 to classify $Z_n = Y_n$, and thus, by Lemma 4.5, classify X_n . First, assuming (B_p) for some $p > 2$, (Q_∞) and (M_L) , by Lemma 4.8 it is clear that (5.1) and (5.2) hold for $Z_n = Y_n$. Furthermore,

$$\begin{aligned} \liminf_{x \rightarrow \infty} 2x\mathbb{E}[Y_{n+1} - Y_n \mid Y_n = x] &= \limsup_{x \rightarrow \infty} 2x\mathbb{E}[Y_{n+1} - Y_n \mid Y_n = x] \\ &= \frac{1}{\pi(0)} \sum_{i \in S} 2c_i \pi(i), \end{aligned}$$

and

$$\liminf_{x \rightarrow \infty} \mathbb{E}[|Y_{n+1} - Y_n|^2 \mid Y_n = x] = \limsup_{x \rightarrow \infty} \mathbb{E}[|Y_{n+1} - Y_n|^2 \mid Y_n = x] = \frac{1}{\pi(0)} \sum_{i \in S} s_i^2 \pi(i).$$

By Lemma 5.1, $\sum_{i \in S} (2c_i - s_i^2)\pi(i) > 0$ implies transience, while $\sum_{i \in S} (2c_i + s_i^2)\pi(i) < 0$ implies positive-recurrence. When $|\sum_{i \in S} 2c_i \pi(i)| < \sum_{i \in S} s_i^2 \pi(i)$, we have

$$\lim_{x \rightarrow \infty} (|2x\mathbb{E}[Y_{n+1} - Y_n \mid Y_n = x]| - \mathbb{E}[|Y_{n+1} - Y_n|^2 \mid Y_n = x]) < 0,$$

which means the middle condition of Lemma 5.1 holds for any $\delta > 0$, and therefore Y_n is null-recurrent.

Now suppose that (Q_∞^+) and (M_L^+) also hold. Then, by Lemma 4.8, we have

$$\begin{aligned} & |2x\mathbb{E}[Y_{n+1} - Y_n \mid Y_n = x]| - \mathbb{E}[|Y_{n+1} - Y_n|^2 \mid Y_n = x] \\ &= \frac{1}{\pi(0)} \left(\left| \sum_{i \in S} 2c_i \pi(i) \right| - \sum_{i \in S} s_i^2 \pi(i) \right) + O(x^{-\delta}) \end{aligned}$$

for some $\delta > 0$, which means that $|\sum_{i \in S} 2c_i \pi(i)| = \sum_{i \in S} s_i^2 \pi(i)$ implies that (Y_n) is null-recurrent, completing the classification of (Y_n) and therefore of (X_n) . \square

For Theorem 2.4 we will apply the following condition.

Lemma 5.2. *Let (Z_n) be an irreducible time-homogeneous Markov chain on \mathbb{Z}_+ . For (Z_n) to be transient, it is sufficient that there exists $\varepsilon > 0$ such that*

$$\begin{aligned} & \sup_z \mathbb{E}[|Z_{n+1} - Z_n|^{1+\varepsilon} \mid Z_n = z] < \infty, \quad \text{and} \\ & \liminf_{z \rightarrow \infty} \mathbb{E}[Z_{n+1} - Z_n \mid Z_n = z] > 0. \end{aligned}$$

We omit the proof of Lemma 5.2, which is similar to the proof of Lemma 5.1 and relies on demonstrating the existence of a suitable Lyapunov function with negative drift outside a bounded set, using Taylor’s formula and some careful truncation.

Proof of Theorem 2.4. Consider the Markov chain (Y_n) . Under the conditions of part (i) of the theorem, Lemma 4.7 implies that the hypotheses of Lemma 5.2 hold for $Z_n = Y_n$, so that (Y_n) is transient. Hence, by Lemma 4.5, (X_n) is also transient.

As mentioned after the statement, part (ii) was obtained by Falin [7]. Our results furnish a different proof: Lemma 4.7 gives positive-recurrence for (Y_n) by Foster’s criterion (e.g. Theorem 2.2.3 of [9]), so, by Lemma 4.5, (X_n) is also positive-recurrent. \square

5.2. Convergence in distribution

The first step in the proof of Theorem 2.6 is to apply a result of Lamperti [20] to obtain a weak limit for the embedded Markov chain (Y_n) . Recall the distribution function $F_{\alpha,\theta}$ as defined at (2.3).

Lemma 5.3. *Suppose (X_n, η_n) is a Markov chain satisfying (B_p) for some $p > 4$, (Q_∞) and (M_L) . Suppose that the matrix q appearing in (Q_∞) is aperiodic. Suppose also that $\sum_{i \in S} (2c_i + s_i^2) \pi(i) > 0$. Define α and θ as at (2.4). Then, for any $x \in \mathbb{R}_+$,*

$$\lim_{n \rightarrow \infty} \mathbb{P} \left[n^{-1/2} Y_n \leq x \right] = F_{\alpha,\theta}(x \sqrt{\pi(0)}).$$

Proof. If (B_p) holds for some $p > 4$, then a consequence of Lemma 4.6 is that

$$\sup_x \mathbb{E} \left[|Y_{n+1} - Y_n|^4 \mid Y_n = x \right] < \infty.$$

Now we apply Theorem 2.1 of [20] to the Markov chain (Y_n) , using the increment moment estimates of Lemma 4.8 and noting the remark preceding the theorem in [20], to obtain

$$\lim_{n \rightarrow \infty} \mathbb{P} \left[n^{-1/2} Y_n \leq x \right] = F_{\alpha,\theta/\pi(0)}(x) = F_{\alpha,\theta}(x \sqrt{\pi(0)}),$$

where the final equality follows from (2.3). \square

Remark 5.4. If in addition (Q_∞^+) and (M_L^+) hold, then in the case $\sum_{i \in S} (2c_i + s_i^2)\pi(i) = 0$ it follows from Lemma 2.1 of [20] that $n^{-1/2}Y_n \rightarrow 0$ in probability; cf. Remark 2.7(iii).

The next goal is to deduce from the weak limit for Y_n a weak limit for X_n . To do so, we need (i) to control the value of the process (X_n) between successive observations of the embedded process, and (ii) to account for the change of time. First we address point (i). For each $n \in \mathbb{Z}_+$, let $N(n) := \max\{k : \tau_k \leq n\}$, so that $\tau_{N(n)} \leq n < \tau_{N(n)+1}$.

Lemma 5.5. *Suppose that (X_n, η_n) satisfies (B_p) for some $p > 2$ and (Q_∞) . Then, as $n \rightarrow \infty$, $n^{-1/2}|X_n - X_{\tau_{N(n)}}| \rightarrow 0$ in probability.*

Proof. Since $\sigma_k \geq 1$, we have that $N(n) \leq n$, a.s. Hence $|X_n - X_{\tau_{N(n)}}| \leq \max_{k \leq n} D_k$, where D_k is as defined at (4.11). Thus it suffices to show that $n^{-1/2} \max_{k \leq n} D_k \rightarrow 0$ in probability. For any $\gamma > 0$, we have

$$\begin{aligned} \max_{k \leq n} D_k &\leq n^\gamma + \max_{k \leq n} (D_k \mathbf{1}\{D_k > n^\gamma\}) \\ &\leq n^\gamma + \sum_{k=0}^n D_k \mathbf{1}\{D_k > n^\gamma\}. \end{aligned}$$

Since (B_p) holds for $p > 2$, Lemma 4.6 shows $\max_k \mathbb{E}[D_k^q] < \infty$ for some $q > 2$, so

$$\mathbb{E} [D_k \mathbf{1}\{D_k > n^\gamma\}] = \mathbb{E} \left[D_k^q D_k^{1-q} \mathbf{1}\{D_k > n^\gamma\} \right] \leq n^{-\gamma(q-1)} \mathbb{E} [D_k^q].$$

It follows that

$$n^{-1/2} \mathbb{E} \max_{k \leq n} D_k = n^{\gamma-(1/2)} + O(n^{(1/2)-\gamma(q-1)}),$$

which is $o(1)$ provided we choose (as we may) $\frac{1}{2(q-1)} < \gamma < \frac{1}{2}$. Thus $n^{-1/2} \max_{k \leq n} D_k \rightarrow 0$ in L^1 , and hence in probability. \square

Next we turn to point (ii) mentioned above. For our purposes, the following renewal-type result will suffice.

Lemma 5.6. *Suppose (X_n, η_n) is a Markov chain satisfying (B_p) for some $p > 2$, (Q_∞) and (M_L) . Suppose also that $\sum_{i \in S} (2c_i + s_i^2)\pi(i) > 0$.*

Then, as $n \rightarrow \infty$, $n^{-1}N(n) \rightarrow \pi(0)$ in probability.

Proof. Under the conditions of the lemma, Theorem 2.5 shows that X_n (and hence Y_n) is null, i.e., null-recurrent or transient. In particular, for any $x \geq 0$,

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\frac{1}{n} \sum_{k=0}^{n-1} \mathbf{1}\{X_{\tau_k} \leq x\} \right] = 0. \tag{5.3}$$

We use an extension of the coupling given in Lemma 4.1 to multiple excursions. We construct on the same probability space (X_n, η_n) together with a sequence $(\eta_{k,n}^*)$ of copies (for $k \in \mathbb{Z}_+$) of the Markov chain (η_n^*) as follows. At each $\tau_k, k \in \mathbb{Z}_+$, start $(\eta_{k,n}^*)_{n \geq 0}$, an independent copy of $(\eta_n^*)_{n \geq 0}$, from $\eta_{k,0}^* = \eta_{\tau_k} = 0 \in S$, coupled to $(\eta_n)_{n \geq \tau_k}$ as described in Lemma 4.1; denote by σ_{k+1}^* the number of steps until $\eta_{k,n}^*$ returns to 0.

Extending the notation E_n defined at (4.5), we write $E_{k,n} = \cap_{0 \leq \ell \leq n} \{\eta_{\tau_k + \ell} = \eta_{k,\ell}^*\}$, the event that the coupling started at τ_k succeeds for n steps.

Now we use this coupling construction and the null property (5.3) to show that $n^{-1}\tau_n \rightarrow \pi(0)^{-1}$ in probability. For $s > 0$, denote $\chi_s(x) := x\mathbf{1}\{x \leq s\}$. Note that

$$\left| \frac{1}{n} \sum_{k=0}^{n-1} \chi_s(\sigma_{k+1}) - \frac{1}{n} \sum_{k=0}^{n-1} \sigma_{k+1} \right| \leq \frac{1}{n} \sum_{k=0}^{n-1} \sigma_{k+1} \mathbf{1}\{\sigma_{k+1} > s\}.$$

Here $\mathbb{E}[\sigma_{k+1} \mathbf{1}\{\sigma_{k+1} > s\}] \leq s^{-1} \mathbb{E}[\sigma_{k+1}^2]$, say, so that, by Lemma 4.3,

$$\lim_{s \rightarrow \infty} \sup_k \mathbb{E}[\sigma_{k+1} \mathbf{1}\{\sigma_{k+1} > s\}] = 0.$$

A similar argument holds for σ_{k+1}^* . Hence, for any $\varepsilon > 0$, there exists $s_0 < \infty$ such that

$$\mathbb{E} \left[\left| \left(\frac{1}{n} \sum_{k=0}^{n-1} \sigma_{k+1} - \frac{1}{n} \sum_{k=0}^{n-1} \sigma_{k+1}^* \right) - \left(\frac{1}{n} \sum_{k=0}^{n-1} \chi_s(\sigma_{k+1}) - \frac{1}{n} \sum_{k=0}^{n-1} \chi_s(\sigma_{k+1}^*) \right) \right| \right] \leq \varepsilon, \tag{5.4}$$

for all $s \geq s_0$ and all n . On the event $E_{k,s}$ (the coupling started at τ_k succeeds for s steps) we have $\chi_s(\sigma_{k+1}) = \chi_s(\sigma_{k+1}^*)$. Then, for any $x > 0$,

$$|\chi_s(\sigma_{k+1}) - \chi_s(\sigma_{k+1}^*)| \leq s \mathbf{1}(E_{k,s}^c) \mathbf{1}\{X_{\tau_k} > x\} + s \mathbf{1}\{X_{\tau_k} \leq x\}.$$

Now

$$\begin{aligned} \mathbb{P}[E_{k,s}^c \cap \{X_{\tau_k} > x\}] &\leq \sup_{y>x} \mathbb{P}[E_{k,s}^c \mid X_{\tau_k} = y, \eta_{\tau_k} = \eta_{k,0}^* = 0] \\ &= \sup_{y>x} \mathbb{P}[E_s^c \mid X_0 = y, \eta_0 = \eta_0^* = 0]. \end{aligned}$$

So for fixed $s \geq s_0$, Lemma 4.1 shows we may choose $x \geq x_0$ large enough such that,

$$\mathbb{E} \frac{1}{n} \sum_{k=0}^{n-1} s \mathbf{1}(E_{k,s}^c) \mathbf{1}\{X_{\tau_k} > x\} \leq \varepsilon,$$

for all n . Combining this with the null property (5.3), we obtain that, for fixed $s \geq s_0$,

$$\limsup_{n \rightarrow \infty} \mathbb{E} \left[\left| \frac{1}{n} \sum_{k=0}^{n-1} \chi_s(\sigma_{k+1}) - \sum_{k=0}^{n-1} \chi_s(\sigma_{k+1}^*) \right| \right] \leq \varepsilon.$$

Thus with (5.4) we conclude that

$$\limsup_{n \rightarrow \infty} \mathbb{E} \left| \frac{1}{n} \sum_{k=0}^{n-1} \sigma_{k+1} - \frac{1}{n} \sum_{k=0}^{n-1} \sigma_{k+1}^* \right| \leq 2\varepsilon.$$

Since $\varepsilon > 0$ was arbitrary, and σ_{k+1}^* are i.i.d. random variables with mean $\pi(0)^{-1}$, it follows that $n^{-1}\tau_n \rightarrow \pi(0)^{-1}$ in probability.

The claimed result now follows by inverting the law of large numbers: for example,

$$\mathbb{P} \left[n^{-1}N(n) > \pi(0) + \varepsilon \right] \leq \mathbb{P} \left[\tau_{\lceil (\pi(0) + \varepsilon)n \rceil} \leq n \right] \leq \mathbb{P} \left[\frac{\tau_{\lceil (\pi(0) + \varepsilon)n \rceil}}{\lceil (\pi(0) + \varepsilon)n \rceil} \leq \frac{1}{\pi(0) + \varepsilon} \right],$$

which tends to 0 as $n \rightarrow \infty$ for any $\varepsilon > 0$; similarly in the other direction. \square

In the proof of Theorem 2.6 we will use two facts about convergence in distribution that we now recall (see e.g. [6, p. 73]). First, if sequences of random variables ξ_n and ζ_n are such that

$\zeta_n \rightarrow \zeta$ in distribution for some random variable ζ and $|\xi_n - \zeta_n| \rightarrow 0$ in probability, then $\xi_n \rightarrow \zeta$ in distribution (this is *Slutsky’s theorem*). Second, if $\zeta_n \rightarrow \zeta$ in distribution and $\alpha_n \rightarrow \alpha$ in probability, then $\alpha_n \zeta_n \rightarrow \alpha \zeta$ in distribution.

Proof of Theorem 2.6. First, since $n^{-1}N(n) \rightarrow \pi(0)$ in probability (Lemma 5.6),

$$\lim_{n \rightarrow \infty} \mathbb{P} \left[\frac{X_{\tau_{N(n)}}}{\sqrt{N(n)}} \cdot \sqrt{\frac{N(n)}{n}} \leq x \right] = \lim_{n \rightarrow \infty} \mathbb{P} \left[\frac{X_{\tau_{N(n)}}}{\sqrt{N(n)}} \leq \frac{x}{\sqrt{\pi(0)}} \right] = F_{\alpha, \theta}(x),$$

by Lemma 5.3 and the fact that $\lim_{n \rightarrow \infty} N(n) = \infty$ a.s. Together with Lemma 5.5 and Slutsky’s theorem, this shows that

$$\lim_{n \rightarrow \infty} \mathbb{P}[n^{-1/2}X_n \leq x] = \lim_{n \rightarrow \infty} \mathbb{P}[n^{-1/2}X_{\tau_{N(n)}} \leq x] = F_{\alpha, \theta}(x). \tag{5.5}$$

Next we prove the joint convergence of (X_n, η_n) . For $m \in \mathbb{Z}_+$, let $R_{n,m} = n^{-1/2}|X_{n-m} - X_n|$. Then, by the $p = 1$ case of (B_p) , we have $\mathbb{E}[R_{n,m}] \leq Cmn^{-1/2}$ for some finite constant C . Hence, for fixed m , as $n \rightarrow \infty$, $R_{n,m} \rightarrow 0$ in L^1 and hence in probability.

Fix $x \in (0, \infty)$. Then, for any $\varepsilon \in (0, x)$,

$$\mathbb{P}[n^{-1/2}X_n > x, \eta_n = k] \leq \mathbb{P}[n^{-1/2}X_{n-m} > x - \varepsilon, \eta_n = k] + \mathbb{P}[R_{n,m} \geq \varepsilon].$$

Here

$$\begin{aligned} &\mathbb{P}[n^{-1/2}X_{n-m} > x - \varepsilon, \eta_n = k] \\ &= \sum_{y: n^{-1/2}y > x - \varepsilon} \mathbb{P}[X_{n-m} = y] \mathbb{P}[\eta_n = k \mid X_{n-m} = y]. \end{aligned} \tag{5.6}$$

Again we use the coupling of Lemma 4.1 and the notation E_n from (4.5). Note that

$$|\mathbb{P}_{y,i,i}[\eta_m = k] - \pi(k)| \leq \mathbb{P}_{y,i,i}[E_m^c] + |\mathbb{P}_{y,i,i}[\eta_m^* = k] - \pi(k)|.$$

Here, since (η_m^*) is an aperiodic, irreducible finite Markov chain with stationary distribution π , $\mathbb{P}_{y,i,i}[\eta_m^* = k] = \mathbb{P}[\eta_m^* = k \mid \eta_0^* = i]$ converges (uniformly over i and y) to $\pi(k)$ as $m \rightarrow \infty$. So, for any $\delta > 0$, we may choose $m_0 < \infty$ such that, for all i and all y ,

$$|\mathbb{P}_{y,i,i}[\eta_{m_0} = k] - \pi(k)| \leq \mathbb{P}_{y,i,i}[E_{m_0}^c] + \delta.$$

By Lemma 4.1, we may then choose $y_0 < \infty$ large enough so that, for all $y \geq y_0$,

$$|\mathbb{P}[\eta_{m_0} = k \mid X_0 = y] - \pi(k)| \leq 2\delta.$$

Now taking n large enough so that $(x - \varepsilon)n^{1/2} > y_0$, it follows from (5.6) that

$$\mathbb{P}[n^{-1/2}X_n > x, \eta_n = k] \leq \mathbb{P}[R_{n,m_0} \geq \varepsilon] + (\pi(k) + 2\delta)\mathbb{P}[n^{-1/2}X_{n-m_0} > x - \varepsilon].$$

We now let $n \rightarrow \infty$ and apply (5.5) to obtain

$$\limsup_{n \rightarrow \infty} \mathbb{P}[n^{-1/2}X_n > x, \eta_n = k] \leq (\pi(k) + 2\delta) (1 - F_{\alpha, \theta}(x - \varepsilon)).$$

Since $\varepsilon > 0$ and $\delta > 0$ were arbitrary, and $F_{\alpha, \theta}$ is continuous, it follows that

$$\limsup_{n \rightarrow \infty} \mathbb{P}[n^{-1/2}X_n > x, \eta_n = k] \leq \pi(k) (1 - F_{\alpha, \theta}(x)), \quad \text{for all } x \in (0, \infty).$$

A similar argument in the other direction, starting from the inequality

$$\mathbb{P}[n^{-1/2}X_n > x, \eta_n = k] \geq \mathbb{P}[n^{-1/2}X_{n-m} > x + \varepsilon, \eta_n = k] - \mathbb{P}[R_{n,m} \geq \varepsilon]$$

yields the complementary lim inf statement, so that

$$\lim_{n \rightarrow \infty} \mathbb{P}[n^{-1/2} X_n > x, \eta_n = k] = \pi(k) (1 - F_{\alpha, \theta}(x)), \quad \text{for all } x \in (0, \infty). \tag{5.7}$$

The statement in the theorem now follows from the fact that, by (5.7),

$$\lim_{n \rightarrow \infty} \mathbb{P}[n^{-1/2} X_n \leq x, \eta_n = k] = \lim_{n \rightarrow \infty} \mathbb{P}[\eta_n = k] - \pi(k) (1 - F_{\alpha, \theta}(x)),$$

where $\lim_{n \rightarrow \infty} \mathbb{P}[\eta_n = k] = \pi(k)$ by taking $x \downarrow 0$ in (5.7). \square

Acknowledgements

This work was supported by the Engineering and Physical Sciences Research Council (grant number EP/J021784/1).

We thank an anonymous referee for his/her careful reading and useful comments.

Appendix. Proof of coupling lemma

In this appendix we give the deferred technical proof of our coupling result, Lemma 4.1.

Proof of Lemma 4.1. As commented on earlier, the proof follows an almost standard coupling argument. Indeed, since the first two statements of the lemma will be satisfied for any coupling of (X_n, η_n) and (η_n^*) on a common probability space, in order to also prove (4.1)/(4.2) it makes sense to use a maximal coupling of η_n and η_n^* , which we will construct in a step-wise fashion. For us, the condition that $q_x(i, j)$ has a limit as $x \rightarrow \infty$ means that the probability of decoupling at any step will be small, provided that X_n stays sufficiently large. This introduces some complications to the standard coupling arguments, as we will need to keep control of the variation of X_n .

We construct the Markov chain (X_n, η_n, η_n^*) by describing a single step:

- If $\eta_n \neq \eta_n^*$ then produce (X_{n+1}, η_{n+1}) from (X_n, η_n) according to the transition probabilities $p(x, i, y, j)$, and produce η_{n+1}^* from η_n^* independently according to the transition probabilities $q(i, j)$.
- Otherwise, given $\eta_n = \eta_n^* = i$ and $X_n = x$, we use a maximal coupling (see, for example, Lindvall [23, pp. 18–20]) to produce $(\eta_{n+1}, \eta_{n+1}^*)$ via

$$\mathbb{P}[\eta_{n+1} = j, \eta_{n+1}^* = k] = \begin{cases} \min\{q_x(i, j), q(i, k)\} & \text{for } j = k, \\ \frac{(q_x(i, j) - q(i, j))^+ (q(i, k) - q_x(i, k))^+}{\frac{1}{2} \sum_{\ell \in S} |q_x(i, \ell) - q(i, \ell)|} & \text{for } j \neq k. \end{cases}$$

Then, given $\eta_{n+1} = j$ we produce X_{n+1} via

$$\mathbb{P}[X_{n+1} = y \mid \eta_{n+1} = j] = \frac{p(x, i, y, j)}{\sum_{z \in \mathbb{Z}_+} p(x, i, z, j)}.$$

It is a simple matter to check that we have constructed a valid coupling of (X_n, η_n) and η_n^* . Indeed, making use of the fact that

$$\sum_{\ell \in S} |q_x(i, \ell) - q(i, \ell)| = \sum_{\ell \in S} (q_x(i, \ell) - q(i, \ell))^+ + \sum_{\ell \in S} (q(i, \ell) - q_x(i, \ell))^+$$

and

$$0 = \sum_{\ell \in S} (q_x(i, \ell) - q(i, \ell)) = \sum_{\ell \in S} (q_x(i, \ell) - q(i, \ell))^+ - \sum_{\ell \in S} (q(i, \ell) - q_x(i, \ell))^+,$$

calculation shows that $\mathbb{P}[\eta_{n+1} = j \mid (X_n, \eta_n) = (x, i)] = q_x(i, j)$ and $\mathbb{P}[\eta_{n+1}^* = j \mid \eta_n^* = i] = q(i, j)$. Then we see that

$$\mathbb{P}[(X_{n+1}, \eta_{n+1}) = (y, j) \mid (X_n, \eta_n) = (x, i)] = \frac{p(x, i, y, j)}{\sum_{z \in \mathbb{Z}_+} p(x, i, z, j)} q_x(i, j) = p(x, i, y, j).$$

This verifies the coupling construction. Note that, with this coupling,

$$\mathbb{P}[\eta_{n+1} \neq \eta_{n+1}^* \mid X_n = x, \eta_n = \eta_n^* = i] = \frac{1}{2} \sum_{j \in S} |q_x(i, j) - q(i, j)|. \tag{A.1}$$

It remains to prove (4.1) and (4.2). First in the case of (4.1), for which we assume (Q_∞) , we give the argument in detail; we will then indicate how to modify the argument to prove (4.2).

Given $\varepsilon > 0$ and $n < \infty$, choose x_0 so that $\max_i \sum_{j \in S} |q_x(i, j) - q(i, j)| \leq \frac{\varepsilon}{n}$ for all $x \geq x_0$; this is possible by assumption (Q_∞) .

Let $A_k = \{X_k \geq x_0\}$, and recall from (4.5) that $E_k = \cap_{0 \leq \ell \leq k} \{\eta_\ell = \eta_\ell^*\}$. Then,

$$\begin{aligned} \mathbb{P}[E_{k+1}^c \mid E_k \cap A_k] &= \mathbb{P}[\eta_{k+1} \neq \eta_{k+1}^* \mid E_k \cap A_k] \\ &\leq \max_i \sup_{x \geq x_0} \mathbb{P}[\eta_{k+1} \neq \eta_{k+1}^* \mid X_k = x, \eta_k = \eta_k^* = i], \end{aligned}$$

so that, given $X_0 = x, \eta_0 = \eta_0^* = i$,

$$\mathbb{P}[E_{k+1}^c] \leq \mathbb{P}[E_{k+1}^c \mid E_k \cap A_k] + \mathbb{P}[E_k^c] + \mathbb{P}[A_k^c] \leq \frac{\varepsilon}{2n} + \mathbb{P}[E_k^c] + \mathbb{P}[A_k^c],$$

which in turn implies that

$$\mathbb{P}[E_n^c] \leq \frac{\varepsilon}{2} + \sum_{k=0}^{n-1} \mathbb{P}[A_k^c] \leq \frac{\varepsilon}{2} + n \mathbb{P}\left[\min_{0 \leq k \leq n-1} X_k < x_0\right].$$

To complete the proof we need to show that, for x sufficiently large,

$$\mathbb{P}\left[\min_{0 \leq k \leq n-1} X_k < x_0 \mid X_0 = x, \eta_0 = \eta_0^* = i\right] \leq \frac{\varepsilon}{2n}, \quad \text{for all } i. \tag{A.2}$$

But

$$\begin{aligned} &\mathbb{P}\left[\min_{0 \leq k \leq n-1} X_k < x_0 \mid X_0 = x, \eta_0 = \eta_0^* = i\right] \\ &\leq \mathbb{P}_{x,i} \left[\max_{0 \leq k \leq n-1} |X_k - X_0| > x - x_0 \right] \\ &\leq \mathbb{P}_{x,i} \left[\bigcup_{0 \leq k \leq n-1} |X_{k+1} - X_k| > \frac{x - x_0}{n} \right] \\ &\leq n \max_{y,j} \mathbb{P}\left[|X_{k+1} - X_k| > \frac{x - x_0}{n} \mid X_k = y, \eta_k = j\right], \end{aligned}$$

so (A.2) will follow from

$$\lim_{r \rightarrow \infty} \max_{x,i} \mathbb{P}[|X_{k+1} - X_k| > r \mid X_k = x, \eta_k = i] = 0,$$

which in turn follows from condition (B_p) with $p > 0$ and Markov's inequality; indeed,

$$\begin{aligned} \max_{x,i} \mathbb{P}[|X_{k+1} - X_k| > r \mid X_k = x, \eta_k = i] &\leq \max_{x,i} \frac{\mathbb{E}[|X_{k+1} - X_k|^p \mid X_k = x, \eta_k = i]}{r^p} \\ &\leq \frac{C_p}{r^p}. \end{aligned}$$

Therefore $\mathbb{P}[E_n^c \mid X_0 = x, \eta_0 = \eta_0^* = i] \leq \varepsilon$ for all i and sufficiently large x , and since ε was arbitrary, this proves (4.1).

The proof of (4.2) is similar, now assuming (Q_∞^+) . We set $n = n(x) = \lfloor A \log x \rfloor$. Now we modify the definition of A_k to be $A_k = \{X_k \geq x/2\}$. Then, (A.1) with (Q_∞^+) gives

$$\mathbb{P}[E_{k+1}^c] \leq \mathbb{P}[E_k^c] + \mathbb{P}[A_k^c] + O(x^{-\delta_0}),$$

from which we have

$$\mathbb{P}[E_{n(x)}^c] \leq O(x^{-\delta_0/2}) + A \log x \mathbb{P}\left[\min_{0 \leq k \leq n(x)-1} X_k < x/2\right].$$

The final probability in the last display we estimate in exactly the same way as in the previous argument, replacing the previous x_0 by $x/2$ and the previous n by $n(x)$, and we again find a term that decays as a power of x . Thus we obtain (4.2). \square

References

- [1] P.C. Allaart, M.G. Monticino, Optimal stopping rules for directionally reinforced processes, *Adv. Appl. Probab.* 33 (2001) 483–504.
- [2] G. Atkinson, Recurrence of co-cycles and random walks, *J. Lond. Math. Soc.* 13 (1976) 486–488.
- [3] A.Y. Chen, E. Renshaw, The general correlated random walk, *J. Appl. Probab.* 31 (1994) 869–884.
- [4] H.E. Daniels, The statistical theory of stiff chains, *Proc. Roy. Soc. Edinburgh Sect. A* 63 (1952) 290–311.
- [5] C. Domb, M.E. Fisher, On random walks with restricted reversals, *Proc. Cambridge Philos. Soc.* 54 (1958) 48–59.
- [6] R. Durrett, *Probability: Theory and Examples*, Wadsworth & Brooks/Cole, Pacific Grove, CA, 1991.
- [7] G.I. Falin, Ergodicity of random walks in the half-strip, *Math. Notes* 44 (1988) 606–608. translated from *Mat. Zametki* 44 (1988) 225–230 (in Russian).
- [8] G. Falin, Heavy traffic analysis of a random walk on a lattice semi-strip, *Comm. Statist. Stochastic Models* 11 (1995) 395–409.
- [9] G. Fayolle, V.A. Malyshev, M.V. Menshikov, *Topics in the Constructive Theory of Countable Markov Chains*, Cambridge University Press, Cambridge, 1995.
- [10] S. Foss, S. Shneer, A. Tyurlikov, Stability of a Markov-modulated Markov chain, with application to a wireless network governed by two protocols, *Stoch. Syst.* 2 (2012) 208–231.
- [11] R. Fürth, Die Brownsche Bewegung bei Berücksichtigung einer Persistenz der Bewegungsrichtung. Mit Anwendungen auf die Bewegung lebender Infusorien, *Z. Phys.* 2 (1920) 244–256.
- [12] J. Gillis, Correlated random walk, *Proc. Cambridge Philos. Soc.* 51 (1955) 639–651.
- [13] S. Goldstein, On diffusion by discontinuous movements, and on the telegraph equation, *Quart. J. Mech.* 4 (1951) 129–156.
- [14] L. Hervé, F. Pène, On the recurrence set of planar Markov random walks, *J. Theoret. Probab.* 26 (2013) 169–197.
- [15] M. Holmes, A. Sakai, Senile reinforced random walks, *Stochastic Process. Appl.* 117 (2007) 1519–1539.
- [16] B.D. Hughes, *Random Walks and Random Environments; Volume 1: Random Walks*, Clarendon Press, Oxford, 1995.
- [17] M. Kac, A stochastic model related to the telegrapher's equation, *Rocky Mountain J. Math.* 4 (1974) 497–509.
- [18] A. Krámli, D. Szász, Random walks with internal degrees of freedom. I. Local limit theorems, *Z. Wahrscheinlichkeitstheor. Verwandte Geb.* 63 (1983) 85–95.

- [19] J. Lamperti, Criteria for the recurrence and transience of stochastic processes I, *J. Math. Anal. Appl.* 1 (1960) 314–330.
- [20] J. Lamperti, A new class of probability limit theorems, *J. Math. Mech.* 11 (1962) 749–772.
- [21] J. Lamperti, Criteria for stochastic processes II: passage-time moments, *J. Math. Anal. Appl.* 7 (1963) 127–145.
- [22] M. Lenci, Recurrence for persistent random walks in two dimensions, *Stoch. Dyn.* 7 (2007) 53–74.
- [23] T. Lindvall, *Lectures on the Coupling Method*, John Wiley & Sons, Inc., New York, 1992.
- [24] V.A. Malyshev, Homogeneous random walks on the product of finite set and a half-line, in: A.N. Kolmogorov (Ed.), *Veroyatnostnye Metody Issledovania (Probability Methods of Investigation)*, vol. 41, Moscow State University, Moscow, 1972, pp. 5–13 (in Russian).
- [25] M.V. Menshikov, I.M. Asymont, R. Iasnogorodskii, Markov processes with asymptotically zero drifts, *Probl. Inf. Transm.* 31 (1995) 248–261. translated from *Problemy Peredachi Informatsii* 31 (1995) 60–75 (in Russian).
- [26] M.F. Neuts, *Structured Stochastic Matrices of M/G/1 Type and their Applications*, Marcel Dekker, Inc., New York, 1989.
- [27] L.C.G. Rogers, Recurrence of additive functionals of Markov chains, *Sankhyā* 47 (1985) 47–56.
- [28] D. Szász, B. Tóth, Persistent random walks in a one-dimensional random environment, *J. Stat. Phys.* 37 (1984) 27–38.