

## Durham Research Online

---

### Deposited in DRO:

28 July 2014

### Version of attached file:

Published Version

### Peer-review status of attached file:

Peer-reviewed

### Citation for published item:

Hickinbottom, S and Simpson, C. J. and James, P. A. and Ibar, E. and Carter, D. and Boselli, A. and Collins, C. A. and Davies, J. I. and Dunne, L. and Eales, S. and Fuller, C. and Mobasher, B. and Peletier, R. F. and Phillipps, S. and Smith, D. J. B. and Smith, R. J. and Valentijn, E. A. (2014) 'A surprising consistency between the far-infrared galaxy luminosity functions of the field and Coma.', *Monthly notices of the Royal Astronomical Society.*, 442 (2). pp. 1286-1293.

### Further information on publisher's website:

<http://dx.doi.org/10.1093/mnras/stu924>

### Publisher's copyright statement:

This article has been accepted for publication in *Monthly Notices of the Royal Astronomical Society* © 2014 The Authors Published by Oxford University Press on behalf of the Royal Astronomical Society. All rights reserved.

### Additional information:

## Use policy

---

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

---

# A surprising consistency between the far-infrared galaxy luminosity functions of the field and Coma

S. Hickinbottom,<sup>1\*</sup> C. J. Simpson,<sup>1</sup> P. A. James,<sup>1</sup> E. Ibar,<sup>2</sup> D. Carter,<sup>1</sup> A. Boselli,<sup>3</sup>  
C. A. Collins,<sup>1</sup> J. I. Davies,<sup>4</sup> L. Dunne,<sup>5</sup> S. Eales,<sup>4</sup> C. Fuller,<sup>4</sup> B. Mobasher,<sup>6</sup>  
R. F. Peletier,<sup>7</sup> S. Phillipps,<sup>8</sup> D. J. B. Smith,<sup>9</sup> R. J. Smith<sup>10</sup> and E. A. Valentijn<sup>7</sup>

<sup>1</sup>*Astrophysics Research Institute, Liverpool John Moores University, IC2, Liverpool Science Park, 146 Brownlow Hill, Liverpool L3 5RF, UK*

<sup>2</sup>*Instituto de Física y Astronomía, Universidad de Valparaíso, Avda. Gran Bretaña 1111, 2360102 Valparaíso, Chile*

<sup>3</sup>*Aix Marseille Université, CNRS, LAM (Laboratoire d'Astrophysique de Marseille) UMR 7326, F-13388 Marseille, France*

<sup>4</sup>*School of Physics and Astronomy, Cardiff University, The Parade, Cardiff CF24 3AA, UK*

<sup>5</sup>*Department of Physics and Astronomy, University of Canterbury, Private Bag 4800, Christchurch 8140, New Zealand*

<sup>6</sup>*Department of Physics and Astronomy, University of California, Riverside, CA 92521, USA*

<sup>7</sup>*Kapteyn Astronomical Institute, University of Groningen, Postbus 800, NL-9700 AV Groningen, the Netherlands*

<sup>8</sup>*Astrophysics Group, School of Physics, University of Bristol, Bristol BS8 1TL, UK*

<sup>9</sup>*Centre for Astrophysics, Science & Technology Research Institute, University of Hertfordshire, Hatfield, Herts AL10 9AB, UK*

<sup>10</sup>*Department of Physics, Durham University, South Road, Durham DH1 3LE, UK*

Accepted 2014 May 6. Received 2014 May 6; in original form 2013 October 1

## ABSTRACT

We present new deep images of the Coma Cluster from the ESA *Herschel Space Observatory* at wavelengths of 70, 100 and 160  $\mu\text{m}$ , covering an area of  $1.75 \times 1.0$  square degrees encompassing the core and south-west infall region. Our data display an excess of sources at flux densities above 100 mJy compared to blank-field surveys, as expected. We use extensive optical spectroscopy of this region to identify cluster members and hence produce cluster luminosity functions in all three photometric bands. We compare our results to the local field galaxy luminosity function, and the luminosity functions from the *Herschel* Virgo Cluster Survey. We find consistency between the shapes of the Coma and field galaxy luminosity functions at all three wavelengths; however, we do not find the same level of agreement with that of the Virgo Cluster.

**Key words:** galaxies: clusters: individual: Coma–galaxies: luminosity function, mass function–infrared: galaxies.

## 1 INTRODUCTION

The study of galaxy clusters is important as they can provide a wealth of information regarding the process of galaxy evolution. It has been well documented that types of galaxies are strongly linked with the nature of their local environments, with a morphology–density relation being found (Dressler 1980; Whitmore, Gilmore & Jones 1993; Dressler et al. 1997; Baldry et al. 2006) such that early-type elliptical and lenticular galaxies are preferentially found in high-density cluster environments. Several processes have been identified for removing or depleting gas in cluster galaxies, such as ram-pressure stripping (Gunn & Gott 1972), ‘harassment’ through multiple high-velocity encounters (Moore, Lake & Katz 1998), ‘starvation’ through removal of large-scale weakly bound gas reservoirs (Larson, Tinsley & Caldwell 1980), rapid consumption of gas in starbursts resulting from galaxy–galaxy interactions (Joseph &

Wright 1985; Keel et al. 1985; Kennicutt et al. 1987) or interactions with the core of the cluster mass distribution (Merritt 1984; Miller 1986; Byrd & Valtonen 1990). These processes are reviewed by Boselli & Gavazzi (2006). The relative importance of the different processes is likely to be a strong function of local environment, with some only occurring in the cores of the richest clusters and others being most influential in less extreme environments, and therefore studying a range of environments is of great importance.

The Coma Cluster is a useful source of information when studying galaxy clusters as it is the richest nearby cluster, at a distance of 100 Mpc (Liu & Graham 2001). The central core regions are nearly completely virialized (Colless & Dunn 1996) and dominated by early-type galaxies. However, there is a clear substructure, the most striking example being a large group centred on NGC 4839 (Neumann et al. 2001), generally referred to as the ‘south-west infall region’. Many studies have looked at the star formation properties of Coma Cluster galaxies, with Kennicutt, Bothun & Schommer (1984) finding several galaxies showing strong  $\text{H}\alpha$  emission indicating ongoing star formation. This activity was confirmed by Caldwell

\* E-mail: [S.A.Hickinbottom@2011.ljmu.ac.uk](mailto:S.A.Hickinbottom@2011.ljmu.ac.uk)

et al. (1993), Gavazzi, Boselli & Kennicutt (1991), Gavazzi et al. (1998) and Moss & Whittle (2005), although Caldwell et al. found very few such galaxies close to the cluster core. More recently, Smith et al. (2009, 2012) have studied the importance of environment within the Coma Cluster for the quenching of star formation, showing that it is very important for lower luminosity galaxies but has little effect on the most massive and highly luminous galaxies.

An additional comparison is provided by the Virgo Cluster, a very well-studied galaxy cluster (Boselli et al. 1995) which primarily lies at a distance of 17 Mpc (Gavazzi et al. 1999). However, through the use of the GOLDMINE data base (Gavazzi et al. 2003; Gavazzi, Franzetti & Boselli 2014), one can see that there is evidence for a substructure within the cluster, with groupings at 17, 23 and 32 Mpc. The Virgo Cluster contains  $\sim 2000$  optically catalogued galaxies (Binggeli, Sandage & Tammann 1985) of both early and late types. The early types have a velocity dispersion of approximately  $589 \text{ km s}^{-1}$ , whereas the late types have a velocity dispersion of approximately  $700 \text{ km s}^{-1}$  (Binggeli, Popescu & Tammann 1993). These authors also note that the Virgo Cluster is not fully virialized, and so these values may be affected by substructures or infall velocities. For comparison, the Coma Cluster has a velocity dispersion of  $1008 \text{ km s}^{-1}$  (Struble & Rood 1999), and it is a fully virialized environment. The Virgo Cluster has been studied in detail at far-infrared wavelengths as part of the *Herschel* Virgo Cluster Survey (HeViCS; Davies et al. 2010). These observations were carried out using the ESA *Herschel Space Observatory* (Pilbratt et al. 2010), using the Photodetector Array Camera and Spectrometer (PACS; Poglitsch et al. 2010) at 100 and  $160 \mu\text{m}$ , as well as the Spectral and Photometric Imaging Receiver (SPIRE; Griffin et al. 2010) at 250, 350 and  $500 \mu\text{m}$ .

Far-infrared emission is a sensitive and powerful tracer of the evolutionary state of galaxies. For late-type galaxies, it principally traces star formation activity (Lonsdale Persson & Helou 1987; Buat & Xu 1996) with the dominant emission coming from dust thermalization and reradiation of energy from high-mass stars. In early-type galaxies, much or all of this emission is instead thought to arise from dust heated by the general radiation field of the older stellar population (Lonsdale Persson & Helou 1987; Walterbos & Greenawalt 1996), and hence far-infrared emission in these galaxies is an indicator primarily of the amount of interstellar medium they have retained. The shape of the far-infrared spectral energy distribution (SED) and the total luminosity at these wavelengths thus give insight into both the cold dust content and the star formation rate (Kennicutt 1998; Dunne et al. 2000; Kennicutt & Evans 2012). In turn, this information can improve our understanding of galaxy evolution in the cluster environment and the effect of the various galaxy interactions upon the morphology.

Initial results from HeViCS indicated a turnover at faint luminosities (Davies et al. 2010), which they interpreted as evidence for the stripping process which removes gas and dust being more effective for lower mass objects. One of the motivations for the present study is to determine the far-infrared luminosity function for the Coma Cluster, and to compare it with that for the Virgo Cluster, including the most recent analysis of the HeViCS data (Auld et al. 2013). Additionally, we will investigate how the Coma Cluster luminosity function compares with that found for field galaxies. Bai et al. (2006, 2009) investigated the Multiband Imaging Photometer for *Spitzer* (MIPS) luminosity function of the Coma Cluster, and found it to be in agreement with the infrared local field galaxy luminosity function. However, it should be noted that only a small fraction of their galaxies were detected at wavelengths longer than  $24 \mu\text{m}$ , and therefore, the total infrared luminosities were extrapolated using relationships based on optical colours.

Our adopted line of enquiry is to compare cluster galaxies with their field counterparts using the luminosity functions, thus providing an understanding of what effect the local environment has on galaxy luminosity distributions. The likely differences lie in the relative numbers of the different morphological types, as well as a deficiency of gas and dust in the cluster galaxies due to various stripping processes. This study will present results from the deepest far-infrared observations ever of the Coma Cluster, which were obtained using the ESA *Herschel Space Observatory*.<sup>1</sup> This paper will discuss the multiwavelength maps of the cluster, and an analysis will be presented of the numbers and luminosities of cluster member galaxies detected at these far-infrared wavelengths. These observations are part of a larger multiwavelength survey that was instigated with the *Hubble Space Telescope* Advanced Camera for Surveys (*HST/ACS*) Coma Cluster Survey (Carter et al. 2008).

The structure of the paper is as follows. In Section 2, the observations and the methodology of the data reduction are described. In Section 3, the analysis of the produced images is discussed along with the extraction of the sources. In Section 4, the Coma Cluster membership is determined via the use of redshifts. In Section 5, the Coma Cluster luminosity functions are derived and compared to both the field and the Virgo Cluster. A summary of the results is given in Section 6.

## 2 OBSERVATIONS AND DATA REDUCTION

The observations were carried out using the ESA *Herschel Space Observatory*, using the PACS instrument at 70, 100 and  $160 \mu\text{m}$ . The observational area was the core of the cluster and the southwest infall region, covering an area of  $1:75$  by  $1:0$ . Two scans were performed with simultaneous imaging at  $100/160 \mu\text{m}$  and two at  $70/160 \mu\text{m}$ , resulting in four separate scans being made. Within each pair, one scan was performed along the long axis of the mapped area, and one along the short axis. The scans were performed at a speed of  $20 \text{ arcsec s}^{-1}$  and the total integration time was 27.2 h, equating to an effective integration time per pixel in the final map of approximately 40 s for 70 and  $100 \mu\text{m}$  and 80 s for  $160 \mu\text{m}$ . The OBSID for the scan containing the  $70 \mu\text{m}$  maps is 1342224628/9, and for the scan containing the  $100 \mu\text{m}$  maps it is 1342233085/6.

The data were reduced using a pipeline written in `JYTHON` (a `PYTHON` implementation written in `JAVA`) that was run within the *Herschel* Interactive Pipeline Environment (HIPE; Ott 2010). The pipeline used in this work follows a similar procedure to that described in Ibar et al. (2010), but using an improved cosmic ray removal method. The data reduction consists of three main stages: flagging and calibration, deglitching and imaging. The first stage involves the masking of the bolometers that are known not to be working or are saturated (aided by housekeeping data for the day of the observations). Then the data are astrometrically calibrated by appending the pointing product to each bolometer timeline. The deglitching process involves the removal of all those read-out signals which deviate by more than  $5\sigma$  from the projected map-pixel contributions. The data are projected using `PHOTPROJECT` after removing large-scale structure from the timelines using a boxcar high-pass filter (HPF; total width of 1.6 arcmin). This filtering is essential to remove the dominant  $1/f$  noise present in the PACS timelines. The projection uses  $3.2 \text{ arcsec}$  pixel sizes for all three wavelengths. The HPF and the map projection is done iteratively in order to

<sup>1</sup> *Herschel* is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA.

mask the read-out contributions to bright pixels before high-pass filtering the timelines, hence removing the sidelobes along the scan directions produced by the boxcar HPF which are clearly seen in the projected images. This process was repeated until the projected map did not differ significantly from the previous iteration. This process results in two different types of sources detected within the map, those which were masked before producing the final version of the map, and those which were not masked and were therefore subject to the high-pass filtering. These two types of sources will be referred to as masked and unmasked sources, respectively.

### 3 DATA ANALYSIS

The data were analysed using the automatic image detection algorithm *SEXTRACTOR* (Bertin & Arnouts 1996), in order to determine the numbers and positions of sources present in each of the three maps. The values used for the detection threshold, the minimum number of contiguous pixels constituting a detected object, and the deblending contrast parameter were  $1.3\sigma$ , 6 and 0.01, respectively. An additional flux limit was imposed by requiring that each source as a whole was detected at a  $5\sigma$  level relative to the root mean square (rms) noise. The background flux level in the final maps is consistent with zero, as is to be expected due to the HPF applied to the timelines before projection. Any sources that were located at the very edges of the maps were discarded due to the increased levels of noise in those regions.

For each map, there is an optimum aperture size that maximizes the signal-to-noise ratio for point sources. For the 70, 100 and 160  $\mu\text{m}$  sources, we determined that the optimum aperture radii are 6, 7 and 10 arcsec (Altieri, private communication). The optimum aperture radius requires that the point source fluxes have a corresponding encircled energy fraction correction applied to them. The encircled energy fractions are 63.7, 64.1 and 61.9 per cent for the 70, 100 and 160  $\mu\text{m}$  point sources, respectively. The final catalogue consists of both masked and unmasked sources, and for unmasked sources this encircled energy fraction will be too large due to the removal of flux during the HPF process. Therefore, a secondary set of corrections was determined by comparing the final flux of sources with the unprocessed map flux and taking an average of the difference between them. This yielded encircled energy fractions for unmasked point sources of 59.9, 59.7 and 54.4 per cent at 70, 100 and 160  $\mu\text{m}$ , respectively.

These aperture sizes and encircled energy fractions were also used to calculate the aperture corrected rms noise values, which were found to be 5.7, 6.7 and 7.9 mJy at 70, 100 and 160  $\mu\text{m}$ , respectively.

The fluxes were measured within either the optimum aperture for point sources and corrected by the factors given in Section 2, or a larger aperture for extended sources, where extended is defined as a full width at half-maximum (FWHM) greater than  $\sqrt{2}$  multiplied by the FWHM of the PACS point spread function (PSF) at the given wavelength. This corresponds to values of 2.47, 2.96 and 4.86 pixels at 70, 100 and 160  $\mu\text{m}$ , respectively. The larger aperture was defined as having a 12.0 pixel (38.4 arcsec) radius for all maps. This value was chosen as it encircles the entirety of the most extended sources that we detect.

The completeness of the data was determined in order to ascertain how many sources had been missed during the source extraction process. Completeness curves were determined for each wavelength band by inserting 2500 fake sources, 100 at a time, into the maps before running the *SEXTRACTOR* algorithm as described above in order to see how many sources were recovered. This was done for

each brightness level between 4 mJy and 1 Jy, with intervals of 0.1 dex.

The fake source was constructed by stacking a number of bright point sources, and then scaling the flux of the object accordingly. This process was completed twice with two separate fake sources: once for masked sources, and once for unmasked sources. This was done to account for sources that were not masked at any point during the HPF process, and as such have a different PSF. The total completeness was determined for each magnitude bin by combining the masked and unmasked completeness using the following equation:

$$C = \frac{c_m c_u (n_m + n_u)}{c_u n_m + c_m n_u}. \quad (1)$$

Here,  $c_m$  and  $c_u$  are the masked and unmasked completenesses, and  $n_m$  and  $n_u$  are the number of masked and unmasked sources in the bin. This results in a 50 (80) per cent completeness of 28.5, 34.5 and 42.0 (34.4, 42.2 and 52.3) mJy for the 70, 100 and 160  $\mu\text{m}$  maps, respectively.

We detect 201, 370 and 507 sources above the  $5\sigma$  limit, in the 70, 100 and 160  $\mu\text{m}$  maps, respectively. The numbers of these confirmed sources as a function of flux in the 100 and 160  $\mu\text{m}$  maps were compared directly to results from the *Herschel* Astrophysical Terahertz Large Area Survey (H-ATLAS; Eales et al. 2010) and the PACS Evolutionary Probe (PEP; Lutz et al. 2011), as described in Rigby et al. (2011) and Berta et al. (2010). This comparison is shown in Fig. 1. The numbers of sources from the Coma data have been corrected for incompleteness, as determined via the method described earlier. We note the excess of sources at the bright end, which is expected given the presence of a rich galaxy cluster in the mapped area. The faint end of our data is consistent with previous studies to within the respective errors.


The Coma Cluster has previously been observed at far-infrared wavelengths using *IRAS*. The observations found 41 galaxies within  $4^{\circ}2$  of the cluster centre, of which 26 were confirmed to be members from their velocities (Wang et al. 1991). Of these 26 confirmed Coma Cluster members found by *IRAS*, only 4 lie within our survey region. All four sources match to the sources in our *Herschel* catalogue to within 1 arcmin, which is the approximate limiting resolution of the *IRAS* survey. There is good flux agreement between *Herschel* and *IRAS* for these four sources; however, they each have a lower flux as determined by *Herschel*, by 8–22 per cent. This is most likely due to the *IRAS* flux being contaminated by other nearby sources due to the large *IRAS* PSF.

The Coma Cluster was also surveyed to shallower depths at 100 and 160  $\mu\text{m}$  by the H-ATLAS survey (Eales et al. 2010). We matched 153 and 231 sources between the maps to within a distance equal to the optimum aperture radius. We find agreement at the level expected given the measured noise on the maps, down to the detection limit, between the fluxes of these sources down to the 100 mJy noise limit of the maps (Smith, private communication).

### 4 CLUSTER MEMBERSHIP

A number of detected sources in the maps will be foreground or background objects, and have no relation to the Coma Cluster. In order to determine which sources are true Coma Cluster members, the data were compared to the Hectospec Coma Redshift catalogue.

Data for this catalogue were obtained with the fibre multi-object spectrograph Hectospec (Fabricant et al. 2005) at the MultiMirror Telescope (now MMT) on the nights of 2007 April 12–15, with additional queue observations being made on subsequent nights. The proposal was PA-07A-0260 (PI: Ann Hornschemeier). These




**Figure 1.** A comparison of *Herschel* Coma source number counts to the H-ATLAS and PEP data at the same wavelength. The top plot shows the 100  $\mu\text{m}$  sources, whilst the bottom plot shows the 160  $\mu\text{m}$  sources. The data for the Lockman and COSMOS fields were taken from Berta et al. (2010). The GOODS field data were taken from Magnelli et al. (2013). GOODS-S refers to the deep scan of the south field only, whereas GOODS-NS is the data from a shallower scan of both the north and south fields.

observations used a 270 lines  $\text{mm}^{-1}$  grating blazed at  $\sim 5000 \text{ \AA}$  to provide a dispersion of  $1.21 \text{ \AA pixel}^{-1}$  over a useful wavelength range of 3800–8900  $\text{\AA}$ . A total of 20 fibre configurations were observed with an integration time of 1 h each and 200 fibres of each configuration were assigned to the redshift survey.

Targets for the redshift survey were selected from a parent catalogue of galaxies with Petrosian magnitudes  $r < 21.3$ , within which the higher priority targets were those with  $r < 20.3$ , those within the footprint of the *HST* Survey (Carter et al. 2008), within the *XMM-Newton* survey (Briel et al. 2001) and those identified with radio sources from the Very Large Array survey (Bravo-Alfaro et al. 2000, 2001). No colour selection criteria were applied.

The data were reduced in a standard manner, and redshifts estimated, using the Hectospec data reduction pipeline HSRED.<sup>2</sup> A second redshift estimate was derived for galaxies in the redshift survey sample using the IRAF cross-correlation task XCSAO. Each spectrum was then inspected visually by two separate members of the *HST* survey team to resolve discrepancies between HSRED and XCSAO redshifts and to assess the quality and reliability of the measured redshifts.



**Figure 2.** A plot of the 100  $\mu\text{m}$  flux against the  $r$ -band magnitude, for confirmed cluster members. The dashed and solid vertical lines indicate the limiting magnitudes of SDSS (17.77) and Hectospec (20.3), respectively. The dashed and solid horizontal lines indicate the 30 and 50 per cent completeness levels, respectively. Ball et al. (2006) determine a value of  $M_r^* = -20.49$  for their sample of galaxies, which corresponds to  $r = 14.51$ ; therefore, as our sources lie around this value, we are not observing the most massive and luminous galaxies.

The Hectospec catalogue was augmented by redshifts from the National Aeronautics and Space Administration (NASA)/IPAC extragalactic data base<sup>3</sup> and an unpublished catalogue by Colless & Dunn (private communication). The final catalogue is approximately 90 per cent complete at  $r = 19.0$ , falling to 50 per cent at  $r = 20.3$ . Incompleteness is due to the lack of spectra for some galaxies because of fibre proximity constraints, and the inability to obtain reliable redshifts for the faintest galaxies.

Any source in our *Herschel* catalogue that could be matched to a Hectospec catalogue member, to within a distance equal to the optimum aperture radius, was classified as a true Coma Cluster member; plausible larger values had no effect on the number of matches. From this process, it was determined that our data contain 50, 64 and 53 cluster members at 70, 100 and 160  $\mu\text{m}$ , respectively, with 46 sources being common to all three wavelength bands.

Fig. 2 shows the infrared flux densities of the 64 sources identified in the 100  $\mu\text{m}$  map against their  $r$ -band magnitudes taken from the Hectospec catalogue. This shows that the limiting magnitude of the Hectospec survey is unlikely to have a significant effect on the number of identifiable sources within our catalogue, and that we can assume that we have identified all Coma members within our *Herschel* sample.

If these confirmed Coma members are removed from the initial sample that was plotted in Fig. 1, and then compared to the PEP survey once again, the bright end excess is no longer present, and the trend follows that seen in the PEP counts.


## 5 LUMINOSITY FUNCTIONS

### 5.1 Coma and the field

Having produced our catalogues of cluster members, we now construct far-infrared luminosity functions at each wavelength, as shown in Fig. 3. The errors in the numbers were determined via the use of low number statistics, as detailed in Gehrels (1986).

<sup>2</sup> <http://www.astro.princeton.edu/~rcool/hsred/>

<sup>3</sup> <http://ned.ipac.caltech.edu/>



**Figure 3.** Luminosity functions at 70, 100 and 160  $\mu\text{m}$  for confirmed Coma Cluster members, shown with blue triangles, green squares and red circles, respectively.

**Table 1.** The Schechter parameter values calculated for Coma Cluster galaxies for the three *Herschel* bands.

Wavelength	$\log \phi^*$ (dex $^{-1}$ )	$\log L^*$ (W)	$\alpha$
70 $\mu\text{m}$	$1.19^{+0.26}_{-0.37}$	$36.53^{+0.31}_{-0.22}$	$-1.15^{+0.27}_{-0.25}$
100 $\mu\text{m}$	$1.07^{+0.26}_{-0.36}$	$36.70^{+0.31}_{-0.22}$	$-1.28^{+0.20}_{-0.19}$
160 $\mu\text{m}$	$1.28^{+0.21}_{-0.28}$	$36.42^{+0.25}_{-0.19}$	$-1.02^{+0.23}_{-0.22}$


Schechter (1976) showed that galaxy numbers as a function of luminosity can be fitted using three parameters;  $\phi^*$ , the number of sources;  $L^*$ , the characteristic luminosity at which a rapid change in the slope of the function is seen and  $\alpha$ , a dimensionless parameter which gives the slope of the function at luminosities less than  $L^*$ . This function is shown in the equation below

$$\phi(\log L)d \log L = \phi^* \ln 10 \left( \frac{L}{L^*} \right)^{\alpha+1} \exp \left( \frac{-L}{L^*} \right) d \log L. \quad (2)$$

We use a maximum likelihood method (Marshall et al. 1983) to fit Schechter parameters to the data at each wavelength independently, present the results in Table 1 and show these fits in Figs 4 and 5. A Schechter fit was chosen as the extra free parameter in the fits used by Saunders et al. (1990) and Soifer et al. (1987) results in the function being unable to converge on a singular set of values, and produces non-physical results for our data.

These data were compared to field galaxy luminosity functions from the literature. In the case of 70 and 160  $\mu\text{m}$ , there are parametrized luminosity function fits available from the *Spitzer* Wide-area Infrared Extragalactic survey (Patel et al. 2013) with an additional data set at 60  $\mu\text{m}$  from *IRAS* (Saunders et al. 1990).

The 100  $\mu\text{m}$  data were compared to the 90  $\mu\text{m}$  binned data presented by Serjeant et al. (2004) and Sedgwick et al. (2011). Due to the luminosity range covered by these data, and the consequent restriction and rebinning of our own data to match it, the number of Coma members available for the comparison is reduced to a level at which a  $\chi^2$  test loses any statistical power. Therefore, we fitted a Saunders et al. (1990) function to the data presented in Serjeant et al. (2004) using  $\chi^2$  minimization, excluding their two lowest bins because they show an unreasonably steep drop in space density that is inconsistent with the local 100  $\mu\text{m}$  luminosity function of Rowan-Robinson, Helou & Walker (1987), even accounting for the quoted



**Figure 4.** A comparison of the Coma and field 70  $\mu\text{m}$  luminosity function. The Coma data are presented as green circles. The green line shows the Schechter function for the Coma data at 70  $\mu\text{m}$ , using the parameters presented in Table 1. The blue dashed line shows the luminosity function derived from field galaxies at 60  $\mu\text{m}$  as taken from Saunders et al. (1990) and converted to 70  $\mu\text{m}$  assuming an M82-like SED. The red dotted line shows the luminosity function as presented by Patel et al. (2013), who use the functional form of Saunders et al. (1990).

uncertainties. We then compared our data to this parametrized function in the same manner as we have done at the other wavelengths. We choose the Serjeant et al. (2004) data because the data in Sedgwick et al. (2011) do not cover a sufficiently large luminosity range to properly constrain the function we fit.


The Coma data and the various parametrized field luminosity functions are plotted for comparison in Figs 4 and 5, with the field galaxy fits having been scaled vertically to fit our data. We run Kolmogorov–Smirnov tests to compare the unbinned luminosity distributions of our cluster members with those predicted by the functions fitted to the field luminosity functions after accounting for completeness. The numbers presented in Table 2 show the probabilities that our sample is drawn from the same parent distribution as the field luminosity functions. The probabilities for the field comparisons were calculated assuming an effective number of objects using the equation shown below, where  $N_{\text{cluster}}$  is the number of galaxies in the Coma Cluster and  $N_{\text{field}}$  is the number of galaxies used to construct the field luminosity function,

$$N_{\text{eff}} = \frac{N_{\text{cluster}} N_{\text{field}}}{(N_{\text{cluster}} + N_{\text{field}})}. \quad (3)$$

We find a high probability that Coma is drawn from the same parent distribution as the field luminosity functions. This similarity between the field and Coma luminosity functions is surprising. One would expect the numerous stripping processes as expected from both theoretical and observational studies, would have a large effect in the dense environment of the cluster. A recent study of the Fornax Cluster by Fuller et al. (2014) finds similar agreement with the field using the PACS 100 and 160  $\mu\text{m}$  bands, as well as the SPIRE 250, 350 and 500  $\mu\text{m}$  bands. Further analysis will be required in order to provide an explanation for this result. In a future paper, we will investigate the masses and morphological types of individual galaxies to look at any underlying reasons for this similarity.

## 5.2 Reassessment of Virgo data

Given the surprising consistency between Coma and the field that suggests the cluster environment has little effect on the far-infrared



**Figure 5.** A comparison of the Coma, Virgo and field 100 and 160  $\mu\text{m}$  luminosity functions. The Coma data are presented as green circles, and the HeViCS data are presented as black crosses. The green line shows the Schechter function fit for the Coma data, using the parameters presented in Table 1. The black dotted line shows our derived Schechter function fit for the Virgo data of Auld et al. (2013). The blue dashed line shows the comparison field luminosity function; Serjeant et al. (2004) for the 100  $\mu\text{m}$  plot (converted from 90  $\mu\text{m}$  assuming an M82-like SED), and Patel et al. (2013) for the 160  $\mu\text{m}$  plot.

**Table 2.** The percentage results of Kolmogorov–Smirnov tests to compare the luminosity distribution of the luminosity functions of Coma, Virgo and the field. The Saunders et al. (1990) 60  $\mu\text{m}$  luminosity function has been converted to 70  $\mu\text{m}$ , and the Serjeant et al. (2004) 90  $\mu\text{m}$  luminosity function to 100  $\mu\text{m}$  assuming the SED of M82 (Silva et al. 1998).

Wavelength	Field LF	Coma/Field	Coma/Virgo	Virgo/Field
70 $\mu\text{m}$	Patel	73		
70 $\mu\text{m}$	Saunders	78		
100 $\mu\text{m}$	Serjeant	86	94	99
160 $\mu\text{m}$	Patel	62	9	0.6

luminosity function, we reanalyse the data from the Virgo cluster as a means of comparison. Davies et al. (2010) showed the luminosity functions that had been derived from the HeViCS data. They showed that at all observed wavelengths, there was evidence of a turnover at lower luminosities. Further evidence of this was seen with the HeViCS catalogue of bright galaxies (Davies et al. 2012). Auld et al. (2013) presented far-infrared fluxes of optically selected Virgo cluster galaxies at 100 and 160  $\mu\text{m}$ . We compare these data to

the equivalent data in the present study in order to ascertain if Virgo still shows a turnover and for comparison with both our Coma Cluster luminosity function and the relevant field galaxy functions. We use the same Kolmogorov–Smirnov test method to complete these comparisons. These two wavelength bands were used as they are the only wavelengths common to both studies. We follow the assumptions made in Auld et al. (2013) that all Virgo Cluster galaxies lie at one of the three distances, 17, 23 or 32 Mpc. For the purposes of this study, we use those Virgo sources which lie at 17 and 23 Mpc and have a measured flux greater than our noise limit. This sample corresponds to that used by the latest HeViCS analysis (Davies et al. 2014). Analysis of the data shows that the inclusion of those sources at 32 Mpc, and the assumption that all sources lie at 17 Mpc, have little effect on the results.

The results of these tests are included in Table 2. The HeViCS data are plotted alongside the present study in Fig. 5, along with a Schechter function fit derived over the range of data included within our comparison. For both Virgo and Coma, we find a clear lack of a turnover at the faint end, in disagreement with the findings of Davies et al. (2010). We find good agreement between the Coma and Virgo Clusters, as well as between the Virgo Cluster and the field, at 100  $\mu\text{m}$ . At 160  $\mu\text{m}$ , however, we find no agreement between Virgo and either the Coma Cluster or the field, which still agree with each other reasonably well. We can provide no explanation as to why there would be agreement for one wavelength data set, but not for the other. However, we note that there is a steeper faint-end slope present in the Virgo 160  $\mu\text{m}$  data, as can be seen clearly in Fig. 5. This disagreement is especially confusing when one considers that the Virgo Cluster is less virialized than the Coma Cluster, and thus might be expected to have properties intermediate between those of the field and Coma.

Urban et al. (2011) showed that the Virgo cluster has a  $r_{200}$  radius value of 1.08 Mpc or  $3:9$  at their adopted distance of 16.1 Mpc, corresponding to a value of 1.16 Mpc at our defined distance of 17.0 Mpc. Arnaud, Pointecouteau & Pratt (2005) showed that the  $r_{200}$  radius value can be adopted as the virial radius for Virgo, even though the cluster is not virialized. The HeViCS observations therefore cover an area out to over twice the virial radius. In comparison, the virial radius of the Coma Cluster is 2.9 Mpc or  $1:7$  (Łokas & Mamon 2003), with our observations covering an area out to approximately this distance. The Virgo observations therefore cover a greater area outside of the core environment relative to the Coma observations. Future work will take this factor into account by looking at subsets of the Virgo observations and comparing similar environments of the clusters.

## 6 SUMMARY

We have presented the deepest far-infrared observations to date of the Coma Cluster of galaxies. The observations are used to produce far-infrared number counts within this area, and in combination with an optical redshift catalogue are used to derive far-infrared luminosity functions at the three *Herschel*/PACS wavelengths (70, 100 and 160  $\mu\text{m}$ ) for the Coma Cluster. These functions are compared with both the field galaxy luminosity function and similar results from the HeViCS survey of the Virgo Cluster. We see that our Coma Cluster luminosity functions have shapes surprisingly consistent with those of the field galaxies at the three far-infrared wavelengths we probe. The Coma Cluster luminosity function also matches well with the newly derived luminosity function for the Virgo Cluster at 100  $\mu\text{m}$ , which no longer shows evidence of a faint-end turnover. The same consistency is not seen with the Virgo Cluster at 160  $\mu\text{m}$ ,

but this cannot be easily explained. We note that it is not due to our sample consisting only of the most luminous galaxies, which would be unaffected by stripping, as Fig. 2 shows that our sources lie around the value of  $L^*$  for the  $r$  band. Future work will look in detail at the dust properties and stellar masses of the detected Coma Cluster members to try to identify a mechanism or process that could result in such similarity between the cluster and field environments.

## ACKNOWLEDGEMENTS

SH acknowledges an STFC studentship. We acknowledge Mustapha Mouhcine for his help in the initial preparation of this study, Matthew W. L. Smith for providing the H-ATLAS data for the source flux comparison, and the anonymous referee for some of his helpful comments.

*Herschel* is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA. PACS has been developed by a consortium of institutes led by MPE (Germany) and including UVIE (Austria); KU Leuven, CSL, IMEC (Belgium); CEA, LAM (France); MPIA (Germany); INAF-IFSI/OAA/OAP/OAT, LENS, SISSA (Italy); IAC (Spain). This development has been supported by the funding agencies BMVIT (Austria), ESA-PRODEX (Belgium), CEA/CNES (France), DLR (Germany), ASI/INAF (Italy) and CICYT/MCYT (Spain). HIPE is a joint development by the *Herschel* Science Ground Segment Consortium, consisting of ESA, the NASA *Herschel* Science Center and the HIFI, PACS and SPIRE consortia.

This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the NASA.

## REFERENCES

- Arnaud M., Pointecouteau E., Pratt G. W., 2005, *A&A*, 441, 893  
Auld R. et al., 2013, *MNRAS*, 428, 1880  
Bai L., Rieke G. H., Rieke M. J., Hinz J. L., Kelly D. M., Blaylock M., 2006, *ApJ*, 639, 827  
Bai L., Rieke G. H., Rieke M. J., Christlein D., Zabludoff A. I., 2009, *ApJ*, 693, 1840  
Baldry I. K., Balogh M. L., Bower R. G., Glazebrook K., Nichol R. C., Bamford S. P., Budavari T., 2006, *MNRAS*, 373, 469  
Ball N. M., Loveday J., Brunner R. J., Baldry I. K., Brinkmann J., 2006, *MNRAS*, 373, 845  
Berta S. et al., 2010, *A&A*, 518, L30  
Bertin E., Arnouts S., 1996, *A&AS*, 117, 393  
Binggeli B., Sandage A., Tammann G. A., 1985, *AJ*, 90, 1681  
Binggeli B., Popescu C. C., Tammann G. A., 1993, *A&AS*, 98, 275  
Boselli A., Gavazzi G., 2006, *PASP*, 118, 517  
Boselli A., Gavazzi G., Lequeux J., Buat V., Casoli F., Dickey J., Donas J., 1995, *A&A*, 300, L13  
Bravo-Alfaro H., Cayatte V., van Gorkom J. H., Balkowski C., 2000, *AJ*, 119, 580  
Bravo-Alfaro H., Cayatte V., van Gorkom J. H., Balkowski C., 2001, *A&A*, 379, 347  
Briel U. G. et al., 2001, *A&A*, 365, L60  
Buat V., Xu C., 1996, *A&A*, 306, 61  
Byrd G., Valtonen M., 1990, *ApJ*, 350, 89  
Caldwell N., Rose J. A., Sharples R. M., Ellis R. S., Bower R. G., 1993, *AJ*, 106, 473  
Carter D. et al., 2008, *ApJS*, 176, 424  
Colless M., Dunn A. M., 1996, *ApJ*, 458, 435  
Davies J. I. et al., 2010, *A&A*, 518, L48  
Davies J. I. et al., 2012, *MNRAS*, 419, 3505  
Davies J. I. et al., 2014, *MNRAS*, 438, 1922  
Dressler A., 1980, *ApJ*, 236, 351  
Dressler A. et al., 1997, *ApJ*, 490, 577  
Dunne L., Eales S., Edmunds M., Ivison R., Alexander P., Clements D. L., 2000, *MNRAS*, 315, 115  
Eales S. et al., 2010, *PASP*, 122, 499  
Fabricant D. et al., 2005, *PASP*, 117, 1411  
Fuller C. et al., 2014, *MNRAS*, 440, 1571  
Gavazzi G., Boselli A., Kennicutt R., 1991, *AJ*, 101, 1207  
Gavazzi G., Catinella B., Carrasco L., Boselli A., Contursi A., 1998, *AJ*, 115, 1745  
Gavazzi G., Boselli A., Scodreggio M., Pierini D., Belsole E., 1999, *MNRAS*, 304, 595  
Gavazzi G., Boselli A., Donati A., Franzetti P., Scodreggio M., 2003, *A&A*, 400, 451  
Gavazzi G., Franzetti P., Boselli A., 2014, preprint ([arXiv:1401.8123](https://arxiv.org/abs/1401.8123))  
Gehrels N., 1986, *ApJ*, 303, 336  
Griffin M. J. et al., 2010, *A&A*, 518, L3  
Gunn J. E., Gott J. R., III, 1972, *ApJ*, 176, 1  
Ibar E. et al., 2010, *MNRAS*, 409, 38  
Joseph R. D., Wright G. S., 1985, *MNRAS*, 214, 87  
Keel W. C., Kennicutt R. C., Jr, Hummel E., van der Hulst J. M., 1985, *AJ*, 90, 708  
Kennicutt R. C., Jr, 1998, *ApJ*, 498, 541  
Kennicutt R. C., Evans N. J., 2012, *A&A*, 50, 531  
Kennicutt R. C., Jr, Bothun G. D., Schommer R. A., 1984, *AJ*, 89, 1279  
Kennicutt R. C., Jr, Roettiger K. A., Keel W. C., van der Hulst J. M., Hummel E., 1987, *AJ*, 93, 1011  
Larson R. B., Tinsley B. M., Caldwell C. N., 1980, *ApJ*, 237, 692  
Liu M. C., Graham J. R., 2001, *ApJ*, 557, L31  
Łokas E. L., Mamon G. A., 2003, *MNRAS*, 343, 401  
Lonsdale Persson C. J., Helou G., 1987, *ApJ*, 314, 513  
Lutz D. et al., 2011, *A&A*, 532, A90  
Magnelli B. et al., 2013, *A&A*, 553, A132  
Marshall H. L., Tananbaum H., Avni Y., Zamorani G., 1983, *ApJ*, 269, 35  
Merritt D., 1984, *ApJ*, 276, 26  
Miller R. H., 1986, *A&A*, 167, 41  
Moore B., Lake G., Katz N., 1998, *ApJ*, 495, 139  
Moss C., Whittle M., 2005, *MNRAS*, 357, 1337  
Neumann D. M. et al., 2001, *A&A*, 365, L74  
Ott S., 2010, in Mizumoto Y., Morita K.-I., Ohishi M., eds, *ASP Conf. Ser. Vol. 434, Astronomical Data Analysis Software and Systems XIX*. Astron. Soc. Pac., San Francisco, p. 139  
Patel H., Clements D. L., Vaccari M., Mortlock D. J., Rowan-Robinson M., Pérez-Fournon I., Afonso-Luis A., 2013, *MNRAS*, 428, 291  
Pilbratt G. L. et al., 2010, *A&A*, 518, L1  
Poglitich A. et al., 2010, *A&A*, 518, L2  
Rigby E. E. et al., 2011, *MNRAS*, 415, 2336  
Rowan-Robinson M., Helou G., Walker D., 1987, *MNRAS*, 227, 589  
Saunders W., Rowan-Robinson M., Lawrence A., Efstathiou G., Kaiser N., Ellis R. S., Frenk C. S., 1990, *MNRAS*, 242, 318  
Schechter P., 1976, *ApJ*, 203, 297  
Sedgwick C. et al., 2011, *MNRAS*, 416, 1862  
Serjeant S. et al., 2004, *MNRAS*, 355, 813  
Silva L., Granato G. L., Bressan A., Danese L., 1998, *ApJ*, 509, 103  
Smith R. J., Lucey J. R., Hudson M. J., Allanson S. P., Bridges T. J., Hornschemeier A. E., Marzke R. O., Miller N. A., 2009, *MNRAS*, 392, 1265  
Smith R. J., Lucey J. R., Price J., Hudson M. J., Phillipps S., 2012, *MNRAS*, 419, 3167



Soifer B. T., Sanders D. B., Madore B. F., Neugebauer G., Danielson G. E.,  
Elias J. H., Lonsdale C. J., Rice W. L., 1987, *ApJ*, 320, 238  
Struble M. F., Rood H. J., 1999, *ApJS*, 125, 35  
Urban O., Werner N., Simionescu A., Allen S. W., Böhringer H., 2011,  
*MNRAS*, 414, 2101  
Walterbos R. A. M., Greenawalt B., 1996, *ApJ*, 460, 696

Wang G., Leggett S. K., Clowes R. G., MacGillivray H. T., Savage A., 1991,  
*MNRAS*, 248, 112  
Whitmore B. C., Gilmore D. M., Jones C., 1993, *ApJ*, 407, 489

This paper has been typeset from a  $\text{\TeX/L\AA\TeX}$  file prepared by the author.