

Durham Research Online

Deposited in DRO:

21 January 2015

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Candy, I. and McClymont, E.L. (2013) 'Interglacial intensity in the North Atlantic over the last 800,000 years : investigating the complexity of the mid-Brunhes Event (MBE).', *Journal of quaternary science.*, 28 (4). pp. 343-348.

Further information on publisher's website:

<http://dx.doi.org/10.1002/jqs.2632>

Publisher's copyright statement:

This is the accepted version of the following article: Candy, I. and McClymont, E.L. (2013), Interglacial intensity in the North Atlantic over the last 800000 years: investigating the complexity of the mid-Brunhes Event. *Journal of Quaternary Science*, 28 (4): 343-348, which has been published in final form at <http://dx.doi.org/10.1002/jqs.2632>. This article may be used for non-commercial purposes in accordance With Wiley Terms and Conditions for self-archiving.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

1 **Interglacial intensity in the North Atlantic over the last 800,000 years: investigating the**
2 **complexity of the mid-Brunhes Event (MBE).**

3 *Ian Candy¹, Erin L. McClymont²*

4 *¹Department of Geography, Royal Holloway, University of London, Egham, Surrey, TW20*
5 *0EX, UK. Tel: 01784 443649, email: ian.candy@rhul.ac.uk*

6 *²Department of Geography, Durham University, Science Laboratories, South Road, Durham,*
7 *DH1 3LE, UK.*

8
9 **Abstract**

10 **The mid-Brunhes Event (MBE) represents a step-like shift in the intensity of**
11 **interglacial warmth that occurs between MIS 13 and MIS 11, with interglacials MIS 19-**
12 **13 being significantly cooler than interglacials MIS 11-1. A transect of palaeoclimatic**
13 **records in the North Atlantic from 40°N through to the Nordic Seas indicates that there**
14 **are strong differences in the expression of the MBE at different latitudes in this region.**
15 **Between 40 and 56°N sea surface temperature and air temperature records suggest that**
16 **all interglacials of the past 800,000 years were characterised by similar levels of**
17 **warmth, therefore, there is no evidence for a MBE in these latitudes of the North**
18 **Atlantic. North of 56°N there is increasing evidence for cooler climates during**
19 **interglacials MIS 19-13 relative to MIS 11-1. This review suggests that the North**
20 **Atlantic was anomalous in comparison to the other records of interglacial climate**
21 **diversity which suggest that the MBE was a global event. Furthermore, the strong**
22 **spatial difference in temperature conditions during interglacials MIS 19-13 in the North**
23 **Atlantic means that this region would have been characterised by strong temperature**
24 **gradients during interglacial episodes of this time interval.**

25 **Introduction**

26 The mid-Brunhes Event (MBE) is the most pronounced climatic shift of the past 800,000
27 years (Jansen et al., 1986; Candy et al., 2010). It represents a step-like change in the intensity
28 of interglacial warmth that occurs between MIS13-11 (EPICA, 2004). This shift is important
29 to our understanding of global warming in the geological past as it shows that interglacials of
30 very different thermal regimes can be generated by similar patterns of insolation. Lang and
31 Wolff (2010) have argued that this is a global event, whereas Meckler et al. (2012) have
32 argued that it is not expressed in low-latitude regions. Climate modelling by Yin and Berger
33 (2012) suggest that the MBE is most strongly expressed in the high-latitudes, is absent in the
34 low-latitudes and has variable expression in the mid-latitudes. Validating these different ideas
35 is problematic because of the paucity of climate records that have a high-enough resolution
36 and long enough duration to investigate interglacial climate diversity over the past 800,000
37 years (Lang and Wolff, 2010). It is only in the north Atlantic that a sufficient density of
38 appropriate records exists to investigate how long-term patterns of interglacial warmth vary
39 between different latitudes (e.g. McManus et al., 1999; Lawrence et al., 2009). This study
40 investigates the expression of the MBE in the North Atlantic along a transect of sites from
41 40°N-62°N.

42

43 **Methodology**

44 Lang and Wolff's (2010) review of interglacial climates of the past 800,000 years concluded
45 that "*strong interglacials are confined to the last 450 ka, and that this is a globally robust*
46 *pattern*". This is the clearest definition of the MBE yet published. As such the
47 intensity/strength of the MBE can be calculated, for any given palaeoclimatic record, by

48 subtracting the mean, maximum interglacial intensity of pre-MBE interglacials from the
49 mean, maximum interglacial intensity of post-MBE interglacials:

$$50 \quad \text{MBE intensity} = \text{MTMAX}_{\text{postMBE}} - \text{MTMAX}_{\text{preMBE}}$$

51 Where:

$$52 \quad \text{MTMAX}_{\text{postMBE}} = (\text{MIS1}_{\text{TMAX}} + \text{MIS5e}_{\text{TMAX}} + \text{MIS7}_{\text{TMAX}} + \text{MIS9}_{\text{TMAX}} + \text{MIS11}_{\text{TMAX}})/5$$

$$53 \quad \text{MTMAX}_{\text{preMBE}} = (\text{MIS13}_{\text{TMAX}} + \text{MIS15}_{\text{TMAX}} + \text{MIS17}_{\text{TMAX}} + \text{MIS19}_{\text{TMAX}})/4$$

54 $\text{MIS}_{n\text{TMAX}}$ = the maximum temperature value recorded in the time interval of MIS_n .

55 In any record where the MBE is present the MBE intensity will always be >0 . In the EPICA
56 deuterium based air temperature anomaly record and the ODP 1090 sea surface temperature
57 record from the southern ocean (Figure 1 and Table 1), MBE intensity = 4.47 and 3.16
58 respectively. This means that, on average, pre-MBE interglacials in Antarctica were 4.47°C
59 cooler than post-MBE interglacials, whilst in the ODP 1090 record of the southern ocean pre-
60 MBE interglacials were, on average, 3.16°C cooler than post-MBE interglacials. This
61 approach does not consider interglacial duration or the relative timing of the interglacial
62 within a warm stage, it simply identifies the maximum temperature value within the
63 chronological boundaries of each warm stage and uses these values in the calculation. This is
64 considered the most appropriate approach as the MBE is primarily defined by shifts in
65 maximum interglacial intensity not by duration of relative warmth.

66

67 In the North Atlantic there are a number of temperature records between 40°N and 57°N
68 (Figure 2 and 3). The majority of these are SST records, generated from a range of proxy
69 data; foraminifera assemblage transfer functions (DSDP 607 (Ruddiman et al., 1989),
70 M23414 (Kandiano and Bauch, 2003) and ODP 552 (Ruddiman et al., 1986)), U^{K}_{37} alkenone
71 data (ODP 982 and 983 (McClymont et al., 2008; Lawrence et al., 2009)) and oxygen

72 isotopic data (ODP 980 (McManus et al., 1999)). Air temperature records are obtainable from
73 palaeoecological assemblages preserved in the British terrestrial record (Coope, 2010; Candy
74 et al., 2010; 2011). Two of these records (ODP 980 and M23414) only span the last
75 500,000yrs (MIS1-MIS13). Although this means it is difficult to reliably calculate MBE
76 intensity values they still provide important information on the presence/absence of the MBE.
77 If, in such records, the peak intensity of MIS13 is as warm or warmer than MIS 1-11 it shows
78 that “*strong interglacials*” are not “*constrained to the last 450ka*” and, therefore, implies that
79 the MBE is absent from that record.

80

81 North of 58°N, there are no continuous temperature records. However, the general
82 temperature characteristics of interglacials can be inferred from multiple sites (Figure 2 and
83 3) through a number of indicators including foraminifera assemblages, particularly the
84 abundance of *Neogloboquadrina pachyderma* (sin.) (Wright and Flower, 2002), and the
85 abundance of an alkenone associated with the presence polar water masses (%C_{37:4};
86 McClymont et al., 2008). Further north in the Norwegian/Greenland Sea the nature of the
87 marine records (heavily influenced by IRD) makes it more difficult to make clear conclusions
88 about long term interglacial diversity, although basic conclusions can be made using ice-
89 rafted detritus (IRD) concentrations, carbonate content and characteristics of biogenic
90 materials, such as percentage of sub-polar foraminifera (see Henrich and Baumann, 1994;
91 Helmke et al., 2003; Bauch, 2013).

92

93 **The MBE in the North Atlantic 40 to 56°N**

94 The key SST records of the mid-latitude North Atlantic (40°-56°N) are shown in Figure 3. In
95 all of these records there is no difference in mean interglacial temperature maxima between
96 MIS11-1 and MIS19-13 (Table 1). In most cases the difference in mean interglacial
97 temperature maxima is minimal, in all cases $<\pm 0.5^{\circ}\text{C}$ difference, and within 1 standard
98 deviation of each other (Table 1). As most of the proxy techniques that are used to
99 reconstruct temperatures have uncertainties in the order of $\pm 1.0^{\circ}\text{C}$ the differences in pre-
100 and post-MBE warmth are all within the uncertainties of the associated techniques. In the
101 SST records from the North Atlantic between 40-56°N there is, therefore, no statistical
102 difference between the magnitude of interglacial temperature maxima during MIS19-13 and
103 MIS11-1. Palaeotemperature reconstructions from the British record are based on a range of
104 proxies so a simple calculation of MBE intensity is not possible. However, even, the coolest
105 of the MIS19-13 temperature reconstructions, West Runton, indicate climates at least as
106 warm as the present day, whilst deposits at Sidestrand and Pakefield indicate climates as
107 warm as the Eemian, the warmest interglacial of the past 500,000 years (Candy et al., 2010).

108

109 **The MBE in the North Atlantic $>56^{\circ}\text{N}$**

110 The key SST record in this region is ODP 982 where a U_{37}^K alkenone-based temperature
111 record has been constructed for the past 4 million years (Lawrence et al., 2009). In ODP 982
112 the interglacials MIS19-13 are clearly cooler than MIS11-1 (Table 1). No other quantified
113 temperature records that continuously span the past 800,000 years exist for the North Atlantic
114 in latitudes higher than 56°N . Although an alkenone SST record does exist for ODP 983
115 (60°N ; McClymont et al., 2008) the focus of the study was the Mid-Pleistocene Transition
116 and, consequently, the SST record finishes at 500,000 yrs. Wright and Flower (2002) have
117 suggested that interglacials MIS 19 to 13, in ODP 984 (61°N) are routinely cooler than

118 interglacials of the past 500,000 years. This is based on percentage concentrations of *N.*
119 *pachyderma* (s). In records of MIS 5 from the nearby core EW9302-8JPC *N. pachyderma* (s)
120 concentrations drop to 0% (Oppo et al., 2001), whilst during MIS 19-13 they are routinely
121 between 25-50%. Higher concentrations of *N. pachyderma* (s) during MIS 19-13 thus suggest
122 that these interglacials were all significantly cooler than those of the late Pleistocene. Wright
123 and Flower (2002) proposed that during interglacials MIS 19 and 17 the polar front lay to the
124 southeast of ODP 984 but to the northwest of ODP 980. This is supported by the %C_{37:4}
125 alkenone concentration record at ODP 983 which indicate cold Arctic waters extending into
126 the North Atlantic during both MIS 19 and 17 (McClymont et al., 2008).

127

128 **The MBE in the Nordic Seas**

129 The pattern of interglacial warmth in the Nordic Seas is more difficult to characterise. A
130 number of authors have argued that a major shift to warmer interglacials occurs from MIS15
131 onwards (Henrich and Baumann, 1994). This suggestion is based on percentage carbonate
132 content, however, this parameter is not simply a proxy for temperature as high concentrations
133 of *N. pachyderma* (s) can generate high percentage carbonate values even during relatively
134 cool episodes. There is, however, a general suggestion from across the Norwegian and
135 Greenland seas that there is a cooler “aspect” to interglacials MIS19-13 (Jansen et al., 1988;
136 Fronval and Jansen, 1994; Henrich and Baumann, 1994; Helmke et al., 2003). In MD992277,
137 for example, MIS19, 17 and 13 appear to be characterised by relatively subdued warmth
138 (Helmke et al., 2003). Although MIS15 appears to be significantly warmer than the other pre-
139 MBE interglacials it is not comparable in magnitude to MIS11, for example, and also appears
140 to be of relatively short duration; ca 5,000 years relative to the 10,000 years of MIS11
141 (Helmke et al., 2003). This pattern of cool interglacials before MIS11 is supported by other

142 records from this region (Jansen et al., 1988; Fronval and Jansen, 1994; Henrich and
143 Baumann, 1994).

144

145 **Discussion**

146 There is no evidence for a mid-Brunhes Event in the mid-latitude North Atlantic, south of
147 56°N (Figure 2). Although these palaeotemperature records have been generated using a
148 range of techniques, the validity of the reconstructions is strengthened by that fact that sites
149 on the same approximate line of latitude produce a consistent picture. For example, in ODP
150 552, 980, M23414 and the British terrestrial sequence MIS13 is at least as warm as the
151 Holocene (Ruddiman et al., 1989; McManus et al., 1999; Flower and Wright, 2002; Candy et
152 al., 2010). North of 56°N there is evidence for cooler temperatures during MIS19-13 (e.g.
153 Lawrence et al., 2009). Although difficult to quantify this pattern is considered robust
154 because it can be seen in alkenone, faunal and lithological proxies. This pattern appears to be
155 true of the Nordic Seas although the nature of the proxy records makes it difficult to be
156 precise about the nature of the thermal regime of interglacials MIS19-13 other than to say
157 that they are ‘cool’ (Helmke et al., 2003). The mid-latitude North Atlantic is, therefore,
158 anomalous in the light of most reviews of interglacial diversity in that “strong interglacials”
159 (cf Lang and Wolff, 2010) occur prior to 450ka ago. Evidence for the MBE in the higher
160 latitudes of this region is, however, consistent with the work of Yin and Berger (2012).

161

162 In the North Atlantic interglacials, MIS19-13 appear to be characterised by a southward
163 expansion of cold polar-water into the North Atlantic relative to MIS11-1 (Wright and
164 Flower, 2002; McClymont et al., 2008). Wright and Flower (2002) have postulated that

165 during some interglacials of the early Middle Pleistocene the Polar Front was situated south
166 of 61°N but north of 56°N. Here, we show that evidence from a range of sites supports this
167 interpretation, but also that the expression of the MBE in the north Atlantic region is a
168 function of the shift, equatorwards, of the polar front. In this respect the interglacial history of
169 the North Atlantic is consistent with that of the Southern Ocean, where cooler interglacial
170 temperatures during MIS19-13 are suggested to correspond with a more northerly position of
171 the polar front relative to its position during MIS11-1 (Kujipers, 1989; Martinez-Garcia et al.,
172 2009).

173

174 The southward expansion of polar waters during MIS19-13 means that the climatic setting
175 that existed in the North Atlantic during these intervals would have been very different to that
176 which occurred during MIS11-1. In particular the climate of the North Atlantic during
177 interglacials MIS19-13 would have been characterised by stronger latitudinal temperature
178 gradients. If, during MIS19-13, sea surface and air temperatures in the region 40-56°N are as
179 warm as MIS11-1 but the regions 57°N and higher are significantly cooler then much steeper
180 temperature gradients than are experienced at the present day would have existed across the
181 North Atlantic (Figure 4).

182

183 The MBE is routinely defined in terms of long-term shifts in the magnitude of interglacial
184 warmth and in this respect there is no evidence for an MBE in, for example, the British Isles.
185 However, the southward intrusion of arctic waters during MIS19-13 may have had impacts
186 on atmospheric circulation, seasonality (of both temperature and rainfall) and annual
187 precipitation. Consequently, although there is no evidence for the MBE in the North Atlantic
188 (40-56°N), this does mean that the climates of MIS19-13 were directly analogous to those of

189 MIS11-5. The climate of MIS19-13 in the North Atlantic region may have been significantly
190 different because of this unusual synoptic setting. As MIS19-13 record the first arrival of
191 early humans into northern Europe and persistently high levels, for warm isotopic stages, of
192 northern hemisphere ice volume the unusual synoptic settings of these interglacial has
193 implications, not just for the diversity of interglacial climates, but also for understanding the
194 context of both early human evolution and long-term ice-sheet dynamics.

195

196 **Captions**

197 *Table 1* – Quantification of the strength of the MBE, see text for methodology, as represented
198 in multiple long temperature time series (s = summer SST, w = winter SST). Temperatures
199 in these records are generated by a number of techniques making the direct comparison of the
200 absolute temperature of an individual interglacial between different records problematic.
201 However, the construction of a temperature difference effectively normalises the record
202 allowing direct comparison of the strength of the MBE to be calculated. EPICA Dome C
203 (EPICA, 2004; Jouzel et al., 2007) is based on comparing deuterium based temperature
204 anomalies. ODP 1090 is in the Southern Ocean and is based on U_{37}^K quantification of sea
205 surface temperatures (Bequey and Gersonde, 2002; Martinez-Garcia et al., 2009) . Sea
206 surface temperatures in DSDP 607 and ODP 552 are based on foraminifera assemblages
207 (Ruddiman et al., 1986;1989). The ODP 980 record only records the past 500,000 years and,
208 therefore, contains a detailed record of MIS 13 but none of the other early Middle Pleistocene
209 interglacials. The temperature difference calculated for this core is, therefore, not comparable
210 with the others, however, it does suggest contain a peak in MIS 13 that is comparable to those
211 of MIS 11 to 1. ODP 980 temperatures are based oxygen isotopic differences between
212 planktonic and benthic foraminifera (McManus et al., 1999). A similar situation is found in

213 MD23414 (53°N, 20°W) where a SST record of the past 500,000 years records MIS 13 SST
214 as warm as MIS 1, 7 and 9 and *N. Pachyderma* (s) concentrations of 0 (Kandiano and Bauch,
215 2003). Wright and Flower (2002) have calculated SST values for the early Middle
216 Pleistocene section of ODP 980 using foraminifera-based transfer functions. Although the
217 calculated temperatures cannot be directly compared with the SST estimates of McManus et
218 al. (1999) they do indicate that MIS 19 to 13 in ODP 980 were at least as warm as modern
219 day and MIS 5e values. The temperature record of ODP 982 is generated from U_{37}^K prime
220 (Lawrence et al., 2009). Although the resolution of the ODP 982 is greatly reduced during
221 MIS19 to13 the reduction in resolution is progressive and, if the lower temperatures were due
222 to the main interglacial peaks being missed, it would be anticipated that interglacial peaks
223 would also gradually decrease. This is not the case and there is a clear step change in
224 interglacial intensity between MIS 13 and 11. Furthermore the pattern seen in ODP 982 is
225 supported by other records in the region (e.g. ODP 984). ODP 984 records % *N. Pachyderma*
226 (s) with higher percentages indicating colder waters (Wright and Flower, 2002). In this region
227 MIS 5 is characterised by *N. Pachyderma* (s) of 0% (Oppo et al., 2001), whereas ODP 984
228 records values between 70 and 30% during MIS 19 to 13, indicating much cooler conditions.
229 Standard deviations for maximum interglacial temperatures of MIS 19 to 13 and 11 to 1 are
230 shown. Significantly the mean interglacial temperature maxima of MIS 19 to 13 and MIS 11
231 to 1 do not overlap even when standard deviations are included for EPICA Dome C, ODP
232 1090 and ODP 982. Mean interglacial temperature maxima of MIS 19 to 13 and MIS 11 to 1
233 do overlap at DSDFP 607, ODP 980 and ODP 552. Statistically, therefore, there is no
234 significant difference between interglacial intensity across the MBE for those North Atlantic
235 sites between 41 and 56°N but a significant difference in interglacial intensity in EPICA
236 Dome C, ODP 1090 and ODP 982. Due to the limited number of interglacials in the pre- and
237 post-MBE intervals no more detailed statistical analysis was possible.

238

239 *Figure 1* – Examples of long palaeoclimate records that show a clear expression of the MBE.

240 1a is the EPICA Dome C deuterium record and shows air temperature variability in
241 Antarctica (EPICA, 2004; Jouzel et al., 2007), 1b is the SST record from ODP 1090 in the
242 southern Ocean (Bequey and Gersonde, 2002; Martinez-Garcia et al., 2009). In Antarctica
243 post-MBE interglacials are, on average, 4.47°C warmer than pre-MBE interglacials. In the
244 ODP 1090 record of the southern ocean post-MBE interglacials are, on average 3.16°C
245 warmer than pre-MBE interglacials.

246

247 *Figure 2* – The location of key palaeoclimate records that span the MBE, discussed in the
248 text, in the North Atlantic and Nordic Seas. and shows the location of the key records that are
249 discussed in the text.

250

251 *Figure 3* – Palaeoclimate records from the North Atlantic that are discussed in the text. DSDP
252 607 (Ruddiman et al., 1989; Lawrence et al., 2010), M23414 (Kandiano and Bauch, 2003),
253 ODP 980 (McManus et al., 1999), ODP 552 (Ruddiman et al., 1986), British
254 Palaeoecological Record (BPR, Candy et al., 2010), ODP 982 (Lawrence et al., 2009), ODP
255 984 (Wright and Flower, 2002).

256

257 *Figure 4* – Summary of the palaeoclimate of early Middle Pleistocene interglacials (MIS 19
258 to 13). Areas shown in red are those that contain evidence to suggest that interglacials MIS
259 19 to 13 were routinely as warm as MIS 11 to 5 (e.g. areas with no evidence for an MBE).
260 Areas shown in blue are those that contain evidence to suggest that interglacials MIS 19 to 13

261 were routinely cooler than MIS 11 to 5 (e.g. areas with a clear expression for an MBE). The
262 current position of the Polar Front is shown for reference.

263

264

265

266 **References**

267 Bauch, H. A., 2013. Interglacial climates and the Atlantic meridional overturning circulation:
268 is there an Arctic controversy? *Quaternary Science Reviews*, 63, 1-22.

269 Candy, I., Rose, J., Coope, G.R., Lee, J.R., Parfitt, S.P., Preece, R.C., and Schreve, D.C.,
270 2010. Pronounced climate warming during early Middle Pleistocene interglacials:
271 investigating the mid-Brunhes event in the British terrestrial sequence. *Earth Science*
272 *Reviews*, 103, 183-196.

273 Candy, I., Silva, B. and Lee, J.R., 2011. Climates of the early Middle Pleistocene in Britain:
274 Environments of the earliest humans in northern Europe. In Ashton, N., Lewis, S.G. and
275 Stringer, C. (eds). *The Ancient Human Occupation of Britain Project. Developments in*
276 *Quaternary Science*, Elsevier, 11-21.

277 Coope, G.R., 2010. Coleopteran faunas as indicators of interglacial climates in central and
278 southern England. *Quaternary Science Reviews*, 29, 1507-1514.

279 EPICA community, 2004. Eight glacial cycles from an Antarctic ice core. *Nature*, 429, 623-
280 628.

281 Fronval, T., and Jansen, E., 1996. Late Neogene paleoclimates and paleoceanography in the
282 Iceland-Horwegian sea: evidence from the Iceland and Voring Plateaus. In: Thiede, J.,
283 Myhre, A.M., Firth, J.V., Johnson, G.L., Ruddiman, W.F. (eds), *Proceedings of the Ocean*
284 *Drilling Program, Scientific Results*, 151. College Station, TX, 455-468.

285 Helmke, J.P., Bauch, H.A., Erlenkeuser, H., 2003. Development of glacial and interglacial
286 conditions in the Nordic seas between 1.5 and 0.35 Ma. *Quaternary Science Reviews*, 22,
287 1717-1728.

288 Henrich, R., and Baumann, K.H., 1994. Evolution of the Norwegian Current and the
289 Scandinavian Ice Sheet during the past 2.6 m.y.: evidence from ODP leg 104 biogenic
290 carbonate and terrigenous records. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 108,
291 75-94.

292 Kuijpers, A., 1989. Southern Ocean circulation and global climate in the Middle Pleistocene
293 (early Brunhes). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 76, 67-83.


- 294 Jansen, J.H.F., Kuijpers, A., Troelstra, S.R., 1986. A Mid-Brunhes Climatic Event: Long-
295 Term Changes in Global Atmosphere and Ocean Circulation. *Science*, 4750, 619-622.
- 296 Jouzel, J., V. Masson-Delmotte, O. Cattani, G. Dreyfus, S. Falourd, G. Hoffmann, B.
297 Minster, J. Nouet, J. M. Barnola, J. Chappellaz, H. Fischer, J. C. Gallet, S. Johnsen, M.
298 Leuenberger, L. Loulergue, D. Luethi, H. Oerter, F. Parrenin, G. Raisbeck, D. Raynaud, A.
299 Schilt, J. Schwander, E. Selmo, R. Souchez, R. Spahni, B. Stauffer, J. P. Steffensen, B.
300 Stenni, T.F. Stocker, J.-L. Tison, M. Werner, Wolff, E.W., 2007. Orbital and millennial
301 Antarctic climate variability over the past 800,000 years. *Science*, 793-796.
- 302 Kandiano E.S., Bauch H.A., 2003. Surface ocean temperatures in the Northeast Atlantic
303 during the last 500,000 years: Evidence from foraminiferal census data. *Terra Nova*, 4, 265-
304 271.
- 305 Lang, N., and Wolff, E.W., 2010. Interglacial and glacial variability from the last 800ka in
306 marine, ice and terrestrial archives. *Climates of the Past*, 7, 361-380.
- 307 Lawrence, K.T., Herbert, T.D., Brown, C.M., Raymo, M.E., Haywood, A.M., 2009. High-
308 amplitude variations in North Atlantic sea surface temperature during the Pliocene warm
309 period. *Paleoceanography*, 24, PA2218.
- 310 Lawrence, K.T., Sosdian, S., White, H.E., Rosenthal, Y., 2010. North Atlantic climate
311 evolution through the Plio-Pleistocene climate transitions. *Earth and Planetary Letters*, 300,
312 329-342.
- 313 Martinez-Garcia. A., Rosell-Mele, A., Geibert, W., Gersonde, R., Masque, P., Gaspari, V.,
314 Barbante, C., 2009. Links between iron supply, marine productivity, sea surface temperature
315 and CO₂ over the last 1.1 Ma. *Palaeoceanography*, 24, PA1207.
- 316 McClymont, E.L., Rosell-Mele, A., Haug, G., Lloyd, J.M., 2008. Expansion of subarctic
317 water masses in the North Atlantic and Pacific oceans and implications for mid-Pleistocene
318 ice sheet growth. *Paleoceanography*, 23, PA4214.
- 319 McManus, J.F., Oppo, D.W., Keigwin, L.D., Cullen, J.L., Bond, G.C., 2002. Thermohaline
320 circulation and prolonged interglacial warmth in the North Atlantic. *Quaternary Research*, 58,
321 17-21.
- 322 Meckler, A.N., Clarkson, M.O., Cobb, K.M., Sodemann, H., Adkins, J.F., 2012. Interglacial
323 Hydroclimate in the Tropical West Pacific Through the Late Pleistocene. *Scienceexpress*
324 (<http://www.sciencemag.org/content/early/recent/3 May 2012/10.1126/science.1218340>.)
- 325 Oppo, D.W., Keigwin, L.D., McManus, J.F., Cullen, J.L., 2001. Persistent suborbital climate
326 variability in marine isotope stage 5 and Termination II. *Paleoceanography*, 16, 2802-92.
327
- 328 Ruddiman, W.F., Shackleton, N.J., McIntyre, A., 1986. North Atlantic sea-surface
329 temperatures for the last 1.1 million years. In: Summerhayes, C.P., Shackleton, N.J. (Eds.),
330 North Atlantic Paleoceanography. Geological Society of America Special Publication,
331 21,155-173.
- 332 Ruddiman, W.F., Raymo, M.E., Martinson, D.G., Clement, B.M., Backman, J., 1989.
333 Pleistocene evolution: Northern Hemisphere ice sheets and North Atlantic Ocean.
334 *Paleoceanography*, 4, 353-412.

335 Wright, A.K., Flower, B.P., 2002. Surface and deep ocean circulation in the subpolar North
336 Atlantic during the mid-Pleistocene revolution. *Paleoceanography*, 17, 1068,
337 doi:10.1029/2002PA000782.

338 Yin, Q.Z. and Berger, A., 2012. Individual contribution of insolation and CO₂ to the
339 interglacial climates of the past 800,000 years. *Climate Dynamics*, 38, 709-724.


340

341 Figure 1



342


343 Figure 2




344

345

a) North Atlantic palaeoclimate records 40 to 57°N



b) North Atlantic and Nordic Sea records >57°N



347

348

349 Figure 4




Table 1. Quantification of the strength of the MBE, see text for methodology, as represented in multiple long temperature time series (s = summer SST, w = winter SST).

Isotope stage	Temp. difference between interglacial peaks at 41°N and 57°N												
	EPICA Dome C	ODP 1090	DSDP 607	DSDP 607	DSDP 607	DSDP 607	ODP 552	M23414	M23414	ODP 982	ODP 643A	ODP 907	ODP 985
Latitude:	-75.100000	-42.913700	41.001200	41.001200	41.001200	56.042700	53.538333	53.538333	57.512667	67.715000	69.241500	66.941400	
Longitude:	123.350000	8.899700	-32.957300	-32.957300	-32.957300	-23.231300	-20.288333	-20.288333	-15.854183	1.033300	-12.698300	-6.450300	
	Deuterium SST	Alkenone SST	Foram SST (s)	Foram SST (w)	Alkenone SST	Foram SST (w)	Foram SST (s)	Foram SST (s)	Alkenone SST	CaCO ₃ (%)	CaCO ₃ (%)	CaCO ₃ (%)	
1	2.12	14.51	24.10	17.20		9.80	14.40	11.00	15.00	10.28	48.86	50.77	
5	5.46	17.14	25.10	17.40		11.00	15.70	11.50	16.20	37.93	17.59	34.82	
7	2.73	10.22	20.50	13.80		11.60	14.40	10.60	15.00	22.10	15.08	24.82	
9	3.75	14.67	23.60	16.40	17.70	9.40	14.00	10.50	15.80	15.90	15.42	17.73	1.90
11	3.15	13.93	26.60	19.50	17.70	11.10	15.40	11.40	15.60	44.34	53.53	60.03	2.10
Mean	3.44	14.09	23.98	16.86	17.70	10.58	14.78	11.00	15.52	26.11	30.10	37.63	2.18
SD	1.14	2.23	2.02	1.84	n/a	0.84	0.65	0.40	0.47	14.52	19.36	17.62	
13	-1.36	10.24	22.30	15.20	17.40	10.60	14.60	10.90	13.70	20.28	10.02	23.84	3.70
15	-0.92	12.07	22.90	16.10	18.80	10.70			14.40	20.03	25.16	36.38	4.40
17	-1.55	11.07	25.20	18.20	17.60	11.10			14.20	11.14	10.87	10.33	3.40
19	-0.54	10.35	24.00	16.80	18.10	9.90			14.10	21.02	8.63	22.71	4.00
Mean	-1.09	10.93	23.60	16.58	17.98	10.58	14.60	10.90	14.10	18.12	13.67	23.32	3.88
SD	0.39	0.73	1.11	1.10	0.54	0.43	0.00	0.00	0.25	4.67	7.72	10.64	
Temp. difference	4.53	3.16	0.38	0.28	-0.28	0.01	0.18	0.10	1.42				