

Durham Research Online

Deposited in DRO:

12 February 2015

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Cowie, D. and Braddick, O. and Atkinson, J. (2008) 'Visual control of action in step descent.', *Experimental brain research.*, 186 (2). pp. 343-348.

Further information on publisher's website:

<http://dx.doi.org/10.1007/s00221-008-1320-1>

Publisher's copyright statement:

The final publication is available at Springer via <http://dx.doi.org/10.1007/s00221-008-1320-1>.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

**Visual control of action in step descent**

Journal:	<i>Experimental Brain Research</i>
Manuscript ID:	EBR-08-0044
Manuscript Type:	Research Note
Date Submitted by the Author:	21-Jan-2008
Complete List of Authors:	Cowie, Dorothy; Institute of Neurology, University College London, Sobell Department Braddick, Oliver; University of Oxford, Experimental Psychology Atkinson, J; University College London, Psychology
Keywords:	human, visuomotor, locomotion, stair, monocular



view

VISUAL CONTROL OF ACTION IN STEP DESCENT**Dorothy Cowie, Oliver Braddick, Janette Atkinson****D Cowie**

Sobell Department, Institute of Neurology, University College London, 8-11 Queen Square, London WC1N 3BG

O Braddick

Department of Experimental Psychology, Oxford University, South Parks Road, Oxford OX1 3UD

J Atkinson

Department of Psychology, University College London, 26 Bedford Way, London WC1E 6BT

Corresponding author:

Dorothy Cowie

E-mail: d.cowie@ion.ucl.ac.uk

Tel.: 020 7837 3611 ext 4463

Fax: 020 7278 9836

Abstract

Visual guidance of forwards, sideways, and upwards stepping has been investigated, but there is little knowledge about the visuomotor processes underlying stepping down actions. In this study we investigated the visual control of a single vertical step. We measured which aspects of the stepping down movement scaled with visual information about step height, and how this visual control varied with binocular vs monocular vision. Subjects stepped down a single step of variable and unpredictable height. Several kinematic measures were extracted including a new measure, 'kneedrop'. This describes a transition in the movement of the lower leg, which occurs at a point proportional to step height. In a within-subjects design measurements were made with either full vision, monocular vision, or no vision. Subjects scaled kneedrop relative to step height with vision, but this scaling was significantly impaired in monocular and no vision conditions. The study establishes a kinematic marker of visually controlled scaling in single-step locomotion which will allow further study of the visuomotor control processes involved in stepping down.

Introduction

Everyday locomotion often involves obstacles or significant changes in the physical environment which must be visually registered and accommodated into the walking pattern. This kind of visual guidance has been examined in anterior-posterior and medio-lateral directions (Lyon & Day, 1997), as well as stepping upwards over an obstacle (Patla et al, 1991). However much less is known about the control of stepping down.

1
2
3 Steps down are encountered frequently in both manmade and natural environments and
4 are of applied importance as one of the most frequent causes of falls during walking
5 (Startzell et al., 2000). This paper examines the visually driven adjustments to
6 locomotion made in response to a single step down. We will concentrate on locomotion
7 down a regular step with distinct 'platforms' and 'risers', though we presume that similar
8 principles apply to less regular surface features that require a descending foot placement
9 in a natural environment.
10
11
12
13
14
15
16
17
18
19

20
21
22 What is the role of vision in step descent? Beyond the binary decision to attempt a step or
23 not (Warren, 1984), visual information about step height enables aspects of movement to
24 be scaled appropriately for the depth of the step. In a series of EMG studies (see Santello,
25 2005 for a review) participants stepped, or made controlled falls, down a step whose riser
26 height varied between trials. These studies show that there is a burst of calf muscle
27 activity just before landing on a step. This activity may increase joint stiffness for
28 landing; crucially, the activity occurs later for deep steps. This ability to scale the onset of
29 the EMG burst to the step height is likely to depend on visual input about the riser height.
30 Craik, Cozzens & Freedman (1982) found that pre-landing EMG activity disappeared
31 when participants were blindfolded, and was reduced when the surround of the step
32 moved down while the participant stepped. However this method is not very robust for
33 quantifying how movement depends on vision condition, since defining the burst onset is
34 difficult when it is weak.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 An alternative approach to quantifying how movement changes with riser height is to
4 measure kinematics during step descent. The biomechanics of staircase descent have been
5 in some respects well characterised (McFadyen & Winter, 1988; Protopapadaki,
6 Drechsler, Cramp, Coutts & Scott, 2007). Furthermore, Riener, Rabuffetti, & Frigo (2002)
7 showed that during stair descent maximum hip and knee flexion angles depended on stair
8 inclination. However, it was unclear whether the dependence was on tread depth or riser
9 height. Several studies have divided the step down into phases. 'Foot placement' (FP),
10 the last phase before foot contact, is associated with extension at lower limb joints, which
11 prepares the body for weight acceptance (McFadyen & Winter, 1988; Zachazewski, Riley
12 & Krebs, 1993). MacFadyen & Winter simply define FP as beginning halfway through
13 the swing phase and ending on foot contact (MacFadyen, personal communication,
14 2006). We tried to define phases more stringently and determine if the transition between
15 them depended on riser height.

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36 One potential source of visual information to step depth is binocular information. It has
37 been claimed that this provides a cue to depth in locomotor tasks. For example
38 participants increase toe clearance over an obstacle when stepping over it with monocular
39 viewing (Patla, Niechwiej, Racco & Goodale, 2002). Likewise in an obstacle avoidance
40 task (Chajka, Vecellio, Hayhoe & Gillam, 2007) monocular viewing causes participants
41 to make longer fixations on obstacles and the floor, and to increase total movement time.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

One potential source of visual information to step depth is binocular information. It has been claimed that this provides a cue to depth in locomotor tasks. For example participants increase toe clearance over an obstacle when stepping over it with monocular viewing (Patla, Niechwiej, Racco & Goodale, 2002). Likewise in an obstacle avoidance task (Chajka, Vecellio, Hayhoe & Gillam, 2007) monocular viewing causes participants to make longer fixations on obstacles and the floor, and to increase total movement time. These authors interpret their findings as showing a role of binocular vision in the guidance of locomotion. However the role of binocular information in stepping down has

not yet been studied. It is likely to be important for perceiving the step's depth, which allows appropriate movement scaling.

This study examined stepping down a single step. In many of the EMG and kinematic studies participants had prior experience of descending each step, so non-visual information about riser height was also available to them. This makes it difficult to infer whether movement scaling to riser height was really visually controlled and predictive. We carefully determined that control is visual by varying riser height between trials rather than in blocks (so vision must be used on every trial) and by measuring movement in a blindfold condition. We present a novel kinematic marker of visual control which captures how movements are planned on the basis of visual information about step height; and, by removing binocular information, we assess the potential contributions of binocular visual cues to the scaling process.

Method

Participants

Ten adults with normal or corrected normal vision took part (mean age 22.1, s.d. 3.6 years, mean height 173.5 cm, s.d. 9.8 cm, five males). All had normal stereo acuity on the TNO test (Institute for Perception TNO 1972). Eye dominance was measured by asking participants to look through a tube with one eye three times; all participants chose to look with same eye for all three trials and this was taken to be their 'dominant' eye.

Equipment

Kinematic data were recorded using a 6-camera motion tracking system (SMART, Milan) operating at 60 Hz. Cameras fixed at ceiling height surrounded a 13m³ testing area, allowing accurate 3-D reconstruction of marker positions. On each leg the participant wore a marker on the Lateral Epicondyl (LE), Lateral Malleolus (LM), Heel (H), and 5th Metatarsal Head (MH). Participants were barefoot and wore shorts to allow easy camera viewing of the kinematic markers. A simple 'step' from an 'upper platform' to a 'lower platform' was constructed. The height of its upper platform was constant for all trials and step height was varied by changing the lower platform between trials, so the step up at the start of each trial was no guide to the height to be descended.

Procedure

The task was to take a single step down from the upper platform to the lower platform. The participant took one practice step down to familiarise them with the basic task before markers were attached. Before each trial the participant waited away from the step, which they could not see. On 'no vision' trials they were then fitted with a blindfold; on monocular trials they were fitted with an eye patch. On all trials they then closed their eyes and were led to the upper platform. On vision and monocular trials they were instructed "open your eyes and step down when you're ready"; on no vision trials they were instructed "step down when you're ready". They were asked to step off the upper platform onto the lower platform as normally as possible, leading with one foot.

Step dimensions

Riser heights were scaled to leg length to allow comparison between participants of different height. Leg length was measured as the distance from ASIS to medial malleolus. The height of the upper platform was 24% leg length + 18mm. The range of riser heights (8 – 24 % leg length) was designed to be as extreme as was compatible with safety and normal stepping behaviour. For an adult of average leg length (90 cm) shallow, medium, and deep steps were ~7cm, 14cm, and 21cm.

Design

Within-subjects factors were riser-height (8, 16, 24% leg length) and vision-condition (vision (V), monocular (M), no vision (NV)). Each participant completed 3 blocks, totalling 27 trials. Trial types were randomised within a block, with each block containing all nine riser-height (3) x vision-condition (3) trial types. Half the group (3 males, 2 females) always had their dominant eye covered, half their nondominant eye (2 males, 3 females).

Data Analysis

Data were analysed using SMART software (BTS, Milan). We extracted several measures from each trial. After analysis, we averaged data from the 3 trials at each combination of riser height and vision condition for an individual participant.

Since the purpose of this study was to discover how movement is scaled to a vertical environmental feature (the step), we developed a new measure, 'kneedrop', to capture the movements of the leg relative to the vertical.

1
2
3
4
5
6 To measure kneedrop (Fig 1a) we first defined the lower leg segment between the Knee
7 (LE marker) and Ankle (LM marker), and measured the angle ('swing') between this
8 segment and the vertical over the course of the stepping movement (Fig 1a). Changes in
9 this angle reflect not only flexion at the knee but also the orientation of the body relative
10 to vertical. For all participants, swing showed the same characteristic pattern during
11 descent – the leg swings outwards to a peak, then swings inwards again. We defined
12 swing peak as the point at which the rate of change in swing angle approached zero (was
13 1.5 deg/s or less). We then calculated the vertical position of the knee (LE marker) as the
14 body descended the step. 'Kneedrop' was defined as the knee's vertical descent from its
15 maximum height to the swing peak. In other words it measures how far the knee has
16 dropped vertically from its peak, at the time when the leg has ceased to swing outwards
17 and is beginning to swing back.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 We hypothesised that the location of the swing peak might change with riser height but
37 that this would depend on the visual information available. To test this we conducted a
38 repeated measures ANOVA on kneedrop with factors riser height (shallow, medium,
39 deep) and vision condition (V,NV,M). Significant results were followed up with further
40 ANOVAs to determine the source of the difference; finally a repeated measures ANOVA
41 on V trials with factor riser height confirmed scaling in the vision condition. We
42 excluded from the dataset any trials which failed to show a swing peak. Since any change
43 in the kneedrop measure might result from a change in knee peak height, we also
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 measured the correlation between each participant's knee peak height and kneedrop on V
4
5 trials.
6
7

8
9
10 We next examined the possibility of learning during the experiment, using a repeated
11
12 measures analysis at each level of riser height, with factors trial number (1,2,3) and
13
14 vision condition (V,M). As an index of overall movement efficiency we analysed
15
16 movement duration in a repeated measures riser height by vision condition (V,M)
17
18 ANOVA. We defined movement onset when the heel was raised 5 mm above the upper
19
20 platform and movement end when the toe was 5 mm above the lower platform. If
21
22 monocular viewing caused systematic misjudgements of riser height, one might expect a
23
24 greater incidence of high-impact, high-velocity landings on the lower platform in the M
25
26 condition than in the V condition. To test whether this was the case we conducted a
27
28 repeated measures riser height by vision condition (V,M) ANOVA on landing speed (MH
29
30 marker resultant speed in three dimensions at movement end).
31
32
33
34
35
36
37
38

39 We expected any effects of riser height to be monotonic, so in all ANOVAs we report
40
41 linear contrasts for height effects unless otherwise stated. For the same reason we report
42
43 linear contrasts for effects of trial number. Main effects are reported for all other factors
44
45 and interactions.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

Kneedrop

14/270 trials were excluded from analysis of kneedrop because swing showed no peak (i.e. kneedrop was not measurable). Figure 1b shows mean kneedrop data from the remaining trials, across all participants, riser heights and vision conditions. Figure 1c shows 'swing' angle unfolding over space in the vision condition, for a sample trial at each riser height. Peak swing occurs further down the step for deep than shallow steps (kneedrop is larger).

An ANOVA including all riser heights and vision conditions showed a vision-condition by riser-height interaction ($F[4,36] = 12.0, p < .001$) as well as effects of vision-condition ($F[2,18] = 3.9, p < .04$) and riser-height ($F[1,9] = 81.9 < .001$). Similarly an ANOVA including riser-height and vision conditions V, NV showed a vision-condition by riser height interaction ($F[2,18] = 27.7, p < .001$) and effects of vision-condition ($F[1,9] = 7.6, p < .03$) and riser height ($F[1,9] = 64.4 < .001$). An ANOVA including riser-height and vision conditions V, M showed a vision-condition by riser-height interaction ($F[2,18] = 4.2, p < .04$), an effect of riser-height ($F[1,9] = 138.7 < .001$), but no effect of vision condition ($F[1,9] = 1.2, p > .3$). An ANOVA with factor riser-height on V trials showed an effect of riser-height ($F[1,9] = 82.9, p < .001$). Taken together, these results show that in the vision condition participants scale their kneedrop to riser height. This scaling is significantly reduced either with no vision or with monocular viewing. Nine participants showed no significant correlation between knee peak height and kneedrop and one

participant showed a negative correlation, so increases in knee peak height could not account for the increases in kneedrop we found.

Learning (Fig 1d) was assessed at each value of riser height. For shallow trials, there was no effect of vision condition (V,M) on kneedrop ($F[1,5] = 1.2, p > 0.3$), an effect of trial number ($F[1,5] = 11.5, p < .02$), with kneedrop reducing as the experiment progressed, and no interaction ($F[2,10] = 0.3, p > 0.7$). Only 6 participants were included in this analysis, since the other 4 had at least one trial on which swing did not peak. For trials with medium step height, there was no effect of vision ($F[1,9] = 0.3, p > 0.5$), an effect of trial number ($F[1,9] = 5.2, p < .05$), with kneedrop reducing as the experiment progressed, and no vision condition by trial number interaction ($F[2,18] = 1.9, p > .1$). For trials with the maximum step height, there was an effect of vision condition ($F[1,9] = 6.4, p < .04$), no effect of trial number ($F[1,9] = 1.6, p > .2$) and no vision condition by trial number interaction ($F[2,18] = 0.7, p > 0.5$).

Secondary measures

An ANOVA on total movement duration (Table 1), showed a significant increase with riser height ($F[1,9] = 54.4, p < 0.001$), no effect of vision condition (V,M) ($F[1,9] = 2.9, p > 0.1$) and no interaction ($F[2,18] = 2.1, p > 0.1$).

Mean landing speed (Table 1) increased with riser height in both V and M conditions. A repeated measures ANOVA on V and M trials showed an effect of riser height ($F[1,9] = 35.8, p < .001$), no effect of vision condition ($F[1,9] = 3.9, p > .08$), and no interaction ($F[1.25,18] = 0.1, p > 0.9$, Greenhouse-Geisser corrected) on landing speed. Thus landing

1
2
3 speed is not larger on M trials than V trials as one might expect if participants were more
4 often making misjudgements on M trials.
5
6
7
8
9

10 11 **Discussion**

12
13
14 We developed a new paradigm to examine the visual control of stepping down a single
15 step. Visual control of the leg's movement during step descent can be captured by the
16 kinematic measure 'kneedrop', the distance dropped by the knee from its peak height to
17 the point where the calf segment reaches its maximum outwards 'swing'. This parameter
18 must be under visual control since (a) its value scales to the riser height of the step, which
19 in our paradigm must be gained using visual information since participants have no non-
20 visual cues to it; (b) scaling does not occur when participants are blindfolded. Scaling is
21 impaired under monocular viewing conditions.
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 This scaling process makes the stepping-down movement an efficient one by combining
37 horizontal and vertical translation in stepping down. Like forward stepping (Lyon & Day
38 1997), it seems that stepping down can be achieved as a controlled fall when the
39 appropriate visual information is present. Kneedrop is not the only measure of visual
40 control in step descent. For example future studies should investigate the relation of the
41 swing peak to pre-landing EMG activity (e.g. Santello, 2005). However, unlike these
42 EMG measures, kneedrop can be reliably extracted in degraded visual conditions, which
43 will allow future experiments to investigate the sources of visual information important
44 for controlling descent.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 While one might have expected monocular viewing to have some general effect on the
4 duration of the step, this was not affected. In contrast covering one eye impaired scaling
5 of kneedrop to step height. One interpretation of this is that binocular information is used
6 to perceive the depth of the step and scale movements to riser height. Thus removing
7 binocular information should cause misperceptions of target distance, which could be
8 responsible for the reduction in scaling we found. An alternative possibility is that with
9 monocular viewing, participants correctly perceive the depth of the step, but add some
10 margin for error in their movement parameters because visual uncertainty caused by
11 reduced field of view leads to cautious movement planning. This kind of effect has been
12 shown in reaching studies (Loftus, Murphy, McKenna & Mon-Williams, 2004). In the
13 present study informal observations and comments made by participants suggested that
14 some depth misperceptions occurred. If depth misjudgments occurred in such a task,
15 future studies could use synoptic viewing (Koenderink, van Doorn & Kappers, 1994) to
16 selectively remove binocular disparity, or prisms to selectively manipulate vergence
17 information, showing which binocular cues were most important for these distance
18 estimates. However, our current results are most consistent with a 'safety strategy'
19 account. Little learning occurred, and when it did it tended towards caution as the
20 experiment progressed. Likewise landing speeds were not high as 'undershooting' the
21 target step would predict. The specific safety strategy used by our participants was to
22 tend towards the mean value of the step depths encountered (or the riser height of an
23 average step, since this was approximated by our medium step). This kind of 'contraction
24 bias' strategy has also been reported in open-loop reaching movements (Tresilian *et al*,
25 1999).
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Our findings are consistent with the results of the few other studies on monocular
4 walking, which report interruptions to locomotor variables with monocular viewing
5 conditions (Patla *et al*, 2002, Chajka *et al*, 2007). In these studies it may also be the case
6 that monocular viewing caused participants to add a safety margin onto their estimates
7 because of a reduced field of view. Indeed Patla *et al* found that participants walking
8 monocularly over an obstacle increased toe clearance over the obstacle, which represents
9 a safety margin during obstacle crossing.
10
11
12
13
14
15
16
17
18
19

20
21
22 In summary, our novel kinematic measure provides a useful tool for assessing the
23 sensitivity of stepping actions to environmental parameters during stair descent,
24 analogous to measures developed for analysing single steps forward, medially or upward
25 over an obstacle. In particular it shows that stepping actions are regulated by visual
26 information about riser height, and demonstrates the kind of response that participants
27 make when visual information is degraded or removed during a step down.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Legends

Fig 1a: 'Swing' is the angle between the calf segment and the vertical. Kneedrop is the distance that the knee descends from its peak while the leg swings outwards. **Fig 1b:** Mean and standard errors of kneedrop in vision (V), monocular (M) and no vision (NV) conditions. **Fig 1c:** Example trials from one participant. As the leg swings towards vertical (x-axis), the knee drops (y-axis). When the leg is closest to vertical, the knee has dropped further for deep steps than shallow. **Fig 1d:** Learning across trials in V and M conditions.

Table 1: Mean and standard errors of movement duration and landing speed in V and M conditions.

References

Chajka, K., Vecellio, E., Hayhoe, M., & Gillam, B. (2007). The Role of Binocular Vision in Navigating Obstacles. *J. Vision*, 7:119a.

Craik RL, Cozzens BA, Freedman W (1982) The role of sensory conflict on stair descent performance in humans. *Exp Brain Res* 45:399-409

Institute for Perception TNO (1972) TNO Test for Stereoscopic Vision. Laméris: Utrecht, The Netherlands.

Koenderink JJ, van Doorn AJ, Kappers AM (1994) On so-called paradoxical monocular stereoscopy. *Perception* 23:583-594

Loftus, A., Murphy, S., McKenna, I., & Mon-Williams, M. (2004). Reduced Fields of View Are Neither Necessary nor Sufficient for Distance Underestimation but Reduce Precision and May Cause Calibration Problems. *Exp. Brain Res.*, 158:328-335

Lyon, I. N. & Day, B. L. (1997). Control of Frontal Plane Body Motion in Human Stepping. *Exp. Brain Res.*, 115:345-356

McFadyen BJ, Winter DA (1988) An integrated biomechanical analysis of normal stair ascent and descent. *J Biomech* 21:733-744

Patla AE, Niechwiej E, Racco V, Goodale MA (2002) Understanding the contribution of binocular vision to the control of adaptive locomotion. *Exp Brain Res* 142:551-561

Patla, A. E., Prentice, S. D., Robinson, C., & Neufeld, J. (1991). Visual Control of Locomotion: Strategies for Changing Direction and for Going Over Obstacles. *J. Exp. Psychol. Hum. Percept. Perform.*, 17:603-634

Protopapadaki A, Drechsler WI, Cramp MC, Coutts FJ, Scott OM (2007) *Clin. Biomech.* 22:203-210

Riener R, Rabuffetti M, Frigo C (2002) Stair ascent and descent at different inclinations. *Gait. Posture* 15:32-44

Santello M (2005) Review of motor control mechanisms underlying impact absorption from falls. *Gait. Posture* 21:85-94


Startzell, JK., Owens, DA., Mulfinger, LM. & Cavanagh, PR. (2000). Stair negotiation in older people: a review. *J Am Geriatr Soc*, 48:567-80

Tresilian, J. R., Mon-Williams, M., & Kelly, B. M. (1999). Increasing Confidence in Vergence As a Cue to Distance. *Proc. Biol. Sci.*, 266:39-44

1
2
3
4
5 Warren WH, Jr. (1984) Perceiving affordances: visual guidance of stair climbing. *J Exp*
6 *Psychol Hum Percept Perform* 10:683-703
7

8 Zachazewski JE, Riley PO, Krebs DE (1993) Biomechanical analysis of body mass
9 transfer during stair ascent and descent of healthy subjects. *J Rehabil Res Dev* 30:412-
10 422
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review



343x379mm (72 x 72 DPI)

	Vision (V) trials			Monocular (M) trials		
	<i>shallow</i>	<i>medium</i>	<i>deep</i>	<i>shallow</i>	<i>medium</i>	<i>deep</i>
Movement duration (sec)	.60 (.03)	.68 (.03)	.74 (.03)	.57 (.02)	.75 (.06)	.85 (.08)
Landing speed (m/sec)	.29 (.03)	.33 (.03)	.49 (.03)	.25 (.02)	.31 (.02)	.48 (.03)

For Peer Review