

Durham Research Online

Deposited in DRO:

29 May 2008

Version of attached file:

Published Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Aharonian,, F. and Akhperjanian, A. G. and Aye, K.-M. and Bazer-Bachi, A. R. and Beilicke, M. and Benbow, W. and Berge, D. and Berghaus, P. and Bernlöhr, K. and Bolz, O. and Boisson, C. and Borgmeier, C. and Breitling, F. and Brown, A. M. and Bussons Gordo, J. and Chadwick, P. M. and Chitnis, V. R. and Chounet, L.-M. and Cornils, R. and Costamante, L. and Degrange, B. and Djannati-Ataï, A. O'C. and Drury, L. and Ergin, T. and Espigat, P. and Feinstein, F. and Fleury, P. and Fontaine, G. and Funk, S. and Gallant, Y. and Giebels, B. and Gillessen, S. and Goret, P. and Guy, J. and Hadjichristidis, C. and Hauser, M. and Heinzlmann, G. and Henri, G. and Hermann, G. and Hinton, J. A. and Hofmann, W. and Holleran, M. and Horns, D. and de Jager, O. C. and Jung, I. and Khélifi, B. and Komin, N. and Konopelko, A. and Latham, I. J. and Le Gallou, R. and Lemoine, M. and Lemièrre, A. and Leroy, N. and Lohse, T. and Marcowith, A. and Masterson, C. and McComb, T. J. L. and de Naurois, M. and Nolan, S. J. and Noutsos, A. and Orford, K. J. and Osborne, J. L. and Ouchrif, M. and Panter, M. and Pelletier, G. and Pita, S. and Pohl, M. and Pühlhofer, G. and Punch, M., M. and Raubenheimer, B. C. and Raue, M. and Raux, J. and Rayner, S. M. and Redondo, I. and Reimer, A. and Reimer, O. and Ripken, J. and Rivoal, M. and Rob, L. and Rolland, L. and Rowell, G. and Sahakian, V. and Saugé, L. and Schlenker, S. and Schlickeiser, R. and Schuster, C. and Schwanke, U. and Siewert, M. and Sol, H. and Steenkamp, R. and Stegmann, C. and Tavernet, J.-P. and Théoret, C. G. and Tluczykont, M. and van der Walt, D. J. and Vasileiadis, G. and Vincent, P. and Visser, B. and Völk, H. J. and Wagner, S. J. (2004) 'Very high energy gamma rays from the direction of Sagittarius A*.', *Astronomy astrophysics.*, 425 (1). L13-L17.

Further information on publisher's website:

<https://doi.org/10.1051/0004-6361:200400055>

Publisher's copyright statement:

Reproduced with permission from *Astronomy Astrophysics*, © ESO

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Durham University Library, Stockton Road, Durham DH1 3LY, United Kingdom

Tel : +44 (0)191 334 3042 | Fax : +44 (0)191 334 2971

<https://dro.dur.ac.uk>

Very high energy gamma rays from the direction of Sagittarius A*

F. Aharonian¹, A. G. Akhperjanian², K.-M. Aye³, A. R. Bazer-Bachi⁴, M. Beilicke⁵, W. Benbow¹, D. Berge¹, P. Berghaus⁶, K. Bernlöhr^{1,7}, O. Bolz¹, C. Boisson⁸, C. Borgmeier⁷, F. Breitling⁷, A. M. Brown³, J. Bussons Gordo⁹, P. M. Chadwick³, V. R. Chitnis^{10,20,*}, L.-M. Chounet¹¹, R. Cornils⁵, L. Costamante^{1,20}, B. Degrange¹¹, A. Djannati-Atai⁶, L. O'C. Drury¹², T. Ergin⁷, P. Espigat⁶, F. Feinstein⁹, P. Fleury¹¹, G. Fontaine¹¹, S. Funk¹, Y. Gallant⁹, B. Giebels¹¹, S. Gillessen¹, P. Goret¹³, J. Guy¹⁰, C. Hadjichristidis³, M. Hauser¹⁴, G. Heinzlmann⁵, G. Henri¹⁵, G. Hermann¹, J. A. Hinton¹, W. Hofmann¹, M. Holleran¹⁶, D. Horns¹, O. C. de Jager¹⁶, I. Jung^{1,14,***}, B. Khélifi¹, Nu. Komin⁷, A. Konopelko^{1,7}, I. J. Latham³, R. Le Gallou³, M. Lemoine¹¹, A. Lemièrè⁶, N. Leroy¹¹, T. Lohse⁷, A. Marcowith⁴, C. Masterson^{1,20}, T. J. L. McComb³, M. de Naurois¹⁰, S. J. Nolan³, A. Noutsos³, K. J. Orford³, J. L. Osborne³, M. Ouchrif^{10,20}, M. Panter¹, G. Pelletier¹⁵, S. Pita⁶, M. Pohl^{17,***}, G. Pühlhofer^{1,14}, M. Punch⁶, B. C. Raubenheimer¹⁶, M. Raue⁵, J. Raux¹⁰, S. M. Rayner³, I. Redondo^{11,20,†}, A. Reimer¹⁷, O. Reimer¹⁷, J. Ripken⁵, M. Rivoal¹⁰, L. Rob¹⁸, L. Rolland¹⁰, G. Rowell¹, V. Sahakian², L. Saugé¹⁵, S. Schlenker⁷, R. Schlickeiser¹⁷, C. Schuster¹⁷, U. Schwanke⁷, M. Siewert¹⁷, H. Sol⁸, R. Steenkamp¹⁹, C. Stegmann⁷, J.-P. Tavernet¹⁰, C. G. Théoret⁶, M. Tluczykont^{11,20}, D. J. van der Walt¹⁶, G. Vasileiadis⁹, P. Vincent¹⁰, B. Visser¹⁶, H. J. Völk¹, and S. J. Wagner¹⁴

¹ Max-Planck-Institut für Kernphysik, PO Box 103980, 69029 Heidelberg, Germany
e-mail: Jim.Hinton@mpi-hd.mpg.de

² Yerevan Physics Institute, 2 Alikhanian Brothers St., 375036 Yerevan, Armenia

³ University of Durham, Department of Physics, South Road, Durham DH1 3LE, UK

⁴ Centre d'Étude Spatiale des Rayonnements, CNRS/UPS, 9 Av. du Colonel Roche, BP 4346, 31029 Toulouse Cedex 4, France

⁵ Universität Hamburg, Inst. für Experimentalphysik, Luruper Chaussee 149, 22761 Hamburg, Germany

⁶ Physique Corpusculaire et Cosmologie, IN2P3/CNRS, Collège de France, 11 place Marcelin Berthelot, 75231 Paris Cedex 05, France

⁷ Institut für Physik, Humboldt-Universität zu Berlin, Newtonstr. 15, 12489 Berlin, Germany

⁸ LUTH, UMR 8102 du CNRS, Observatoire de Paris, Section de Meudon, 92195 Meudon Cedex, France

⁹ Groupe d'Astroparticules de Montpellier, IN2P3/CNRS, Université Montpellier II, CC85, Place Eugène Bataillon, 34095 Montpellier Cedex 5, France

¹⁰ Laboratoire de Physique Nucléaire et de Hautes Énergies, IN2P3/CNRS, Universités Paris VI & VII, 4 place Jussieu, 75231 Paris Cedex 05, France

¹¹ Laboratoire Leprince-Ringuet, IN2P3/CNRS, École Polytechnique, 91128 Palaiseau, France

¹² Dublin Institute for Advanced Studies, 5 Merrion Square, Dublin 2, Ireland

¹³ Service d'Astrophysique, DAPNIA/DSM/CEA, CE Saclay, 91191 Gif-sur-Yvette, France

¹⁴ Landessternwarte, Königstuhl, 69117 Heidelberg, Germany

¹⁵ Laboratoire d'Astrophysique de Grenoble, INSU/CNRS, Université Joseph Fourier, BP 53, 38041 Grenoble Cedex 9, France

¹⁶ Unit for Space Physics, North-West University, Potchefstroom 2520, South Africa

¹⁷ Institut für Theoretische Physik, Lehrstuhl IV: Weltraum und Astrophysik, Ruhr-Universität Bochum, 44780 Bochum, Germany

¹⁸ Institute of Particle and Nuclear Physics, Charles University, V Holesovickach 2, 180 00 Prague 8, Czech Republic

¹⁹ University of Namibia, Private Bag 13301, Windhoek, Namibia

²⁰ European Associated Laboratory for Gamma-Ray Astronomy, jointly supported by CNRS and MPG

Received 29 June 2004 / Accepted 12 August 2004

* Now at Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400 005, India.

** Now at Washington Univ., Department of Physics, 1 Brookings Dr., CB 1105, St. Louis, MO 63130, USA.

*** Now at Department of Physics and Astronomy, Iowa State University, Ames, Iowa 50011-3160, USA.

† Now at Department of Physics and Astronomy, Univ. of Sheffield, The Hicks Building, Hounsfield Road, Sheffield S3 7RH, UK.

Abstract. We report the detection of a point-like source of very high energy (VHE) γ -rays coincident within $1'$ of Sgr A*, obtained with the HESS array of Cherenkov telescopes. The γ -rays exhibit a power-law energy spectrum with a spectral index of $-2.2 \pm 0.09 \pm 0.15$ and a flux above the 165 GeV threshold of $(1.82 \pm 0.22) \times 10^{-7} \text{ m}^{-2} \text{ s}^{-1}$. The measured flux and spectrum differ substantially from recent results reported in particular by the CANGAROO collaboration.

Key words. gamma-rays: observations – Galaxy: centre

1. Introduction

The Galactic Centre (GC) region (Melia & Falcke 2001) harbours a variety of potential sources of high-energy radiation including the supermassive black hole Sgr A* of $2.6 \times 10^6 M_{\odot}$ (see e.g. Schödel et al. 2002), which has been identified as a faint source of X-rays (Baganoff et al. 2003) and infrared radiation (Genzel et al. 2003). Emission from Sgr A* is presumably powered by the energy released in the accretion of stellar winds onto the black hole (Melia 1992; Yusef-Zadeh et al. 2000; Yuan et al. 2003).

High (Mayer-Hasselwander et al. 1998) and very high (Tsuchiya et al. 2004; Kosack et al. 2004) energy γ -ray emission have also been detected from the GC region. The γ -radiation could result from acceleration of electrons or protons in shocks in these winds, in the accretion flow or in nearby supernova remnants, followed by interactions of accelerated particles with ambient matter or radiation. Alternative mechanisms include the annihilation of dark matter particles accumulating at the GC (Bergström et al. 1998; Ellis et al. 2002; Gnedin & Primack 2003) or curvature radiation of protons near the black hole (Levinson 2000).

2. Observations and results

The observations presented here were obtained in Summer 2003 with the High Energy Stereoscopic System (HESS), consisting of four imaging atmospheric Cherenkov telescopes (Hofmann 2003; Bernlöhr et al. 2003; Vincent et al. 2003) in Namibia, at $23^{\circ}16' \text{ S } 16^{\circ}30' \text{ E}$. At this time, two of the four telescopes were operational, the other two being under construction. During the first phase of the measurements (June 6 to July 7, 2003), the telescopes were operated independently and images were combined offline using GPS time stamps (4.7 h on source, “June/July” data set). In the second phase (July 22 to August 29, 2003), a hardware coincidence required shower images simultaneously in both telescopes (11.8 h on source, “July/August” data set). The resulting background suppression allowed us to lower the telescope trigger thresholds, yielding a post-cuts energy threshold of 165 GeV (for typical Sgr A* zenith angles of 20°) as compared to 255 GeV for the “June/July” data set.

Shower images are parametrised by their centres of gravity and second moments, followed by the stereoscopic reconstruction of shower geometry, providing an angular resolution of $\approx 0.1^{\circ}$ for individual γ -rays. γ -ray candidates are selected based on the shape of shower images, allowing effective suppression of cosmic-ray showers. The γ -ray energy is estimated from the image intensity and the reconstructed shower geometry, with a typical resolution of 15–20%.




Fig. 1. Angular distribution of γ -ray candidates for a 3° field of view centred on Sgr A*. Both data sets (“June/July” and “July/August”) are combined, employing tight cuts to reduce the level of background. The significance of the feature extending along the Galactic Plane is under investigation.

The GC region is characterised by high night-sky brightness (NSB), varying across the field of view and potentially interfering with image reconstruction. Simulations of a range of NSB levels show, however, that the stereoscopic reconstruction is insensitive to this feature, resulting in variations of the measured flux and spectrum that are well within the systematic errors quoted here.

The performance and stability of HESS have been confirmed by observations of the Crab Nebula (a standard candle in γ -ray astronomy). The absolute calibration of the instrument has been verified using muon images (Leroy et al. 2003) which provide a measurement of the absolute photon detection efficiency, and by the measured cosmic ray detection rates (Funk et al. 2004), which are in excellent agreement with simulations.

Figure 1 shows the distribution of γ -ray candidates for a 3° window around Sgr A*. A clear excess of events in the Sgr A* region is observed. Here, tight γ -ray selection cuts are applied to minimise background at the expense of γ -ray efficiency. For the analysis of the flux and spectrum of the central point source, looser cuts are used which reject 96% of the cosmic-ray background and retain 50% of the γ -rays. Using a ring around the assumed source location to estimate background, we find – with loose cuts – a 6.1σ excess in the “June/July” data set and a 9.2σ excess in the “July/August” data set, both centred on Sgr A*. The γ -ray excess is located at RA $17^{\text{h}}45^{\text{m}}41.3^{\text{s}} \pm 2.0^{\text{s}}$, Dec $-29^{\circ}0'22'' \pm 32''$, or $l = 359^{\circ}56'53''$, $b = -0^{\circ}2'57''$, within




Fig. 2. Centre of gravity of the VHE signal (triangle), superimposed on a 8.5' by 8.5' Chandra X-ray map (Muno et al. 2003) of the GC. The location of Sgr A* is indicated by a cross. The contour lines indicate the 68% and 95% confidence regions for the source position, taking into account systematic pointing errors of 20". The white dashed line gives the 95% confidence level upper limit on the rms source size. The resolution for individual VHE photons – as opposed to the precision for the centre of the VHE signal – is 5.8' (50% containment radius).

$14 \pm 30''$ in b and $12 \pm 30''$ in l from Sgr A* (Fig. 2). There is no evidence in our data for an energy dependence of this position. A conservative pointing error of less than 20" in RA and Dec has been estimated using stars (Gillissen et al. 2003), and verified by reconstructing the location of known VHE sources such as the Crab Nebula and the AGN PKS 2155–304.

Given the high density of potential sources over the central square degree of the Galaxy, an important question is whether the VHE γ -ray signal shows signs of source extension. Figure 3 shows the angular distribution of detected γ -rays relative to Sgr A*. The width and shape of this distribution are consistent with point source simulations. These simulations have been verified using the strong signals from the Crab Nebula and PKS 2155–304. Assuming a Gaussian distribution of source brightness, $\rho \propto \exp(-\theta^2/2\sigma_{\text{source}}^2)$, we find an upper limit $\sigma_{\text{source}} < 3'$ for the source size (95% CL), corresponding to < 7 pc at the distance of the GC (dashed white line in Fig. 2). The apparent point-like nature of the central source does not exclude the possibility of non-azimuthally symmetric tails in the emission.

The measured energy spectrum is shown in Fig. 4. Data are fit by a power-law, $F(E) = F_0 E_{\text{TeV}}^{-\alpha}$, with a spectral index $\alpha = 2.21 \pm 0.09$ and $F_0 = (2.50 \pm 0.21) \times 10^{-8} \text{ m}^{-2} \text{ s}^{-1} \text{ TeV}^{-1}$ for the “July/August” data set (full circles), with a $\chi^2/\text{d.o.f.}$ of 0.6. The flux above the 165 GeV threshold is $(1.82 \pm 0.22) \times 10^{-7} \text{ m}^{-2} \text{ s}^{-1}$, equivalent to 5% of the Crab Nebula flux at this threshold. The smaller “June/July” data set gives consistent results ($\alpha = 2.11 \pm 0.19$ and $F_0 = (2.76 \pm 0.33) \times 10^{-8} \text{ m}^{-2} \text{ s}^{-1} \text{ TeV}^{-1}$). We estimate systematic errors of $\Delta\alpha \approx 0.15$ and $\Delta F/F \approx 25\%$, with the latter mainly governed by the precision of the energy




Fig. 3. Angular distribution of VHE γ -rays relative to the location of Sgr A*. Inset: distributions in θ^2 where θ is the angle between the γ -ray direction and Sgr A*; a uniform background results in a flat distribution in θ^2 . Full points: signal region; open points: background region. The main figure shows background-subtracted excess counts. The solid line indicates the distribution expected for a point source of γ -rays at the position of Sgr A*.

calibration of the instrument. The energy reconstruction and flux determination have been tested with the Crab Nebula; we reconstruct a power law with index $\alpha = 2.63 \pm 0.04$ and a flux above 1 TeV of $(1.98 \pm 0.08) \times 10^{-7} \text{ m}^{-2} \text{ s}^{-1}$ for the Crab, in very good agreement with previous measurements (see Aharonian et al. 2000, and references therein). Fitting the GC γ -ray spectrum as a power law with an exponential cutoff, we find a lower limit for the cutoff energy of 4 TeV. Within statistics, there are no indications for time variability of the GC signal.

3. Discussion and conclusions

The CANGAROO collaboration recently reported the detection of sub-TeV gamma rays from within 0.1° of the GC based on 67 h of (on-source) data taken in July 2001 and July/August 2002 (Tsuchiya et al. 2004). The reported spectrum is very steep, $F(E) \propto E^{-4.6 \pm 0.5}$. The rather hard HESS spectra are obviously not consistent with the steep spectrum obtained with CANGAROO-II (Fig. 4); the large flux at low energies implied by the CANGAROO result would have been detected with HESS in a matter of minutes. At higher energies, above 2.8 TeV, a marginal detection with a significance of 3.7σ resulting from 26 h of large-zenith-angle observations in the years 1995 through 2003 was reported by the Whipple collaboration (Kosack et al. 2004), consistent with Sgr A* within the 15' 95% C.L. error circle. The Whipple flux is a factor 3 above that implied by our spectra. Taking all data at face value, one would conclude that the source underwent significant changes over the timescale of one year (2002 to 2003). However, this seems unlikely since none of the individual experiments detects significant variability. Implications of the CANGAROO and Whipple data are discussed in Hooper et al. (2004).

At lower energies around 100 MeV, the EGRET instrument detected a strong excess from the central part of the Galaxy




Fig. 4. Energy spectrum $E^2 dN/dE$ of γ -rays from the Galactic Centre. Full circles: HESS “July/August 2003” data set. Full triangles: HESS “June/July 2003” data set. The line indicates a power-law fit to the “July/August” spectrum. Open squares: CANGAROO-II spectrum from Summer 2001 and 2002 (Tsuchiya et al. 2004). Open triangle: Whipple flux from 1995 through 2003 (Kosack et al. 2004), converted to a differential flux at the peak detection energy assuming a Crab-like spectrum. The inset shows the EGRET flux from 1991 to 1996 (Mayer-Hasselwander et al. 1998) (circles) compared to fits to the CANGAROO-II (dashed line) and HESS (solid line) spectra. Due to the poor angular resolution of EGRET (1°) the flux shown may include other sources.

(Mayer-Hasselwander et al. 1998), consistent within its error circle with Sgr A*, but with angular resolution of 1° also covering other potential sources (and a solid angle ~ 100 times larger than the emission region seen by HESS). In the analysis of a high-energy sub-sample of EGRET data (Hooper & Dingus 2002), the source was found to be offset from the GC by 0.21° , excluding Sgr A* at 99.9% C.L.

Models for the wide-band spectra of Sgr A* include Advection Dominated Accretion Flow (ADAF) models (Narayan et al. 1998; Yuan et al. 2003), possibly combined with a jet extracting energy from the accretion disk (Yuan et al. 2002). Shocks in the accretion flow (Markoff et al. 1997) or in the jet could accelerate particles. γ -rays are generated in proton interactions, but predicted spectra tend to fall off rapidly in the TeV region (Markoff et al. 1997).

Another source of VHE γ -rays should be diffuse emission from the entire central region, in which case year time-scale variability should not occur. γ -rays may result from interactions of accelerated protons and nuclei (Fatuzzo & Melia 2003) with the ambient matter with a density as large as $n = 10^3 \text{ cm}^{-3}$ (Maeda et al. 2002). Only modest overall energy, $W_p \approx 5 \times 10^{47} (10^3 \text{ cm}^{-3}/n) \text{ erg}$, in TeV protons is needed to explain the observed γ -ray flux from this region (the 1–10 TeV luminosity is $\sim 10^{35} \text{ ergs/s}$). An obvious candidate for the proton accelerator could be the young (10^4 yr) and unusually powerful (total explosion energy $\approx 4 \times 10^{52} \text{ erg}$) supernova remnant Sgr A East

(Maeda et al. 2002). The measured spectral index of TeV emission, $\alpha \approx 2.2$, is close to the spectrum of shock-accelerated particles. For a 10^4 yr source age, the modest source extension and the hard spectrum imply that particle diffusion in the central region proceeds much slower compared to the diffusion in the Galactic Disk. While detailed modelling remains to be done, estimates show that for magnetic fields up to $\sim 1 \text{ mG}$ the X-ray and radio emission resulting from secondary electrons generated in interactions of such accelerated protons are below the measured diffuse luminosities integrated over Sgr A East. This consistency criterion is, however, much more challenging if one considers Sgr A* as the source, with its significantly larger magnetic fields and lower (quiescent-state) X-ray flux. In either case, the explanation of the EGRET flux (Fig. 4) requires a second source component with a cutoff close to the highest EGRET energies, located well within the HESS field of view (and therefore excluded as a strong TeV source), but not necessarily coincident with Sgr A East or Sgr A*. We note that Tsuchiya et al. (2004) describe the EGRET and CANGAROO fluxes jointly in a model of a diffuse proton flux with a spectral cutoff of a few TeV, interacting with ambient gas. In such a model, one would, however, not expect fast variability.

Alternative mechanisms invoke the hypothetical annihilation of super-symmetric dark matter particles (Bergström et al. 1998; Ellis et al 2002; Gnedin & Primack 2003) or curvature radiation of protons in the vicinity of the central super-massive black hole (Levinson 2000).

The spectrum of γ -rays from hypothetical annihilation of neutralinos of mass M_χ consists of a γ -ray continuum and two lines at $E = M_\chi$ and $E = M_\chi(1 - m_Z^2/4M_\chi^2)$. The continuum spectra generated by the DarkSusy program (Gondolo et al. 2004) are well approximated by $F \sim E^{-\alpha} e^{-(E/M_{\text{cut}})}$ with $\alpha = 2.2 \dots 2.4$ and $M_{\text{cut}} = 0.15 \dots 0.3 M_\chi$ depending on the annihilation channel¹. Assuming that the observed γ -rays represent a continuum annihilation spectrum, the lower limit of 4 TeV on the cutoff implies $M_\chi > 12 \text{ TeV}$, a range which is presently disfavoured due to particle physics and cosmology arguments (Ellis et al. 2003). Supersymmetric dark matter annihilation as the main source of the observed γ -rays is therefore unlikely, but not excluded.

The spectrum of the proton curvature radiation (Levinson 2000) depends, to a large extent, on the configuration of magnetic fields near the gravitational radius of the black hole, and detailed predictions are lacking; as a characteristic feature, one would expect time variability. Further observations of the GC region are a high priority for HESS in the near future.

Acknowledgements. The support of the Namibian authorities and of the University of Namibia in facilitating the construction and operation of HESS is gratefully acknowledged, as is the support by the German Ministry for Education and Research (BMBF), the Max Planck Society, the French Ministry for Research, the CNRS-IN2P3 and the Astroparticle Interdisciplinary Programme of the CNRS, the

¹ Bergström et al. (1998) use $\alpha = 1.5$ and $M_{\text{cut}} = 0.13 M_\chi$ to describe spectra for $E < 0.5 M_\chi$, but with larger α values a better fit up to $E = 0.9 M_\chi$ is obtained.

UK Particle Physics and Astronomy Research Council (PPARC), the IPNP of the Charles University, the South African Department of Science and Technology and National Research Foundation, and by the University of Namibia. We appreciate the excellent work of the technical support staff in Berlin, Durham, Hamburg, Heidelberg, Palaiseau, Paris, Saclay, and in Namibia in the construction and operation of the equipment.

References

- Aharonian, F. A., Akhperjanian, A. G., Barrio, J. A., et al. 2000, *ApJ*, 539, 317
- Baganoff, F. K., Maeda, Y., Morris, M., et al. 2003, *ApJ*, 591, 891
- Bergström, L., Ullio, P., & Buckley, J. H. 1998, *A&A*, 335, 161
- Bernlöhr, K., Carrol, O., Cornils, R., et al. 2003, *A&A*, 409, 111
- Ellis, J. R., Feng, J. L., Ferstl, A., Matchev, K. T., & Olive, K. A. 2002, *Eur. Phys. J.*, C24, 311
- Ellis, J., Olive, K. A., Santoso, Y., & Spanos, V. C. 2003, *Phys. Lett. B*, 565, 176
- Fatuzzo, M., & Melia, F. 2003, *ApJ*, 596, 1035
- Funk, S., Hermann, G., Hinton, J., et al. 2004, *A&A*, submitted
- Genzel, R., Schödel, R., Ott, T., et al. 2003, *Nature*, 425, 934
- Gillessen, S. for the HESS collaboration 2003 Proc. 28th ICRC, Tsukuba (Tokyo: Univ. Academy Press), 2899
- Gnedin, O. Y., & Primack, J. R. 2003, *Phys. Rev. Lett.*, submitted [arXiv:astro-ph/0308385]
- Gondolo, P., Edsjö, J., Ullio, P., et al. 2004, *JCAP* 0407, 008
- Hofmann, W. 2003, Proc. 28th ICRC, Tsukuba (Tokyo: Univ. Academy Press), 2811
- Hooper, D., & Dingus, B. 2002 [arXiv:astro-ph/0212509]
- Hooper, D., de la Calle Perez, I., Silk, J., Ferrer, F., & Sarkar, S. 2004 [arXiv:astro-ph/0404205]
- Kosack, K., Badran, H. M., Bond, H. I., et al. 2004, *ApJ*, 608, L97
- Leroy, N., Bolz, O., Guy, J., et al. 2003, Proc. 28th ICRC, Tsukuba (Tokyo: Univ. Academy Press), 2895
- Levinson, A. 2000, *Phys. Rev. Lett.*, 85, 912
- Maeda, Y., Baganoff, F. K., Feigelson, E. D., et al. 2002, *ApJ*, 570, 671
- Markoff, S., Melia, F., & Sarcevic, I. 1997, *ApJ*, 489, L47
- Mayer-Hasselwander, H. A., Bertsch, D. L., Dingus, B. L., et al. 1998, *A&A*, 335, 161
- Melia, F. 1992, *ApJ*, 387, L25
- Melia, F., & Falcke, H. 2001, *ARA&A*, 39, 309
- Muno, M. P., Baganoff, F. K., Bautz, W. M., et al. 2003, *ApJ*, 589, 225
- Narayan, R., Mahadevan, R., Grindlay, J. E., Popham, R. G., & Gammie, C. 1998, *ApJ*, 492, 554
- Schödel, R., Ott, T., Genzel, R., et al. 2002, *Nature*, 419, 694
- Tsuchiya, K., Enomoto, R., Ksenofontov, L. T., et al. 2004, *ApJ*, 606, L115
- Vincent, P., Denance, J.-P., Huppert, J.-F., et al. 2003, Proc. 28th ICRC, Tsukuba (Tokyo: Univ. Academy Press), 2887
- Yuan, F., Markoff, S., & Falcke, H. 2002, *A&A*, 383, 854
- Yuan, F., Quataert, E., & Narayan, R. 2003, *ApJ*, 598, 301
- Yusef-Zadeh, F., Melia, F., & Wardle, M. 2000, *Science*, 287, 85