

Durham Research Online

Deposited in DRO:

16 June 2015

Version of attached file:

Other

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Ilderton, Anton and Mansfield, Paul (2005) 'Time evolution in string field theory and T-duality.', *Physics letters B.*, 607 (3-4). pp. 294-301.

Further information on publisher's website:

<http://dx.doi.org/10.1016/j.physletb.2004.12.066>

Publisher's copyright statement:

NOTICE: this is the author's version of a work that was accepted for publication in *Physics Letters B*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Physics Letters B*, 607, 3-4, 10 February 2005, 10.1016/j.physletb.2004.12.066.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Time evolution in string field theory and T-Duality

Anton Ilderton* and Paul Mansfield†

Centre for Particle Theory, University of Durham,
Durham, DH1 3LE, UK

Introduction

In the mechanics of particles and fields it is natural to consider the evolution in time of arbitrary configurations. In second quantised string theory this is not so straightforward, for example in Witten's theory [1] the natural time variable is that at the mid-point of the string rather than a global time for the whole string. In this letter we will construct the time evolution operator for second quantised strings by analogy with that for field theory. We begin by showing that when the field theory Schrödinger functional is written in terms of propagators expressed in first quantised form then these describe particles moving on a timelike orbifold $\mathbb{S}^1/\mathbb{Z}_2$. The first quantised propagators have an immediate generalisation to string theory, suggesting that the Schrödinger functional for second quantised strings can be expressed in terms of first quantised strings moving on this orbifold. To strengthen the analogy we give a graphical construction of the field theory Schrödinger functional which extends to both open and closed string theory. This avoids using a Lagrangian formulation of string field theory. Finally we study the effect of T-duality on time evolution and describe the nature of BRST invariance in our approach.

1 Time Evolution in QFT

Consider a bosonic scalar field ϕ in $D + 1$ dimensions. It will be convenient to work in a basis in which $\hat{\pi}(\mathbf{x}) = \dot{\phi}(\mathbf{x})$, the momentum canonically conjugate to the field (the problem of defining a momentum in string field theory goes hand in hand with the definition of a global time), is diagonal

$$\langle \pi | \hat{\pi}(\mathbf{x}) = \pi(\mathbf{x}) \langle \pi |, \quad i \frac{\delta}{\delta \pi(\mathbf{x})} \langle \pi | = \langle \pi | \hat{\phi}(\mathbf{x}).$$

*a.b.ilderton@dur.ac.uk, corresponding author

†p.r.w.mansfield@dur.ac.uk

a Lagrangian derivation we will give a graphical derivation of the field theory result which can be taken over into string theory.

We appeal to a fundamental property of field theory, which follows from the observation that paths from t_3 to t_1 must cross the plane at time t_2 for $t_3 > t_2 > t_1$, so that formally the sum over paths in (3) can be factorised,

$$\sum_{\text{paths AB}} e^{-\text{length(AB)}} = \sum_C \left(\sum_{\text{paths AC}} e^{-\text{length(AC)}} \right) \left(\sum_{\text{paths CB}} e^{-\text{length(CB)}} \right).$$

The explicit result, which we refer to as the gluing property, is

$$\int d^D \mathbf{y} G_0(\mathbf{x}_3, t_3; \mathbf{y}_2, t_2) \left(-i \overleftrightarrow{\frac{\partial}{\partial t_2}} \right) G_0(\mathbf{y}, t_2; \mathbf{x}_1, t_1) = G_0(\mathbf{x}_3, t_3; \mathbf{x}_1, t_1), \quad (4)$$

for $t_3 > t_2 > t_1$. More generally, if the endpoints x_3 and x_1 are on opposite sides of the plane at time t_2 , the propagators are glued to form the usual propagator. If they are on the same side gluing produces the image propagator G_I equal to the free space propagator for the points x_3 and the reflection of x_1 in the plane at t_2 . The cases are summarised below,

$$\int d^D \mathbf{y} G_0(\mathbf{x}_2, t_2; \mathbf{y}, t) \frac{\partial}{\partial t} G_0(\mathbf{y}, t; \mathbf{x}_1, t_1) = \begin{cases} \mp \frac{i}{2} G_0(\mathbf{x}_2, t_2; \mathbf{x}_1, t_1) & t_2 \geq t \geq t_1 \\ \mp \frac{i}{2} G_I(\mathbf{x}_2, t_2; \mathbf{x}_1, t_1) & t \geq t_1, t_2 \end{cases} \quad (5)$$

Applying (5) twice we obtain

$$\begin{aligned} \int d^D(\mathbf{x}_3, \mathbf{x}_2) G_0(\mathbf{x}_4, t_4, \mathbf{x}_3, t_3) \left(4 \frac{\partial^2}{\partial t_3 \partial t_2} G_0(\mathbf{x}_3, t_3, \mathbf{x}_2, t_2) \right) G_0(\mathbf{x}_2, t_2, \mathbf{x}_1, t_1) \\ = G_0(\mathbf{x}_4, t_4, \mathbf{x}_1, t_1) \quad \text{for } t_4 > t_3 > t_2 > t_1. \end{aligned}$$

Taking all the t_i to zero gives a useful relation which may be expressed as

where the heavy line is the plane at time $t = 0$, the unbroken line is the free space propagator and a black dot is -2 times a time derivative. Thus

From this we deduce that the inverse of the free space propagator at equal time is

We can now show that time evolution is captured by the gluing rules and the Feynman diagram expansion. Consider calculating the free theory two-point function at unequal times,

$$\langle \pi(\mathbf{x}, t)\pi(\mathbf{y}, 0) \rangle = \int \mathcal{D}(\pi_2, \pi_1)\Psi_0[\pi_2]\pi_2(\mathbf{x})\mathcal{S}[\pi_2, t; \pi_1, 0]\pi_1(\mathbf{y})\Psi_0[\pi_1]. \quad (8)$$

The vacuum wave functional $\Psi_0[\pi]$ can be constructed by requiring that it yield G_0 at equal times as a vacuum expectation value,

$$\begin{aligned} G_0(\mathbf{x}, 0; \mathbf{y}, 0) &= \langle \phi(\mathbf{x}, 0)\phi(\mathbf{y}, 0) \rangle = - \int \mathcal{D}\pi \Psi_0[\pi] \frac{\delta}{\delta\pi(\mathbf{x})} \frac{\delta}{\delta\pi(\mathbf{y})} \Psi_0[\pi] \\ \implies \Psi_0[\pi] &= \exp\left(- \text{diagram} \right). \end{aligned} \quad (9)$$

The π_1 integration in (8) is Gaussian in the free theory,

$$\begin{aligned} \int \mathcal{D}\pi_1 \pi_1(\mathbf{y}) \exp\left(\text{diagram}_1\right) \exp\left(-\frac{1}{2} \text{diagram}_2 - \text{diagram}_3\right) \\ = K^{-1}(\mathbf{y}, \mathbf{z}) \left(\text{diagram}_4\right) \exp\left(\frac{1}{2} \text{diagram}_5 K^{-1}(\mathbf{a}, \mathbf{b}) \text{diagram}_6\right). \end{aligned}$$

The symmetric operator K and it's inverse are

$$K(\mathbf{x}, \mathbf{y}) = 2 \left(\text{diagram}_7 + \text{diagram}_8 \right) \implies K^{-1}(\mathbf{x}, \mathbf{y}) = -\frac{1}{4} \left(\text{diagram}_9 + \frac{1}{4} \text{diagram}_{10} \right)$$

which can be checked using the gluing rules (5) and the corollary (7). The exponential term is

$$\begin{aligned} \frac{1}{2} \text{diagram}_{11} K^{-1} \text{diagram}_{12} &= 2 \left(\text{diagram}_{13} + \text{diagram}_{14} + \dots \right) \left(\text{diagram}_{15} - \text{diagram}_{16} \right) \\ &\quad \times \left(\text{diagram}_{17} + \text{diagram}_{18} + \dots \right) \\ &= \frac{1}{2} \left(\text{diagram}_{19} - \text{diagram}_{20} \right) \end{aligned}$$

which leaves us with another Gaussian integral in π ,

$$\begin{aligned}
\int \mathcal{D}\pi_2 \pi_2(\mathbf{x}) &= \frac{1}{4} \left(\text{diagram 1} - \text{diagram 2} \right) \text{diagram 3} \exp \left(-2 \text{diagram 4} \right) \\
&= \frac{1}{4} \left(\text{diagram 1} - \text{diagram 2} \right) \text{diagram 5} \left(\frac{1}{4} \text{diagram 6} \right) \\
&= \text{diagram 7} = \langle \pi(\mathbf{x}, t) \pi(\mathbf{y}, 0) \rangle
\end{aligned}$$

The diagrams are Feynman diagrams for a Gaussian integral. Diagram 1: A horizontal line from y to z at time 0, with a grey dot at y and a black dot at z , and a curved line connecting them. Diagram 2: A horizontal line from y to z at time $2t$, with a grey dot at y and a black dot at z , and a vertical dashed line from z to time 0. Diagram 3: A horizontal line from z to z at time 0, with a grey dot at z and a black dot at z , and a vertical dashed line from z to time t . Diagram 4: A horizontal line from π_2 to π_2 at time 0, with a grey dot at π_2 and a black dot at π_2 , and a curved line connecting them. Diagram 5: A horizontal line from y to z at time 0, with a grey dot at y and a black dot at z , and a curved line connecting them. Diagram 6: A horizontal line from w to x at time 0, with a grey dot at w and a black dot at x , and a curved line connecting them. Diagram 7: A horizontal line from x to y at time t , with a grey dot at x and a black dot at y , and a vertical dashed line from y to time 0.

We arrive at the Feynman diagram of the free propagator with the correct coefficient of unity; the grey dot denotes a time derivative (with no factor) which appears since we are computing correlation functions of $\pi = \dot{\phi}$.

So using the gluing property alone we have shown that the expression (1) for the Schrödinger functional leads to the correct result for the two point function at unequal times. This argument is invertible; If we know the two point function we can construct the Schrödinger functional as in (1) provided the gluing property holds. If we can generalise the gluing property to string theory we can repeat the diagrammatic arguments and construct the second quantised string Schrödinger functional.

2 The String Field Propagator

The string field propagator can be constructed in much the same way as in QFT [3], [5] as the transition amplitude $G(X_f; X_i)$ between arbitrary spacetime curves $X_i(\sigma)$ and $X_f(\sigma)$. We denote the propagator with boundary conditions $X^0(\sigma) = \text{constant}$, between arbitrary spacelike curves $\mathbf{X}(\sigma)$ as

$$G_{t_f - t_i}(\mathbf{X}_f; \mathbf{X}_i) = \int \mathcal{D}(X, g) e^{-\frac{1}{4\pi\alpha'} \int d^2\sigma \sqrt{g} g^{ab} \partial_a X^\mu \partial_b X_\mu} \Bigg|_{\substack{\mathbf{X}=\mathbf{X}_f(\sigma), X^0=t_f \\ \mathbf{X}=\mathbf{X}_i(\sigma), X^0=t_i}} \quad (10)$$

At tree level the worldsheet is a finite strip (cylinder) for open (closed) strings. An arbitrary metric can be written as a diff \times Weyl transformation (orthogonal to the CKV for the closed string) of a reference metric $\hat{g}_{ab}(T)$ for some value of the Teichmüller parameter T . We take $\hat{g}_{ab}(T) = \text{diag}(1, T^2)$ so that T represents the intrinsic length of the worldsheet. The propagator is [6]

$$G_t(\mathbf{X}_f; \mathbf{X}_i) = \int_0^\infty dT \text{Jac}(T) (\text{Det}' \hat{P}^\dagger \hat{P})^{\frac{1}{2}} (\text{Det} \hat{\Delta})^{-13} \int \mathcal{D}\xi e^{-S_{\text{cl}}[X_{\text{cl}}, \hat{g}(T)]}. \quad (11)$$

The measure on Teichmüller space is $\text{Jac}(T)$ given by

$$\text{Jac}(T)_{\text{open}} = \frac{(h_{ab}|\chi_{ab})}{(h_{ab}|h_{ab})^{1/2}}, \quad \text{Jac}(T)_{\text{closed}} = \frac{(h_{ab}|\chi_{ab})}{(V^a|V^a)^{1/2}(h_{ab}|h_{ab})^{1/2}}$$

where h_{ab} is the zero mode of \hat{P}^\dagger , χ_{ab} is the symmetric traceless part of $\hat{g}_{ab,T}$ and V^a is the CKV on the cylinder. X_{cl} satisfies the wave equation in metric \hat{g} with boundary conditions $X_{\text{cl}}^\xi|_{\tau=0} = X_i(\sigma)$, $X_{\text{cl}}^\xi|_{\tau=1} = X_f(\sigma)$. The remaining ξ integral is over reparametrisations of the boundary data. If we attach reparametrisation invariant functionals $\Pi_i[\mathbf{X}_i]$, $\Pi_f[\mathbf{X}_f]$ to the boundaries of the worldsheet then this integral can be done trivially to give an (infinite) constant factor, for then

$$\begin{aligned} & \int \mathcal{D}(X_f, X_i) \int \mathcal{D}\xi e^{-S_{\text{cl}}[X_{\text{cl}}, \hat{g}]} \Pi_i[X_f] \Pi_f[X_i] \\ &= \int \mathcal{D}(X_{\text{cl}}|_{\tau=1}, X_{\text{cl}}|_{\tau=0}) e^{-S_{\text{cl}}[X_{\text{cl}}, \hat{g}]} \Pi_f[X_{\text{cl}}|_{\tau=1}] \Pi_i[X_{\text{cl}}|_{\tau=0}] \int \mathcal{D}\xi. \end{aligned} \quad (12)$$

The same applies when we sew two worldsheets together, since G itself is reparametrisation invariant. Carlip's sewing method [7], required for correctly combining moduli spaces, involves integrating over all boundary values of X^0 which in our problem is not appropriate. We wish to generalise (4), the key to which is the correct identification of the degrees of freedom on the boundary. The Alvarez conditions [8] on the reparametrisations, $n^a \xi_a = n^a t^b P(\xi)_{ab} = 0$, split into orthogonal pieces on the strip (cylinder), the τ - components of which do not couple to reparametrisations of the boundary. When we sew two worldsheets only half of the determinant of $P^\dagger P$ is sewn together, the remainder cancelling the effects of not integrating over X^0 .

We can make this precise using ghosts. The rôle of the ghosts in string theory is to cancel the undesirable effects of including the X^0 oscillators. Our ghosts will do the same thing, although to a different end. Take the usual representation of the metric integral

$$(\text{Det}' P^\dagger P)^{1/2} = \int \mathcal{D}(b, c) e^{-\frac{1}{2} \int b_{ab} P(c)^{ab}} \quad (13)$$

and change variable $b = P\gamma$ for Grassmann vector γ , which turns the ghost sector of the path integral into

$$(\text{Det}' P^\dagger P)^{1/2} = \int \mathcal{D}(\gamma, c) (\text{Det}' P^\dagger P)^{-1/2} e^{-\frac{1}{2} \int P(\gamma)_{ab} P(c)^{ab}}. \quad (14)$$

Now represent the determinant on the RHS of the above by a bosonic vector integral,

$$(\text{Det}' P^\dagger P)^{-1/2} = \int \mathcal{D}f e^{-\frac{1}{2} \int P(f)_{ab} P(f)^{ab}}. \quad (15)$$

For the closed string this change of variable is defined only up to shifts $J^a \rightarrow J^a + \lambda V^a$ for $J^a \in \{c^a, \gamma^a, f^a\}$, so we choose $(J|V) = 0$ removing the c.o.m. from the classical pieces. This is our new ghost system. In accord with the Alvarez conditions we integrate out the fields J^τ and fix the values of J^σ on the boundaries so the propagator in the extended Hilbert space interpolates

between arbitrary values of $\mathbf{X}(\sigma), \{J^\sigma(\sigma)\}$ at particular times. Letting \mathbf{B} denote boundary values of \mathbf{X}^i , the set J^σ and the X^0 oscillators (zero) it can be shown [10] that the Euclidean generalisation of (4) holds in string theory as

$$\int \mathcal{D}\mathbf{B} G_{t_2-t}(\mathbf{B}_2; \mathbf{B}) \overleftrightarrow{\frac{\partial}{\partial t}} G_{t-t_1}(\mathbf{B}; \mathbf{B}_1) = G_{t_2-t_1}(\mathbf{B}_2; \mathbf{B}_1), \quad t_2 > t > t_1. \quad (16)$$

We can check our method using the cancellation of the Weyl anomaly. Even in the critical dimension (10) has a dependence on the Liouville field at the corners of the open string worldsheet [9]. The sewing prescription we have described cancels the anomaly on the boundaries being sewn, ensuring that the sewn worldsheet carries no anomaly in the bulk.

Our factorisation of the ghosts may seem ad-hoc but in fact follows from the gauge choice

$$\int d^2\sigma \sqrt{g} g^{ab} \hat{h}_{ab} \equiv (\hat{h}_{ab}|g_{ab}) = 0, \quad \hat{P}^\dagger \left(\frac{\sqrt{g} g^{rs}}{\sqrt{\hat{g}}} \right)^a = 0, \quad (17)$$

where a hat denotes use of the fiducial metric (this is equivalent to the usual choice $\sqrt{g} g^{ab} = \sqrt{\hat{g}} \hat{g}^{ab}(T)$). The corresponding gauge fixed action is

$$S_{\text{BRST}} = \frac{1}{2} \int d^2\sigma \sqrt{\hat{g}} \hat{g}^{ab} \partial_a X \partial_b X + \frac{1}{2} \int d^2\sigma \sqrt{\hat{g}} \hat{P}(\gamma)_{ab} \hat{P}(c)^{ab}. \quad (18)$$

The BRST transformations are

$$\delta_Q X = c^a \partial_a X, \quad \delta_Q c^a = c^b \partial_b c^a, \quad \delta_Q \hat{P}(\gamma)_{ab} = -2\hat{T}_{ab}, \quad (19)$$

where \hat{T}_{ab} is the usual string energy momentum tensor with $b_{ab} = \hat{P}(\gamma)_{ab}$. This set of transformations are non-local but have the natural interpretation of generating local reparametrisations of the boundary. Consider the string field propagator written as

$$G(\mathbf{X}_f; \mathbf{X}_i) = \int \mathcal{D}(X, \gamma, c) (\text{Det } \hat{P}^\dagger \hat{P})^{-1/2} e^{-S_{\text{BRST}} - S_J} \Big|_{X=X_i}^{X=X_f} \quad (20)$$

where S_{BRST} is as in (18) and S_J is a source term which generates boundary values of the ghosts,

$$S_J = \frac{1}{2} \int d^2\sigma \sqrt{\hat{g}} (\hat{P}^\dagger \hat{P} \gamma)_a c_{\text{cl}}^a + \gamma_{\text{cl}}^a (\hat{P}^\dagger \hat{P} c)_a = \int_{\text{bhd}} d\Sigma^s (\hat{P} \gamma)_{rs} c_b^s + \gamma_b^s (\hat{P} c)_{rs}. \quad (21)$$

In the above, c_{cl}^a obeys $\hat{P}^\dagger \hat{P} = 0$ and equals the boundary values of the ghosts on the Dirichlet sections of the worldsheet. The bulk action S_{BRST} is BRST invariant for arbitrary boundary values of X, c, γ . The source term S_J does not respect this symmetry, so we are led to expect a Ward identity resulting from a shift in integration variables corresponding to (19), $\langle \delta_Q S_J \rangle = 0$. In addition to

the usual short distance divergences in the quantum pieces, the corner anomaly leads to finite, non-zero contributions from the image charges. We find the Ward identity can be written as an operator acting on the propagator,

$$\left[\int_0^\pi d\sigma c_b^\sigma \mathbf{X}'_b \frac{\delta}{\delta \mathbf{X}_b} + (c_b^\sigma \gamma_b^\sigma)' \frac{\delta}{\delta \gamma_b^\sigma} + \frac{1}{2} c_b^\sigma c_b^{\sigma'} \frac{\delta}{\delta c_b^\sigma} + \frac{26}{8} (c^{\sigma'}(0) + c^{\sigma'}(\pi)) \right] G = 0, \quad (22)$$

where a subscript b indicates boundary data. This operator describes the transformation of 25 scalars \mathbf{X} and the tangential component of a vector γ^σ under a reparametrisation of the boundary generated by the ghost c^σ , with quantum corrections. Demanding BRST invariance here does not put the string field on shell, as the reparametrisations are only a subset of those described by BRST in the usual formalism.

3 T-Duality in the Schrödinger Functional

The sewing rule (16) confirms that despite the extended nature of the string a Schrödinger representation makes sense for string field theory, and we can carry over our diagrammatic arguments so that

$$\mathcal{S}[\Pi_f, t; \Pi_i, 0] = \exp \left(-\frac{1}{2} \left(\text{diagram 1} + \text{diagram 2} - \frac{1}{2} \text{diagram 3} \right) \right) \quad (23)$$

for momentum string fields $\Pi[\mathbf{X}, J^\sigma]$, and the double line represents the orbifolded propagator for either the open or closed string. The normalisation constant is discussed below. The orbifold leads naturally to the question of what rôle T-duality plays. We now show that T-duality exchanges the states attached to the propagators with backgrounds in the dual picture, and vice versa.

The closed string Schrödinger functional is T-dual to the loop diagrams appearing in the normalisation of the open string Schrödinger functional. We set $t = \pi R$, making the orbifold radius explicit, and use Poisson resummation and a change in modular parameter to convert the closed propagators into open loops so that, in an obvious notation, the Schrödinger functional becomes

$$\begin{aligned} \log \mathcal{S}_{\text{closed}} &= -\frac{1}{2} \sum_{n \text{ even}} \Pi_f G_{\pi R n} \Pi_f + \sum_{n \text{ odd}} \Pi_f G_{\pi R n} \Pi_i - \frac{1}{2} \sum_{n \text{ even}} \Pi_i G_{\pi R n} \Pi_i \\ &= -\frac{1}{2} \sum_{n \text{ even}} \Pi_f G_{\pi \tilde{R} n} \Pi_f + \sum_{n \text{ even}} \Pi_i G_{\pi \tilde{R} n} e^{in\pi/2} \Pi_f - \frac{1}{2} \sum_{n \text{ even}} \Pi_i G_{\pi \tilde{R} n} \Pi_i \end{aligned} \quad (24)$$

where $\tilde{R} = \alpha'/R$ and G in the second line is an open string contribution. The new exponent comes from the resummation and the fields glued onto the Dirichlet sections of the closed string propagator become an averaging over backgrounds characterised by Π_i , Π_f coupling to the ends of the open string. These

backgrounds, as they must be, are the same at each end of the string, for we can write the above as

$$\log \mathcal{S}_{\text{closed}} = -\frac{1}{2} \sum_{n \text{ even}} (\Pi_i - e^{iA \int dX^0} \Pi_f) G_{\pi \tilde{R} n} (\Pi_i - e^{iA \int dX^0} \Pi_f) \quad (25)$$

with Wilson line value $A = (2\tilde{R})^{-1}$. Let us give an explicit example. We take reparametrisation invariant boundary states $\Pi_{i,f}[\mathbf{X}] = \delta^p(\mathbf{X}(\sigma) - \mathbf{q}_{i,f})$ for $\mathbf{q}_{i,f}$ constant p -vectors. These are pointlike states in p directions and Neumann states in $25 - p$ directions, $0 \leq p \leq 25$. The closed string Schrödinger functional is

$$\begin{aligned} \log \mathcal{S}_{\text{closed}} = & -\text{Vol}^{25-p} \int_0^\infty \frac{dT}{T^{\frac{p+1}{2}}} e^{2T} \prod_{m=1} (1 - e^{-2mT})^{-24} \sum_{n \text{ even}} e^{-\frac{\pi^2 \tilde{R}^2}{2\alpha' T} n^2} \\ & + \text{Vol}^{25-p} \int_0^\infty \frac{dT}{T^{\frac{p+1}{2}}} e^{-\frac{\delta \mathbf{q}^2}{2\alpha' T} + 2T} \prod_{m=1} (1 - e^{-2mT})^{-24} \sum_{n \text{ odd}} e^{-\frac{\pi^2 \tilde{R}^2}{2\alpha' T} n^2} \end{aligned} \quad (26)$$

with $\delta \mathbf{q} = \mathbf{q}_f - \mathbf{q}_i$. Since the open string runs from $\sigma = 0 \dots \pi$ and the closed string from $\sigma = 0 \dots 2\pi$ we must scale the closed string worldsheet to interpret (26) as an open loop. We include this in a change of modular parameter $S := 2\pi^2/T$. After this and a Poisson resummation we find

$$\begin{aligned} \log \mathcal{S}_{\text{closed}} = & -\text{Vol}^{25-p} \int_0^\infty \frac{dS}{S} \frac{1}{S^{\frac{26-p}{2}}} e^S \prod_{m=1} (1 - e^{-mS})^{-24} \\ & \times \sum_{n \text{ even}} e^{-\frac{\pi^2 \tilde{R}^2}{4\alpha' S} n^2} \left(1 - e^{\frac{in\pi}{2}} e^{-\frac{\delta \mathbf{q}^2 S}{4\pi\alpha'}} \right). \end{aligned} \quad (27)$$

Now consider an open string loop. The measure on Teichmüller space is dS/S (this gives the logarithm of the worldsheet propagator). If the string has Neumann conditions in $26 - p$ directions (including X^0) and Dirichlet conditions in p directions, as for a string on a $D(25 - p)$ -brane then the trace over \mathbf{X} gives the eta function and the factor $(S^{-1/2} \text{Vol})^{(25-p)}$ from the $25 - p$ zero modes. The sum and remaining factor of $S^{-1/2}$ come the trace over X^0 in the co-ordinate representation. We arrive at (27), if the term in large brackets represents an averaging over backgrounds of Wilson lines and $D(25 - p)$ -branes of separation $\delta \mathbf{q}$.

We interpret the open string duality as taking us from one Schrödinger functional to another with an exchange of boundary states and backgrounds. Explicit examples are difficult to construct since the corner anomaly, not an issue for the closed string, forces us to find conformally invariant states and

backgrounds, but we can give an outline. Poisson resummation implies

$$\left(\frac{\pi R^2}{\alpha' T}\right)^{1/2} \sum_{n \text{ even}} e^{-\frac{\pi^2 R^2}{4\alpha' T} n^2} = \sum_{n \text{ even}} e^{-\frac{\bar{R}^2 T}{4\alpha'} n^2} + \sum_{n \text{ odd}} e^{-\frac{\bar{R}^2 T}{4\alpha'} n^2} \quad (28)$$

(for the odd sum the plus on the R.H.S. becomes minus) where $\bar{R} = 2\alpha'/R$. Following a modular transformation $S := \pi^2/T$ these are the sums in the open string Schrödinger functional. Again the states now represent an averaging over backgrounds. The open string Schrödinger functional becomes

$$\log \mathcal{S}_{\text{open}} = -\frac{1}{2} \sum_{n \text{ even}} (\Pi_i - \Pi_f) G_{n\pi\bar{R}} (\Pi_i - \Pi_f) - \frac{1}{2} \sum_{n \text{ odd}} (\Pi_i + \Pi_f) G_{n\pi\bar{R}} (\Pi_i + \Pi_f) \quad (29)$$

and the new momentum states are characterised by the original Neumann condition on the open string ends. To interpret this as strings moving in a single background we can introduce a Wilson line, $\Pi_i - e^{iA \int dX^0} \Pi_f$, with value $A = (\bar{R})^{-1}$.

In summary we have shown that the Schrödinger functional describing evolution through time t of second quantised strings can be written in terms of first quantised strings moving on the orbifold $\mathbb{S}^1/\mathbb{Z}_2$ and that the consequent T-duality interchanges t with $1/t$ and momentum fields with Dp -brane backgrounds. BRST transformations describe reparametrisations of boundary data and for the open string are sensitive to the Weyl anomaly even in the critical dimension. We have worked only in the free theory but interactions will be discussed in [10].

References

- [1] E. Witten, “Noncommutative Geometry And String Field Theory,” Nucl. Phys. B **268** (1986) 253
- [2] K. Symanzik, “Schrodinger Representation And Casimir Effect In Renormalizable Quantum Field Theory,” Nucl. Phys. B **190**, 1 (1981)
- [3] L. Brink, P. Di Vecchia and P. S. Howe, “A Locally Supersymmetric And Reparametrization Invariant Action For The Spinning String,” Phys. Lett. B **65**, 471 (1976)
- [4] W. Taylor and B. Zwiebach, “D-branes, tachyons, and string field theory,” arXiv:hep-th/0311017.
- [5] A.M. Polyakov, “Gauge fields and strings”, Harwood Academic Publishers, 1987, ISBN 3-7186-0492-6
- [6] G. W. Moore and P. Nelson, “Absence Of Nonlocal Anomalies In The Polyakov String,” Nucl. Phys. B **266**, 58 (1986)

- [7] S. Carlip, “Bordered surfaces, off-shell amplitudes, sewing and string field theory”, published in “Functional integration, geometry and strings”, Birkhäuser Verlag Basel, 1989, ISBN 3-7643-2387-6
- [8] O. Alvarez, “Theory Of Strings With Boundaries: Fluctuations, Topology, And Quantum Geometry”, Nucl. Phys. B **216**, 125 (1983)
- [9] P. Mansfield, “Bosonic Ghosts And Witten’s Noncommutative Geometry,” Nucl. Phys. B **306**, 630 (1988)
- [10] A. . Ilderton and P. Mansfield, “Timelike T-duality in the string field Schroedinger functional,” arXiv:hep-th/0411166