

## Durham Research Online

---

### Deposited in DRO:

23 February 2016

### Version of attached file:

Accepted Version

### Peer-review status of attached file:

Peer-reviewed

### Citation for published item:

Reissland, N. and Francis, B. and Buttanshaw, L. and Austen, J. M. and Reid, V. (2016) 'Do fetuses move their lips to the sound that they hear? An observational feasibility study on auditory stimulation in the womb.', *Pilot and feasibility studies.*, 2 (1). p. 14.

### Further information on publisher's website:

<https://doi.org/10.1186/s40814-016-0053-3>

### Publisher's copyright statement:

© 2016 Reissland et al. Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

### Additional information:

### Use policy

---

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

# Pilot and Feasibility Studies

## Do fetuses move their lips to the sound that they hear? An observational feasibility study on auditory stimulation in the womb

--Manuscript Draft--

<b>Manuscript Number:</b>	PAFS-D-15-00008R2	
<b>Full Title:</b>	Do fetuses move their lips to the sound that they hear? An observational feasibility study on auditory stimulation in the womb	
<b>Article Type:</b>	Research	
<b>Funding Information:</b>	Economic and Social Research Council (GB) (ES/L003155/1)	Dr Vincent Reid
<b>Abstract:</b>	<p><b>Background:</b> We investigate in this feasibility study whether specific lip movements increase prenatally when hearing a particular sound. We hypothesised that fetuses would produce more mouth movements resembling those required to make the sound stimulus they heard (i.e. mouth stretch) compared with a no-sound control group who heard no specific auditory stimuli. Secondly, we predicted that fetuses hearing the sound would produce a similar number of mouth movements unrelated to the sound heard (ie lip pucker) as the no-sound group of fetuses.</p> <p><b>Methods:</b> In an observational feasibility study, 17 fetuses were scanned twice at 32 and 36 weeks gestation, and two different types of mouth movements recorded. Three fetuses received an auditory stimulus, and 14 did not. A generalised mixed effects log-linear model was used to determine statistical significance.</p> <p><b>Results:</b> Fetuses in the sound group performed one specific mouth movement (mouth stretch) significantly more frequently than fetuses in the no-sound group. A significant interaction between group and gestational age indicates that there was differential change in this specific movement as age increases (<math>X^2 = 7.58</math> on 1 df, <math>p = .006</math>), with the no sound group showing a decline of 76% between 32 weeks and 36 weeks (<math>p &lt; .001</math>), whereas the sound group showed no significant change over time (<math>p = 0.41</math>). There was no significant difference between the sound group and no sound group in the frequency of lip puckering - the second, unrelated mouth movement (<math>p = 0.35</math>).</p> <p><b>Conclusions:</b> These results suggest that a sound stimulus is associated with an increase in specific, rather than general, mouth movements. The results are informative for the development of infant speech, and potentially could also lead to a diagnostic test for deafness in utero. More research is needed to replicate this research with a randomised design and with a range of different auditory stimuli which would be produced with different mouth movements, such as "o" which would be seen as pursed lips.</p>	
<b>Corresponding Author:</b>	Nadja Reissland UNIVERSITY OF DURHAM UNITED KINGDOM	
<b>Corresponding Author Secondary Information:</b>		
<b>Corresponding Author's Institution:</b>	UNIVERSITY OF DURHAM	
<b>Corresponding Author's Secondary Institution:</b>		
<b>First Author:</b>	Nadja Reissland	
<b>First Author Secondary Information:</b>		
<b>Order of Authors:</b>	Nadja Reissland	
	Brian Francis	
	Louisa Buttanshaw	

	Joe Austen
	Vincent Reid

[Click here to view linked References](#)

**Do fetuses move their lips to the sound that they hear? An observational feasibility study on  
auditory stimulation in the womb.**

Nadja Reissland<sup>1\*</sup>, Brian Francis<sup>2</sup>, Louisa Buttanshaw<sup>1</sup>, Joe M. Austen<sup>1</sup>, Vincent Reid<sup>3</sup>

<sup>1</sup>Department of Psychology, Durham University, Durham, United Kingdom

<sup>2</sup>Department of Maths and Stats, Lancaster University, Lancaster, United Kingdom

<sup>3</sup>Department of Psychology, Lancaster University, Lancaster, United Kingdom

\*Corresponding Author Nadja Reissland: [n.n.reissland@durham.ac.uk](mailto:n.n.reissland@durham.ac.uk)

Brian Francis: [b.francis@lancaster.ac.uk](mailto:b.francis@lancaster.ac.uk)

Louisa Buttanshaw: [louisabuttanshaw@gmail.com](mailto:louisabuttanshaw@gmail.com)

Joe M. Austen: [j.m.austen@durham.ac.uk](mailto:j.m.austen@durham.ac.uk)

Vincent Reid: [v.reid@lancaster.ac.uk](mailto:v.reid@lancaster.ac.uk)

**Running title:** Auditory stimulation in the womb

# Abstract

**Background:** We investigate in this feasibility study whether specific lip movements increase prenatally when hearing a particular sound. We hypothesised that fetuses would produce more mouth movements resembling those required to make the sound stimulus they heard (i.e. mouth stretch) compared with a no-sound control group who heard no specific auditory stimuli. Secondly, we predicted that fetuses hearing the sound would produce a similar number of mouth movements unrelated to the sound heard (ie lip pucker) as the no-sound group of fetuses.

**Methods:** In an observational feasibility study, 17 fetuses were scanned twice at 32 and 36 weeks gestation, and two different types of mouth movements recorded. Three fetuses received an auditory stimulus, and 14 did not. A generalised mixed effects log-linear model was used to determine statistical significance.

**Results:** Fetuses in the sound group performed one specific mouth movement (mouth stretch) significantly more frequently than fetuses in the no-sound group. A significant interaction between group and gestational age indicates that there was differential change in this specific movement as age increases ( $\chi^2 = 7.58$  on 1 df,  $p = .006$ ), with the no sound group showing a decline of 76% between 32 weeks and 36 weeks ( $p < .001$ ), whereas the sound group showed no significant change over time ( $p = 0.41$ ). There was no significant difference between the sound group and no sound group in the frequency of lip puckering - the second, unrelated mouth movement ( $p = 0.35$ ).

**Conclusions:** These results suggest that a sound stimulus is associated with an increase in specific, rather than general, mouth movements. The results are informative for the development of infant speech, and potentially could also lead to a diagnostic test for deafness in utero. More research is needed to replicate this research with a randomised design and with a range of different auditory stimuli which would be produced with different mouth movements, such as "o" which would be seen as pursed lips.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Keywords:** prenatal stimulation; fetal hearing; fetal mouth movement

# Background

From birth, infants produce silent movements resembling the lip movements necessary for speech.

A seminal study reported by Trevarthen [1] in which one 7 week old girl silently produced lip and mouth movements which resembled the mouth movements of a female speaker reading a word list, has been interpreted as the basis of the intention to speak. These mouth movements indicated that very young infants when hearing language in their first weeks after birth produce mouth and lip movements similar to those necessary to replicate speech. In addition, infants just after birth have been shown to imitate silently mouth movements required to produce language sounds, even when no sound was produced [2].

One way in which this ability could develop prenatally was proposed by Green and Wilson [3]. They argued that randomly produced lip and jaw movements during fetal development could conceivably create sensorimotor pathways that could serve as precursors of early speech. They based their conjecture on the well-established suggestion that neuronal firing resulting in leg and hand movements are precursors of walking and grasping movements (e.g. [4;5]). Neuronal activation elicits early limb movements and these limb movements in turn help to consolidate the pathways which shape purposeful movements such as reaching or walking (e.g. [4;5]). Hence, early activation of specific mouth and jaw movements such as a jaw drop with elongation of lips in the vertical axis could be a precursor necessary for speech sounds such as “a” and pursing lips could be a precursor to producing the lip movement necessary for the sound “o”.

Prenatal cognitive development has been tested in relation to sound and light stimulation. For example, Horimoto et al. [6] reported that fetuses between 32 and 34 weeks gestation showed a high incidence of mouth movements that were later during gestation correlated with non-rapid eye movements. However, the mouth movements which have been reported in previous studies are general movements that do not have specific definitions, such as a smile or grimace.

1 The following sequential order of development has been found for specific fetal oral movements,  
2 namely jaw opening, jaw closing, tongue movement, and lip movement [7], whereby spontaneous  
3  
4 movements of the jaw appear as early as 11 weeks during prenatal development [8]. Regarding first  
5  
6 vocalizations in infancy, jaw opening and closing are primary movements during babbling [9] which  
7  
8 can be distinguished from other mouth openings such as smiling [10]. Green, Moore and Reilly [11]  
9  
10 investigated the sequential development of jaw and lip control and found in an analysis of children  
11  
12 aged 1 to 6 years of age developed sequentially, with jaw movements preceding lip movements.  
13  
14  
15  
16  
17

18 By the second trimester, the fetal auditory cortex is reorganized by external stimulation (e.g.  
19  
20 [12;13]). In the study by Kisilevsky et al., [13] which examined maturation of fetal responding to  
21  
22 airborne auditory stimuli, differential responding occurred as a result of fetal maturation during the  
23  
24 third trimester. Hence, we selected two gestational ages in the third trimester to examine whether  
25  
26 we might find maturational changes in reaction to an auditory stimulus. If fetuses were exposed to  
27  
28 auditory rhythmic stimulation [14], then this might be reflected in their production of movement  
29  
30 patterns, specifically movement patterns of the jaw and lips. Given that fetal mouth movements  
31  
32 develop but have to date not been analysed to the level of specificity required, it is essential to  
33  
34 establish whether there is a relationship between mouth movements and sound stimulation. If jaw  
35  
36 and lip movements are produced prenatally in response to sound before the ability to produce  
37  
38 speech develops, then this would support the argument that precursors of language are rooted in  
39  
40 fetal development. In particular, we argue that if types of jaw and lip movements vary between  
41  
42 fetuses who do not hear any sounds during scans and fetuses who hear specific sounds, there might  
43  
44 be reason to believe that lip and jaw movements are pre-cursors of silent pre-speech movements  
45  
46 which can be observed in neonates.  
47  
48  
49  
50  
51  
52  
53

54  
55  
56 The production of the auditory stimulus presented in this study involved predominantly jaw  
57  
58 movements, allowing us to determine whether the response to this stimulus was specific (i.e., only  
59  
60  
61  
62  
63  
64  
65



1 jaw movements were produced) or general (i.e., jaw and lip movements were produced). In order to  
2 investigate this question we studied two groups of fetuses: one that was presented with the  
3 auditory stimulus during scanning and a second no-sound control group of fetuses that was not  
4 presented with any sound stimulus. We expected that those fetuses presented with the specific  
5 sound would produce mouth movements consistent with the sound when contrasted with another  
6 mouth movement that typically manifests with the same frequency in normally developing fetuses.  
7 This feasibility study was designed to establish whether it is possible to test fetal fine grained  
8 reactions to specific sound stimulation.  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20

## 21 **Methods**

### 22 **Ethics**

23  
24  
25  
26  
27  
28 Ethical permission for the feasibility study was obtained from the County Durham and Tees Valley 2  
29 Research Ethics Committee (REC Ref: 08/H0908/31 and County Durham and Tees Valley 2 Research  
30 Ethics Committee REC Ref: 11/NE/03/61) and the research and development department of James  
31 Cook University Hospital, as well as the Durham University (Department of Psychology ethics  
32 committee). All mothers gave informed written consent.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

### 43 **Stimulus**

44  
45  
46 The auditory stimulus consisted of multiple presentations of the sound MA (/ma:/ in the  
47 International Phonetic Alphabet). The MA sound was spoken in a female voice, was 0.40s in length,  
48 and was repeated 8 times with 0.80s of silence between each presentation. This cluster of eight MA  
49 sounds was then repeated for the duration of the scan, with 6.0s of silence between each cluster.  
50  
51  
52  
53

54  
55 The auditory stimulus was played on a Sandisk Sansa Clip portable MP3 player, attached to a  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

[15], who report research showing that the speaker should not be placed on the mother's abdomen, for this study the speaker was held at a distance of 3cm above the mother's abdomen near the ear of the fetus for the duration of the stimulus presentation. The sound pressure level at a distance of 3cm from the speaker was 94dB, measured with a Precision Gold N05CC Digital Sound Meter (with a measurement range of 30 - 100dB with an accuracy of  $\pm 1.5$ dB), although uterine attenuation will have reduced the sound level for the fetus by approximately 20 - 35dB [16;17]. The auditory stimulus contained frequencies between 0Hz and 11kHz, with most output in the 0.6 - 1.6kHz and 2.6 - 3.6kHz regions. These frequencies are audible to fetuses from 29 weeks gestational age [16]. Although, in a study of newborn auditory matching, Chen, Striano and Rakoczy [18] only measured responding to an auditory stimulus during its presentation, we measured responding during the presentation of sound and during the intervening seconds of silence. The inclusion of short periods of silence following the presentation of sound was deemed appropriate as fetuses may be slower to respond to sound stimuli than newborns due to being less developmentally mature. The short periods of silence allow the fetuses time to respond to the recently presented auditory stimuli. The relative frequencies of jaw and lip movements during the presentation of the auditory stimuli were analysed and compared with the no-sound control group who did not experience the auditory stimulus.

## Participants

Mothers who had completed their normal 20-week anomaly scans were invited to participate in this study. All fetuses participating in this study were completely healthy as determined by their 20 week scan. A convenience sample of seventeen mothers was recruited for this feasibility study; three mothers whose fetuses were exposed to the auditory stimulus (1 boy and 2 girls), the sound group and 14 mothers whose fetuses were not exposed to any stimulus (7 boys and 7 girls), the no-sound group. The no-sound group participants were recruited through the midwives of the antenatal unit of the James Cook University Hospital, Middlesbrough, UK, and sound group participants through the Hypnobirthing group in London at the Harley Street Ultrasound Centre

1 (www.thewisehippo.com) following approved ethical procedures. During consent and before each  
2 procedure, mothers were made aware that the scans were for research purposes and were not  
3  
4 routine medical scans. Given that maternal stress, attachment, anxiety, and depression are known to  
5  
6 affect fetal behavior (e.g., [19, 20, 21, 22]), the two groups were assessed for these factors with the  
7  
8 Perceived Stress Scale (PSS [23]), Antenatal Maternal Attachment Scale (AMAS [24]), and the  
9  
10 Hospital Anxiety and Depression Scale (HADS [25]).  
11  
12  
13  
14  
15  
16  
17

## 18 Procedure

19  
20  
21 All participating mothers received scans at 32 and 36 weeks gestational age, with fetuses being  
22  
23 scanned for approximately 900-1200 seconds, with the maximum time of the scan determined by  
24  
25 the British Medical Ultrasound Society (BMUS) guidelines. The scan times of 32 and 36 weeks were  
26  
27 chosen as Kisilevsky et al [13] identified that it was more likely to get reactions to sound in the latest  
28  
29 stages of pregnancy. The scans took place either in the radiography department of James Cook  
30  
31 University Hospital, where mothers had previously undergone their routine 12 and 20 week medical  
32  
33 scans, or in the London Ultrasound centre. The scanning took place with mothers lying in a darkened  
34  
35 room on their back or on their side, depending on the position of the fetus and how comfortable  
36  
37 mothers were. The fetal face and upper torso were visualized both by means of 4-D color full frontal  
38  
39 or facial profile ultrasound recordings, as well as sequences of traditional 2-D monochrome images.  
40  
41 The scans were recorded for off line analysis with a GE Voluson E8 Expert Ultrasound System using a  
42  
43 GE RAB4–8L Macro 4D Convex Array Transducer. Mothers were provided with a DVD copy of their  
44  
45 scans.  
46  
47  
48  
49  
50  
51  
52  
53

## 54 Measures

55  
56  
57 Scan recordings were used to code mouth movements using the Fetal Observable Movement System  
58  
59 (FOMS) [26], an adaptation of the Facial Action Coding System [27], which has been found to be  
60  
61  
62  
63  
64  
65

1 reliable in previous research [28;29]. Following established procedures [29] two types of mouth  
2 movements were identified for analysis: *mouth stretch* and *lip pucker*.  
3

4  
5  
6  
7 Mouth stretch is defined by the lower jaw being pulled down by the action of the external pterygoids  
8 and digastricus muscles, so that the mouth is actively opened. The opening is stretched such that  
9 the longest axis is the vertical plane. The cheeks are stretched and flattened and the skin on the chin  
10 also may become bulged.  
11  
12  
13  
14  
15

16  
17  
18 Lip pucker in contrast, is defined by the lips narrowing and pursing with the lips protruding forwards.  
19 This is caused by the incisivii labii superioris and incisivii labii inferioris muscles, which pull the  
20 corners of the lips medially. The lips usually appear as if contracted and the mouth opening will look  
21 smaller and rounded. There also may be some bulging of the chin as the skin of the chin is pulled  
22 upwards towards the lips. In normally developing fetuses, these two mouth movements are  
23 produced with relatively equal frequencies at the gestational ages scanned in this study [26].  
24  
25  
26  
27  
28  
29  
30  
31  
32

33  
34  
35 The aim in this feasibility study was to analyse up to 600 seconds of codable scan for each fetal scan.  
36 Codable sections of the scan for the control scans were sections where the fetal face was visible, and  
37 where the pocket of amniotic fluid was present to allow a clear image. For the sound group, coding  
38 occurred during the presentation of sound and during the intervening seconds of silence. As fetal  
39 movements differ as a function of the movement state of the fetus [30], it is essential that all of the  
40 fetuses were in an active state during scanning. Assessment of the movement behaviour showed  
41 that all fetuses were in the active states of 2F or 3F during scanning, as assessed by their gross body  
42 movements and eye movements according to Nijhuis et al.'s four-state categorisation [30], and not in  
43 states 1F (totally passive) or 4F (overly active).  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58

## 59 Statistical analysis

60  
61  
62  
63  
64  
65

1 Reflecting the longitudinal structure of the measurements and the non-normality of the count  
2 responses, a Poisson log-linear mixed effects analysis [31] was used to assess the effect of  
3  
4 experimental group and gestational age, and the interaction between them using the *glmer* function  
5  
6 of the *lme4* library in R [32]. A mixed effects analysis has been shown to be superior to standard  
7  
8 repeated measures analysis of variance for experimental data [33]. Moreover, imbalance in the  
9  
10 number of participants in each treatment arm, and in the number of scans contributed by each  
11  
12 mother can easily be accounted for. The analysis models the number of *mouth movements* of  
13  
14 different types as a count variable adjusted by the length of analysed scan as an exposure variable,  
15  
16 and a random individual fetus effect. The individual random fetus effect allows for individual  
17  
18 variability between fetuses in their overall propensity to mouth movements and is assumed to be  
19  
20 normally distributed.  
21  
22  
23  
24  
25  
26  
27

28 Formally, we can write the model as  
29

$$30 \quad M_{it} \sim \text{Poisson}(\lambda_{it})$$

31 with  
32

$$33 \quad \log(\lambda_{it}) = \log(\text{scan length}_{it}) + \beta_0 + \beta_1 \text{ gestational age}_{it} + \beta_2 \text{ treatment}_i$$

$$34 \quad + \beta_3 (\text{gestational age} * \text{treatment})_{it} + u_i$$

$$35 \quad u_i \sim \text{Normal}(0, \sigma_f^2),$$

36 where  $M_{it}$  are the mouth movement counts for fetus  $i$  at gestational age  $t$ ,  $\lambda_{it}$  is the underlying  
37  
38 Poisson rate,  $\beta_0$  to  $\beta_3$  are unknown regression coefficients and  $\sigma_f^2$  is the individual within-fetus  
39  
40 variance. The indices to the individual covariates show which of them vary over time in our model.  
41  
42 Thus gestational age, is recorded at each scan, whereas the treatment condition (sound/no sound) is  
43  
44 constant for each fetus.  
45  
46  
47  
48  
49  
50  
51  
52

53  
54  
55  
56  
57 A test for overdispersion for count data was carried out on the full interaction model using the  
58  
59 methodology of Mancuso[34]. If the overdispersion test indicated no overdispersion, then  
60  
61  
62  
63  
64  
65

1 significance of terms was assessed through analysis of deviance likelihood ratio test, examining  
2 changes of deviance between fitted models, and comparing to a chi-squared distribution with the  
3 appropriate number of degrees of freedom. Bolker et al [35] provides full details of fitting and  
4 testing Poisson log-linear mixed effects for count data.  
5  
6  
7  
8  
9

## 10 11 **Results**

12  
13 Comparing mothers in the two groups we analysed their results on a number of scales. On the  
14 Perceived Stress Scale (PSS [23]), mothers in the sound group were similar to those in the no-sound  
15 group at 32 weeks ( $t(15) = 0.16, p = .87$ ) and at 36 weeks ( $t(15) = 0.18, p = .86$ ). Scores on the  
16 Antenatal Maternal Attachment Scale (AMAS [24]) also did not differ between mothers in the sound  
17 and no-sound control groups either at 32 weeks ( $t(15) = 0.85, p = .41$ ) or 36 weeks ( $t(15) = 0.50, p =$   
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

1 movements per hundred seconds of codable scan time. Thus, frequency of movements can be  
2 readily compared between groups with differing scan times in the descriptive Tables 1 and 2.. Below  
3  
4 we report the results for “mouth stretch” and “lip pucker”.  
5  
6

## 7 Mouth Stretches

8  
9  
10 The mouth stretch means from the fetuses at 32 and 36 weeks gestational age can be seen in Table  
11 1 and Fig. 1a. These suggest that fetuses at both gestational ages produce more frequent mouth  
12 stretches in the sound group compared with the no-sound control group. These observations were  
13 confirmed by the main effects Poisson linear mixed model. The overdispersion test gave a  
14 dispersion parameter estimate of 0.89, which is under one, and therefore indicates no  
15 overdispersion. There was a significant main effect of group,  $X^2= 5.78$  on 1 df ,  $p = .01$ , and of  
16 gestational age ( $X^2= 10.78$  on 1 df ,  $p = .001$ ), demonstrating that fetuses in the sound group  
17 performed mouth stretches more frequently than fetuses in the no-sound control group, and there  
18 was a general trend towards fewer mouth movements as gestational age increased. When an  
19 interaction model was fitted, the interaction between group and gestation was also significant ( $X^2=$   
20  $7.58$  on 1 df ,  $p = .006$ ), indicating that there was differential change in mouth stretch rate as  
21 gestational age increases. For the sound group there was no evidence of a change in the rate of  
22 mouth stretch ( $\beta = 0.49$ , 95%  $CI = [-0.71, 1.69]$ ,  $\exp(\beta) = 1.63$  ,  $p = .41$ ); for the no-sound group  
23 there was a decrease of 76% in the rate of mouth stretch ( $\beta = -1.41$ , 95%  $CI = [-2.17, -0.65]$ ,  
24  $\exp(\beta) = 0.24$ ,  $p < .001$ ).  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

48 **Table 1 about here**

49  
50  
51  
52  
53 **Figure 1 about here**

54  
55  
56  
57  
58 These data indicate that fetuses presented with the auditory stimulus MA produce more mouth  
59 stretches than fetuses presented with any specific auditory stimulus. However, it is possible that  
60  
61  
62  
63  
64  
65

1 the presentation of the auditory stimulus triggered an increase in the number of mouth movements  
2 in general. In order to test this hypothesis, the frequency with which fetuses performed the second  
3 mouth movement, a lip pucker, was established. The lip pucker is a valid mouth movement for  
4 comparison as fetuses at 32 and 36 weeks of age not exposed to any specific stimulation show  
5 similar frequencies of mouth stretches and lip puckers. To corroborate this, we examined  
6 frequencies of lip pucker and mouth stretch in the control group. A paired-samples t-test indicated  
7 that, pooled over the two gestational ages, there was no significant difference between the  
8 frequency of mouth stretches ( $M = .163$  per minute) and lip puckers ( $M = .218$  per minute),  $t(13) =$   
9  $0.63, p = .54$ .

## 23 Lip Pucker

24 The lip pucker mean counts from the fetuses at 32 and 36 weeks gestational age can be seen in  
25 Table 2 and Fig. 1b showing only small differences in the frequencies with which fetuses pucker their  
26 lips depending on gestational age and group. Across both gestational ages, it appears that a similar  
27 amount of lip puckers were produced by the control group and the sound group. This was tested  
28 using the mixed effects Poisson model as before. Again, the overdispersion test indicated that no  
29 overdispersion was present in the data, with the dispersion parameter of 0.516 being less than one.  
30 Results indicate no statistically significant main effect of group,  $X^2 = 0.87$  on 1 df,  $p = .35$ , no  
31 significant main effect of gestational age,  $X^2 = 0.01$  on 1 df,  $p = .93$ , and no interaction between  
32 group and gestation,  $X^2 = 1.64$  on 1 df,  $p = .20$ .

33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51 **Table 2 about here**  
52  
53

54 These data suggest that the frequency with which fetuses show lip pucker lip movements are  
55 independent of the presentation of an auditory stimulus. Additionally, there is no evidence that  
56 gestational age affects the rate of lip puckers. From this result we can infer that the presentation of  
57  
58  
59  
60  
61  
62  
63  
64  
65



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

the auditory stimulus MA does not cause an increase in the frequency of mouth movements in general. Rather, it increases the frequency of a specific mouth movement corresponding to the MA sound, with a jaw drop and the mouth stretching in the vertical plane.

## Discussion

Results of this study indicate that fetuses respond to a specific sound MA with a specific mouth movement which mimics the sound heard, namely a mouth stretch which involves a jaw drop. There are a number of researchers who argue that given the precocity of the functional development of the auditory system, the abilities shown in new-born babies must have their origin in prenatal life [36;13]. Given that research [37] has established that the fundamental frequency (F0) of vowels are well transmitted to the fetus, using the sound MA seemed to be an ideal candidate in this study.

However, in order to eliminate the possibility that fetuses would respond with more mouth movements in general we analysed the occurrence of another type of mouth movement, namely pursing of lips, which in the control no-sound group occurred with similar frequencies to mouth stretching. We found that upon hearing MA, fetuses did not respond with an increase in pursing of their lips. Given the results by Green et al. [11] who found that control over vertical movements of lips and jaw during speech developed sequentially with jaw movements preceding lip movements, this might explain why we found a relationship between hearing the MA sound and producing the mouth stretch. Ferronato, Domellöf and Rönqvist [38] suggest that postnatally speech stimuli are special stimuli which elicit specific behavioural reactions. They argue that the pairing of certain acoustic stimuli with defined motor activities (e.g., rhythmic sounds with rhythmic movements) demonstrate that the “human brain is primed with the body” (p. 3). This according to Ferronato et al. [38] could indicate that auditory input and behavioural output might play a role in the integration between external and internal information which is essential for learning in general and language acquisition in particular. In terms of differential responding to sounds over the duration of the third

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

trimester, fetuses did not show a significant increase in the production of mouth movements as a consequence of maturation. However, there is a numerical increase in mean responses as gestational age increases, which needs to be further investigated.

Turning to the observed changes over time, we identified a significant interaction between the sound and no-sound groups in the slope of the rate of mouth stretch in response to the stimulus of “MA”. The no sound group showed declining rate of mouth stretch, whereas the sound group showed no such decline. Fetal movements in general tend to decrease with gestational age. Our results provide evidence that appropriate mouth movements associated with specific stimuli do not decrease in this way.

This feasibility study was not a randomised control trial but used a convenience sample. However we consider it unlikely that fetuses of mothers who did not consent to participate were different to those who participated in our study, and so there was minimal selection bias.

## Conclusion

This feasibility study established that it is possible to test specific fetal fine grained reactions to sound stimulation. More research is needed to develop this feasibility study. Firstly a randomised controlled trial is needed with balanced numbers of treatment and control participants. Additionally, a range of different auditory stimuli should be examined to determine whether the fetus is reacting to a specific MA sound, or to any general auditory stimulus such as white noise. Additionally, a greater range of specific sounds could also be examined in order to fully explore how the fetus produces not only random mouth movements but specific pre-speech movements.

The potential implications of this work are twofold. Firstly, this work is likely to provide knowledge on the relationship of prenatal reaction to specific sounds and postnatal language development.

Secondly, there is a possibility that lack of reaction of the fetus to specific sounds could be used as a pre-natal diagnostic test for deafness.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Authors' contributions

NR and VR conceived the study. NR and BF wrote the paper. The coding of the fetal scans was carried out by LB and JA. BF also carried out the statistical analysis . All authors read and approved the final manuscript.

## Acknowledgements

We thank all parents who took part in this study, the Wise Hippo hypnobirthing group for advertising our study, and the London Ultrasound Centre for their support. We also thank James Cook University Hospital, in particular the scanning unit and consultants for their support This work was supported by the UK Economic and Social Research Council (grant number ES/L003155/1).

## Declarations

'The authors declare that they have no competing interests'

## References

- 1  
2  
3 [1] Trevarthen C. Communication and cooperation in early infancy: A description of primary  
4 intersubjectivity. In: Bullowa M, editor. Before speech: The beginning of interpersonal  
5 communication, London: CUP; 1979, p. 321-47.  
6  
7  
8  
9  
10  
11 [2] Reissland NN. Neonatal imitation in the first hour of life: Observations in rural Nepal. *Dev Psychol*  
12 1988;24:464–9.  
13  
14  
15  
16 [3] Green JR, Wilson EM. Spontaneous Facial Motility in Infancy: A 3D Kinematic Analysis. *Dev*  
17 *Psychobiol* 2006;48:16–28.  
18  
19  
20  
21  
22 [4] Bhat AN, Heathcock J, Galloway JC. Toy-oriented changes in hand and joint kinematics during the  
23 emergence of purposeful reaching. *Infant Behav Dev* 2005;28:445–65.  
24  
25  
26  
27  
28 [5] Thelen E. Rhythmical stereotypes in normal human infants. *Anim Behav* 1979;27:699–715.  
29  
30  
31 [6] Horimoto N, Koyanagi T, Nagata S, Nakahara H, Nakano H. Concurrence of mouthing movement  
32 and rapid eye movement/non-rapid eye movement phases with advance in gestation of the human  
33 fetus. *Am J Obstet Gynecol* 1989;161:344–51.  
34  
35  
36  
37  
38  
39 [7] Humphrey T. The development of human fetal activity and its relation to postnatal behavior. *Adv*  
40 *Child Dev Behav* 1970;5:1–57.  
41  
42  
43  
44 [8] de Vries JIP, Visser GHA, Prechtl HFR. The emergence of fetal behaviour. II. Quantitative aspects.  
45 *Early Hum Dev* 1985;12:99–120.  
46  
47  
48  
49  
50 [9] Green JR, Moore CA, Reilly KJ, Higashikawa M, Steeve RW. The physiologic development of  
51 speech motor control: Lip and jaw coordination. *J Speech Lang Hear Res* 2000;43:239–55.  
52  
53  
54  
55 [10] Holowka S, Pettito LA. Left Hemisphere Cerebral Specialization for Babies while Babbling.  
56 *Science* 2002;297:1515.  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 [11] Green JR, Moore CA, Reilly KJ. The sequential development of jaw and lip control for speech. J  
2 Speech Lang Hear Res 2002;45:66–79.  
3

4  
5 [12] Hepper PG, Scott DS, Shahidullah S. Newborn and fetal response to maternal voice. J Reprod  
6 Infant Psychol 1993;11:147–54.  
7

8  
9  
10 [13] Kisilevsky BS, Pan L, Hains SMJ. Maturation of human fetal responses to airborne sound in low-  
11 and high-risk fetuses. Early Hum Dev 2000;58:179–95.  
12

13  
14 [14] Provasi J, Anderson DI, Barbu-Roth M. Rhythm perception, production, and synchronization  
15 during the perinatal period. Front Psychol 2014;5:1048.  
16

17 [15] Kruger C, Horesh E, Crosland BA. Safe sound exposure in the fetus and preterm infant. J Obstet  
18 Gynecol Neonatal Nurs 2012;41:166–70.  
19

20 [16] Gerhardt KJ, Abrams RM. Fetal exposures to sound and vibroacoustic stimulation. J Perinatol  
21 2000;20:S21-30.  
22

23 [17] Richards DS, Frentzen B, Gerhardt KJ, McCann ME, Abrams RM. Sound levels in the human  
24 uterus. Obstet Gynecol 1992;80:186–90.  
25

26 [18] Chen X, Striano T, Rakoczy H. Auditory-oral matching behaviour in newborns. Dev Sci 2004;7:  
27 42–7.  
28

29 [19] DiPietro JA, Hilton SC, Hawkins M, Costigan KA, Pressman EK. Maternal stress and affect  
30 influence fetal neurobehavioral development. Dev Psychol 2002;38:659–68.  
31

32 [20] Groome LJ, Swiber MJ, Bentz LS, Holland SB, Atterbury JL. Maternal anxiety during pregnancy:  
33 Effect on fetal behavior at 38 to 40 weeks of gestation. J Dev Behav Pediatr 1995;16:391–6.  
34

35 [21] Kinsella MT, Monk C. Impact of maternal stress, depression and anxiety on fetal  
36 neurobehavioral development. Clin Obstet Gynecol 2009;52:425–40.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 [22] Glover V. Maternal depression, anxiety and stress during pregnancy and child outcome; what  
2 needs to be done. *Best Pract Res Clin Obstet Gynaecol* 2014;28:25–35.  
3

4  
5 [23] Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav*  
6 1983;24:385–96.  
7

8  
9  
10 [24] Honjo S, Arai S, Kaneko H, Ujiie T, Murase S, Sechiyama H, et al. Antenatal depression and  
11 maternal-fetal attachment. *Psychopathology* 2003;36:304–11.  
12

13  
14 [25] Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand*  
15 1983;67:361–70.  
16

17  
18 [26] Reissland N, Francis B, Buttanshaw L. The Fetal Observable Movement System (FOMS). In:  
19 Reissland N, Kisilevsky BS, editors. *Advancing Research on Fetal Development*, New York, NY:  
20 Springer; in press.  
21

22 [27] Ekman P, Friesen WV. *Manual for the Facial Action Coding System*. Paolo Alto, CA: Consultant  
23 Psychologists Press; 1978.  
24

25 [28] Reissland N, Francis B, Aydin E, Mason J, Schaal B. The development of anticipation in the fetus:  
26 A longitudinal account of human fetal mouth movements in reaction to and anticipation of touch.  
27 *Dev Psychobiol* 2013;56: 955–63.  
28

29 [29] Reissland N, Francis B, Mason J, Lincoln K. Do facial expressions develop before birth? *PLOS ONE*  
30 2011;6:e24081.  
31

32 [30] Nijhuis JG, Prechtl HFR, Martin Jr. CB, Bots RSGM. Are there behavioural states in the human  
33 fetus? *Early Hum Dev* 1982;6:177–95.  
34

35 [31] Stroup WW. *Generalized Linear Mixed Models: Modern Concepts, Methods and Applications*.  
36 Boca Raton, FL: Chapman and Hall; 2012.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 [32] Bates DM, Maechler M, Bolker B, Walker S. (2013) lme4: Linear mixed-effects models using  
2 Eigen and S4. R package version 1.0-5. <http://CRAN.R-project.org/package=lme4>  
3

4  
5 [33] Baayen RH, Davidson DJ, Bates DM. Mixed-effects modeling with crossed random effects for  
6 subjects and items. *J Mem Lang* 2008;59:390–412.  
7

8  
9  
10  
11 [34]. Mancuso K. Checking overdispersion when modelling proportions. Response to query on [R-  
12 sig-ME] mailing list. Jan 29, 2012. Online at [https://stat.ethz.ch/pipermail/r-sig-mixed-  
13 models/2012q1/017639.html](https://stat.ethz.ch/pipermail/r-sig-mixed-models/2012q1/017639.html)  
14  
15

16  
17  
18 [35] Bolker, BM, Brooks ME, Clark CJ, Geange SW, Poulsen, JR, Stevens MHH, White JS. Generalized  
19 linear mixed models: a practical guide for ecology and evolution. *Trends Ecol Evol* . 2009; 24:127-  
20  
21  
22  
23  
24 135.

25  
26 [36] Granier-Deferre C, Ribeiro A, Jacquet AY, Bassereau S. Near-term fetuses process temporal  
27 features of speech. *Dev Sci* 2011;14:336–52.  
28  
29

30  
31 [37] Querleu D, Renard X, Boutteville C, Crepin G. Hearing by the human fetus? *Semin Perinatol*  
32  
33  
34 1989;13:409–20.  
35

36  
37 [38] Ferronato PA, Domellöf E, Rönqvist L. Early influence of auditory stimuli on upper-limb  
38 movements in young human infants: an overview. *Front Psychol* 2014;18:1043.  
39  
40

41  
42 [39] Breslow NE, Day NE. *Statistical methods in cancer research, volume II: The design and analysis*  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
of cohort studies. Lyon: International Agency for Research on Cancer, World Health Organisation,  
1987.



**Table Legends**

**Table 1. Average number of mouth stretches per scan and rates of mouth stretching per 100 second of scan [with 95% confidence intervals] for the sound and control groups by gestational age**

**Table 2. Average number of lip puckers per scan and rates of lip puckering per 100 second of scan [with 95% confidence intervals] for the sound and control groups by gestational age**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Figure legends

**Fig. 1.** The effect of gestational age and presentation of an auditory stimulus on the frequency of (a) mouth stretches and (b) lip puckers. Inset images provide examples of (a) mouth stretch and (b) lip pucker movements.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Table 1. Average number of mouth stretches per scan and rates of mouth stretching per 100 second of scan [with 95% confidence intervals] for the sound and control groups by gestational age**



	Average Number of Movements Observed per Scan		Relative Frequency (movements per 100 seconds of scan)	
	32 weeks	36 weeks	32 weeks	36 weeks
<b>Sound</b>	1.67 [0.63, 3.65]	2.67 [1.26, 5.23]	0.89 [0.33, 1.94]	1.38 [0.65, 2.60]
<b>Control</b>	2.50 [1.77, 3.44]	0.64 [0.32, 1.17]	0.44 [0.31,0.60]	0.11 [0.15,0.20]


Note: 95% confidence intervals calculated using Byar's method for rates [39]

**Table 2. Average number of lip puckers per scan and rates of lip puckering per 100 second of scan [ with 95% confidence intervals] for the sound and control groups by gestational age**

	Average Number of Movements Observed per Scan		Relative Frequency (movements per 100 seconds of scan)	
	32 weeks	36 weeks	32 weeks	36 weeks
<b>Sound</b>	0 [0.00, 0.82]	0.33 [0.03, 1.55]	0 [0.00, 0.43]	0.17 [0.02, 0.80]
<b>Control</b>	2.14 [1.47, 3.01]	2.21 [1.53, 3.10]	0.38 [0.26, 0.53]	0.37 [0.26, 0.52]

Note: 95% confidence intervals calculated using Byar's method for rates [39]





Click here to access/download  
**Supplementary Material**  
STROBE-Checklist-v4-MS-Word.doc

