

Durham Research Online

Deposited in DRO:

03 June 2016

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Howard, A.J. and Kinsey, M. and Carey, C. (2015) 'Preserving the legacy of historic metal-mining industries in the light of the Water Framework Directive and future environmental change in mainland Britain : challenges for the heritage community.', *The historic environment : policy practice.*, 6 (1). pp. 3-15.

Further information on publisher's website:

<http://dx.doi.org/10.1179/1756750514Z.00000000061>

Publisher's copyright statement:

This is an Accepted Manuscript of an article published by Taylor Francis Group in *The Historic Environment: Policy Practice* on 13/04/2015, available online at: <http://www.tandfonline.com/10.1179/1756750514Z.00000000061>.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Preserving the Legacy of Historic Metal-Mining Industries in the Light of the Water Framework Directive and Future Environmental Change in mainland Britain: Challenges for the Heritage Community

Andy J. Howard

Landscape Research & Management; and the University of Durham

Mark Kinsey

University of Durham

Chris Carey

University of Brighton

Contemporary global metal mining is a source of environmental pollution, but in Britain it is our historic mining industry that has left a legacy of contamination in the landscape, around both the immediate mine sites as well as within the river valley floors that drain these orefields. It has been estimated that the levels of lead and zinc stored within some northern British river systems represent values comparable to present-day reserves of economically viable ore deposits and exposure to them can be detrimental to human health. Despite the prevalence and significance of these deposits, they have been neglected by the cultural heritage community, in favour of more easily interpretable remains such as mine buildings, technologies of ore procurement and processing, and the final products of manufacture. This paper argues that in the light of future climate change and legislation associated with the EU Water Framework Directive, heritage managers and industrial archaeologists have to start investigating these deposits as part of their studies and to engage with the environmental science and geomorphological communities who are, at present, setting the agenda in terms of strategies for pollution mitigation and landscape remediation.

KEYWORDS Metal Mining, Water Framework Directive, Climate Change, Heritage Management

Introduction

Across the globe, ore deposits have been exploited for their metalliferous minerals since later prehistory, a process that has shaped the social, economic and political development of contemporary human society. Nowhere is this more evident than in Britain, where the origins of mining for the major ores of copper, tin and lead and other associated metals date back beyond the Roman period to the Bronze Age, although its precise character and timing is still the focus of much debate.¹ What is unequivocal is the intensification and peak output of this activity was during the Industrial Revolution of the 18th and 19th centuries.

In England and Wales, it has recently been estimated that there are over 3000 abandoned metal mines,² concentrated in 12 main ore producing regions (Figure 1). Many of these remains form an important component of our landscape of industrial heritage³ and are often used as part of a wider portfolio of historic assets to promote conservation, tourism and education.⁴ Whilst the economic boost that this legacy provides is undoubtedly beneficial, particularly in regions that have long suffered industrial decline, the preservation of these remains has also left a significant legacy of industrial contamination associated with extraction, ore processing and smelting,

around both the immediate sites and across the wider landscape, posing significant risks to natural ecosystems and human health.⁵

Despite the prevalence of industrial remains associated with metal mining across the orefields of Britain and the formation of the *Industrial Archaeology Society* in 1973, until recently few practicing archaeologists have sought to consider the implications of metal mining contamination upon the wider environment. This is despite the fact that archaeologists are probably the best placed profession to understand the intricacy of these remains and likely spatial variability in contamination levels across former mining sites; for example, the ability to identify crushing and dressing floors, which are likely to be highly contaminated. However, despite this expertise, traditional archaeological approaches to this environment have largely focused on recording the fabric of the built environment, the technologies of ore extraction, power, processing and manufacture, and the final metal products themselves.⁶ In the past two decades environmental archaeologists have sought to redress this balance by investigating the link between pollution histories, the historic environmental record and palaeoecology, particularly the exploitation of woodland, which was a source of fuel for ore processing.⁷ Such environmental studies are starting to view geochemical contamination from metalworking processes as an artefact in its own right and a resource to be studied and utilised, for example, Romano-British metalworking geochemistry.⁸ Such concepts have already been established by the osteoarchaeological and geochemical communities who have combined forces to consider the human health effects of historical contamination.⁹

Whilst there is no doubt that this important aspect of our industrial heritage should be preserved, the combined predicted effects of future climate change¹⁰ and, in Europe, stricter pollution control associated with the Water Framework Directive¹¹ provide significant challenges for its longer-term sustainable management. The aim of this paper is to explore these issues by highlighting the scale of potential problems, possible solutions and to consider how the wider heritage community can contribute to this debate.

The Legacy of Pollution

Although the heritage community has shown limited interest in the wider contamination associated with metal mining, it is the geomorphological, geochemical and environmental science communities that have long recognized the importance of these industries for environmental pollution and the sensitivity of these landscapes to change. A number of geomorphological studies have considered specific physical processes of erosion, for example by wind and water around discrete mine sites,¹² whilst the link between geomorphology and wider contamination studies was established in the mid 1970s;¹³ since then, numerous empirical investigations across the orefields of Britain have demonstrated the levels of contamination and the importance of river systems in facilitating the dispersal, storage and remobilisation of metals transported as particulate fractions or binding to sediments in a variety of chemical forms.¹⁴ In addition to sediment transport processes, the spatial distribution and hence concentration of contaminated sediments within valley floors is intimately linked to river planform and style of sedimentation as well as flood regime; these factors have led geomorphologists to distinguish between valley floors characterised by 'passive dispersal' where the addition of contaminated materials seems to have a negligible impact on the natural system and those 'actively transformed', though both are merely end members of a continuous spectrum.¹⁵ Therefore, archaeologists need to be aware of the contaminants within the mine sites themselves as well

as those that have already been dispersed, but are stored within the wider catchment environment (Plate 1a&b).

The scale of particulate contamination of the environment is considerable; it has been estimated that in northern England alone, over 12,000 km² of catchments are affected by the legacy of historic metal mining.¹⁶ Within the Yorkshire Ouse basin, which drains the North Pennine Orefield and is without doubt the most intensely studied metalliferous region of Britain, it is estimated that 620 million tonnes of lead and 640 million tonnes of zinc are stored within its floodplains, representing values comparable to present day reserves of economically viable ore deposits.¹⁷ Within the sub-catchment of the River Swale, it is estimated that approximately 55% of the agriculturally important floodplain is significantly contaminated and that 28% of the lead produced within the catchment remains stored within the valley floor.¹⁸ Scenarios developed using computational modelling suggest that 70% of contaminants deposited within the Swale system have remained for more than 200 years following mine closure¹⁹ and more generally, contamination from historic 'low-scale' mining can still be detected in alluvial sediments over 1000 years after deposition.²⁰

In addition to the particulate load deposited upon the valley floors, metal contaminants can be carried in suspension and solution and are known collectively as 'acid mine drainage'.²¹ A comprehensive government-funded inventory of pollution discharges across England and Wales identified 338 discharges from 4923 abandoned metal mines.²² Whilst contaminant profiles were only available for around 30% of those discharges, they suggested a minimum of 193 tonnes of zinc, 18.5 tonnes of lead, 0.64 tonnes of cadmium, 19.1 tonnes of copper, 551 tonnes of iron, 72 tonnes of magnesium and 5.1 tonnes of arsenic are discharged into surface waters annually.²³

However, other studies have demonstrated the complexity of unravelling the sources of pollution showing that during periods of low flow, diffuse sources are relatively unimportant with more contaminations being released from point source mine waters; in contrast, during periods of higher flow 90% of contaminants are attributed to diffuse sources with contributions from waste materials and groundwater discharge.²⁴

The Impact of Climate Change and the Implementation of the Water Framework Directive

Recent geological history demonstrates that the earth's climate has been shaped by natural cycles of climate change,²⁵ but it is now widely acknowledged that human adaptation of the planet is causing significant changes to current and future climate,²⁶ leading geoscientists to define this period as the Anthropocene.²⁷ Although challenges remain in refining predictions, it seems likely that both the frequency and intensity of weather events will become more pronounced, with extremes of both temperature and rainfall common place.²⁸

Despite levels of contamination being shown to be high, at present the majority of these mining age sediments in northern and western Britain are 'locked down' in the landscape, protected from erosion by vegetation (predominantly rough pasture or improved grassland). However, the autumn 2001 floods in the Yorkshire Ouse basin provided an insight into the potential problems that might occur as a result of future climatic conditions with post-flood point sampling at 35 sites along the River Swale recording lead, zinc and cadmium concentrations exceeding government safety levels.²⁹

In addition to the issues of climate change, the challenges of dealing with the legacy of metal mining have been compounded further by the requirements of the Water Framework Directive.³⁰ Although this significant piece of legislation, which provides a structure for the protection, improvement and sustainable management of the water environment came into effect in December 2000, and has therefore been on the statute books for a considerable period of time, the ramifications of this legislation for heritage are only now beginning to take shape since there is a need for all surface waters to maintain 'good status' by 2015. This status is measured using a range of criteria including ecological targets across 12 River Basin Districts, which provide the strategic level of management and those that fail will need to achieve good status by 2021 or 2027 and the challenges are considerable.³¹

Remediation, its cost and the impact on Heritage

Whilst metal mining is not the only issue driving water quality,³² the empirical evidence described above clearly demonstrates that it has resulted in the introduction of significant volumes of industrial pollutants into the natural environment, but that the environmental science community has developed a thorough understanding of levels of toxicity, as well as current and future pathways of contaminant delivery. However, in contrast to other countries where contemporary mining is causing significant environmental problems,³³ the majority of contamination in Britain is associated with historic mining remains and whilst as point sources they are no longer directly contributing contaminants into the environment, waste tips and mine drainage continue to provide a source of pollutants. In the UK, this historical context creates a key management dilemma since it is only since 1999 that mine operators have been under a legal obligation to mitigate pollution from abandoned mines.³⁴ Even if obligations were met, the financial burden is considerable; a series of joint reports commissioned by DEFRA and the Welsh Government in collaboration with the Environment Agency estimates that at present-day economic rates the total cost to remediate all of the water-related environmental problems associated with abandoned non-coal mines would be around 370 million over an initial 10 year period, excluding operating costs.³⁵

Given the lack of contemporary workings, greater emphasis is placed in the literature on diffuse sources, recognising the importance of contaminants already distributed beyond the mines themselves,³⁶ a point corroborated by a range of empirical studies. For example, analysis of flood sediments after the 2003 floods on the River Swale indicated that the highest concentrations of contaminants were found at the mouths of heavily mined tributaries with elevated levels continuing for 5-10km downstream of those inputs.³⁷

However, whilst focusing on diffuse sources within the floodplains and wider valley floors is important, there seems to have been little attempt to explore diffuse pollution sources within the perimeters of mining sites themselves. This is certainly an area of research where the input of heritage specialists would be exceedingly valuable, especially if levels of contamination could be correlated with industrial processes and archaeological features. Recent work using a combination of multi-spectral satellite imagery, lidar and aerial photography, augmented by ecological survey and geochemical analysis has been undertaken to assess the condition of industrial archaeological remains on Alston Moor in the North Pennines.³⁸ The results indicated extensive processes of gullying, mass movement and deflation all contributing to environmental degradation and acting as pathways for pollutant dispersal. The study also highlighted the importance of water management features within the landscape, including sites of hushing, a

hydraulic mining technique used to expose and clear overburden from above ore deposits³⁹. Whilst water management features were important at the time of mining they clearly have a contemporary role to play as pathways of pollutant movement. This study⁴⁰ demonstrates the potential for detailed large-scale survey and provides an exemplar of how combining geomorphological, geochemical and heritage methodologies can provide a much greater understanding of the landscape rather than considering a mine site as one simple, large point source of pollution. Understanding slope-channel coupling and sediment storage is of paramount importance and Lidar clearly offers a valuable technique to assist in landscape analysis.

As English Heritage note on their website, they have been protecting ‘industrial sites’ since the early 1950s through a combination of both scheduling and listing. This approach has been augmented by initiatives such as the annual ‘*Heritage at Risk*’ programme (established 2008), which has a sub-theme specifically devoted to industrial remains (<http://www.english-heritage.org.uk/caring/heritage-at-risk/industrial-heritage-at-risk/protecting-industrial-sites/>) and the publication of other literature aimed at informing protection and conservation strategies for the industrial environment.⁴¹ However, current protection measures invariably provide a best-fit designation for all site types, periods and locations and in the case of metal mining they don’t consider contaminants although this does come under the umbrella of ‘associated site issues’. At present, with funding from English Heritage, the *National Association of Mining History Organisations* (NAHMO) are in the process of producing ‘*The Research Framework for the Archaeology of the Extractive Industries of England*’ with the aim of informing decision-making with respect to future archaeological research and heritage (management) of mining landscapes. This 18 month programme offers the heritage community a real opportunity to engage with the wider contamination issues of historic metal mining, which is considered as a discrete theme and the research agenda does mention the wider landscape impacts; however, draft chapters available via the NAHMO website (<http://www.namho.org/research.php>) suggest that this opportunity to take a holistic assessment of such remains may not be realized fully.

Having specifically led the way with contaminant studies, the published literature indicates that it is the environmental science community who are largely driving the agenda focused around strategies for mitigation and remediation. To date, the majority of this work has been undertaken to reduce the effects of pollution from coal mines; in 2008, the Environment Agency reported that together with the Scottish Environmental Protection Agency and Coal Authority 54 mine-water treatment plants had been built preventing 2500 tonnes of iron and other metals from entering the hydrological system.⁴² With respect to metal mines, progress has been slower but is now considered a priority with the treatment plant at Wheal Jane tin mine (Cornwall), which prevents 670 tonnes of iron and 150 tonnes of zinc from entering the tidal Restronguet Creek annually, providing an exemplar of what can be achieved.⁴³

Whilst the environmental community has the technological ability to undertake these important schemes, it is unclear from the literature as to how heritage stakeholders are contributing their specialist knowledge to remediation proposals; solutions can be active or passive, dividing into those dealing with particulate waste and those focused upon acid mine drainage.⁴⁴ Where pollution can be identified to a discrete point source, active solutions usually use chemical reagents to facilitate cleaning within treatment plants, whereas passive solutions use a combination of gravity and natural biogeochemical processes operating across artificial lagoons

and wetlands to achieve the same outcome, though in reality, most solutions include a combination of the two.

However, it is estimated that some 60% of metal mining pollution in England and Wales is from diffuse sources.⁴⁵ A popular mitigation proposal in the literature is for relatively discrete areas such as spoil heaps to be capped with solid, inert materials such as waste rock, clay, organic waste or neutralizing agents such as limestone with an optional vegetative cover.⁴⁶ Where spoil heaps are being directly eroded next to adjacent stream channels, rock-filled gabions have been used to reduce erosion, but this is at a considerable cost; approximately £3 million for a 1-2km stretch of the River Nent in Cumbria; where diffuse pollution is shown to be concentrated along particular pathways of runoff, buffer strips (e.g. grassed waterways, infiltration trenches) can be used to prevent runoff entering the river system.⁴⁷

The potential of extreme rainfall events to flush contaminated waters out from mine systems was highlighted by the events at Wheal Jane tin mine in 1992,⁴⁸ a problem that could be amplified under scenarios of future climate change and exacerbated if adits are blocked. Mayes and Jarvis⁴⁹ note the need for precise and accurate mapping of all mines to mitigate this issue but this plea highlights the lack of overlap between the environmental and heritage communities; much of this information is readily available via local authority Historic Environment Records (HERs), the National Mapping Programme (NMP) and volunteer groups such as the Northern Mine Research Society (<http://www.nmrs.org.uk/>).

Whilst the approaches to mitigation are well-developed, the challenges these provide for heritage are many and some scenarios may not be desirable to preservation of the historic environment. For example, technologies designed to deal with point sources require relatively large spaces for infrastructure, which is often lacking within the metal-rich uplands of northern and western Britain. As well as impacting directly on heritage designations (e.g. Scheduled Monuments, Listed Buildings), solutions for both point source and diffuse pollution may impact significantly on other designations related to the aesthetics, ecology and wilderness value (e.g. ANOB, SSSI), especially since many of the areas sit within National Parks and those in Devon and Cornwall have World Heritage Site status. Furthermore, phytotoxic plant communities, which have evolved to withstand high metal concentrations, are well-research phenomena with Calaminarian Grasslands now recognized as a Biodiversity Action Plan habitat type.⁵⁰ Consequently, remediation of heavy metal pollution in these areas would result in the destruction of areas of unique biodiversity (Plate 2), whilst maintenance of these habitats requires a degree of ongoing contamination to be maintained within the environment.

There are further conflicts of interest that may arise from mitigating pollution sources and remediation of existing pollution deposition. Primarily, the metal pollution, whilst a contaminant is also an historic artefact. It provides an insight to the past, in terms of environmental consideration, environmental legislation and economic activities. Indeed, geochemical pollution from the past is actively sought after as a key artefact with which to understand the past.⁵¹

As noted earlier, many of these waste deposits contain higher levels of metal than many commercially extracted ore deposits;⁵² therefore, in a world of diminishing resources, increasing technological innovation, and the need to remove contaminants, it is not inconceivable to consider that future proposals might include the re-mining of deposits, which can have a variety

of uses in addition to their metal content.⁵³ Furthermore, in the North Pennines over 1 million US dollars is currently being invested by one company who believe that *'there is significant untested potential for such mineralisation approximately 300-400m below previous, adit-accessed workings, and such deposits could be significantly larger than any previously discovered'* (<http://www.minco.ie/projects/North%20Pennine%20Project.html>). If proven, such developments would offer significant economic opportunities for this area, which might be difficult to resist and offer significant challenges for the historic environment.

Conclusions

This paper has sought to highlight the challenges facing the metal mining heritage of Britain in the light of future climate change and ever tightening pollution legislation; in Europe, this is associated with the implementation of the Water Framework Directive, but in other parts of the World may involve comparable legislation. These challenges are considerable; in the Swale catchment alone, it is estimated that it will take in excess of 5000 years for all this metal-rich sediment to be removed by natural processes.⁵⁴

Review of the published literature demonstrates the environmental science community has led the way in engaging with the pollution challenges created by the legacy of mining, but that these initiatives appear to take little account of the detailed heritage record. For example, the development of a national strategy for the identification, prioritisation and management of pollution from abandoned non-coal mine sites in England and Wales using a GIS framework to capture mine data from geological and Ordnance Survey maps fails to consider local authority Historic Environment Records⁵⁵; as the authors acknowledge, stakeholder engagement is critical and it is important for the heritage community to engage with and contribute to such initiatives. Where large stakeholder consultations have been undertaken, they have yielded significant results; for example, the mining strategy for Wales, which brought together 20 stakeholder groups to consider the top 50 polluting metal mining sites in the country.⁵⁶

Of course, contaminant problems are not restricted to mainland Britain; historic metal mining is a feature of a number of countries⁵⁷ and climate change will certainly affect all parts of the globe. In Europe, the WFD adds an additional dimension to these issues and is acute where mobilisation of contaminants is transnational⁵⁸, but if metal mining remains across the globe are to be managed in a sustainable way for the long term, heritage professionals need to set an agenda beyond simply preservation.

Acknowledgements

Harry Buglass, formerly of the Institute of Archaeology & Antiquity, The University of Birmingham, drafted Figure 1. Additional thanks must go to Dr Martin Smith (University of Bournemouth) who discussed pollution issues associated with human health and Dr Jen Heathcote (English Heritage) for the clarity of the *'Strategy for Water and Wetland Heritage'* (part of English Heritage's Thematic Research Strategies), which provided an additional impetus to write this paper. The comments of an anonymous referee and Dr Roger White are also acknowledged; however, the views expressed in this paper are solely those of the authors.

Notes

¹ Andrew Dutton, Peter Fasham, David Jenkins, Astrid Caseldine. and Sheila Hamilton-Dyer. 'Prehistoric copper mining on the Great Orme, Llandudno, Gwynedd', *Proceedings of the Prehistoric Society*, 60 (1994), 245-286; Varyl Thorndycraft, Duncan Pirrie, and Antony Brown 'Tracing the record of early alluvial tin mining on Dartmoor, UK' in *Geoarchaeology: exploration, environments, resources*, ed. by Mark Pollard (Geological Society of London, Special Publications 165, 1999), pp.91-102; Tim Mighall, Simon Timberlake, Sarah Clark, and Astrid Caseldine 'A palaeoenvironmental investigation of sediments from the prehistoric mine of Copa Hill, Cwmystwyth, mid Wales', *Journal of Archaeological Science*, 29 (2002), 1161-1188; Tim Mighall, Lisa Dunmayne-Peaty, and David Cranstone 'A record of atmospheric pollution and vegetation change as recorded in three peat bogs from the Northern Pennines PB-Zn Orefield', *Environmental Archaeology*, 9 (2004), 13-38.

² Adam Jarvis, Adrian Fox, Emma Gozzard, Steve Hill, William Mayes, and Hugh Potter 'Prospects for effective national management of abandoned metal mine water pollution in the UK', in *International Mine Water Association Symposium 2007: Water in the Mining Environment*, ed. by R. Cidu and F. Frau,(Mako Edizioni, Cagliari, 2007), pp. 77-81.

³ *Managing the Industrial Heritage*, ed. by Marilyn Palmer, and Peter Neaverson, (Leicester Archaeology Monograph 2, 1995); Peter Neaverson and Marilyn Palmer, *Industrial Archaeology: Principles and Practice* (Routledge, London, 1998); Kate Clarke 'The Workshop of the World. The industrial revolution', in *The Archaeology of Britain. An introduction from the Upper Palaeolithic to the Industrial Revolution*, ed. by John Hunter, and Ian Ralston (Routledge, London, 1999), pp. 280-296; David Cranstone 'Industrial archaeology – manufacturing a new society', in *The Historical Archaeology of Britain* ed. by Richard Newman, David Cranstone, and Christine Howard-Davis, C. (Sutton, Stroud, 2001), pp. 183-210; Marilyn Palmer, Mike Nevell, and Mike Sissons, *Industrial Archaeology, a Handbook* (Council for British Archaeology, York, 2012).

⁴ Robert White, *The Yorkshire Dales Landscapes Through Time* (Batsford/English Heritage, London, 1997); *Dartmoor's Past Tin Industry* (Dartmoor National Park Authority Factsheet, 2001); John Barnatt, and Ken Smith, *The Peak District Landscapes Through Time*. (Batsford/English Heritage, London, 2004).

⁵ Karen Hudson Edwards, Heather Jamieson, and Bernd Lottermoser 'Mine Wastes: Past, Present and Future', *Elements*, 7 (2011), 375-380; Geoff Plumlee, and Suzette Morman 'Mine wastes and human health', *Elements*, 7 (2011), 399-404.

⁶ David Crossley, Priorities for post-Medieval Archaeology in North Derbyshire, in *Recent Developments in the Archaeology of the Peak District*, ed. by Richard Hodges and Ken Smith (Sheffield Archaeology Monograph 2, Sheffield, 1991), pp. 123-134; Palmer, and Neaverson, 1995; Neaverson, and Palmer, 1998; David Cranstone, 'From Newby Hall to Navy Camp: Power, Pots and People in Post-Medieval Archaeology', in *The archaeology of Yorkshire: an assessment at the beginning of the 21st century*, ed. by Terry Manby, Stephen Moorhouse, and Patrick Ottaway (Yorkshire Archaeology Society Occasional Paper 3, Leeds, 2003), pp215-221; Palmer, Nevell, and Sissons,2012.

⁷ Tim Mighall, and Frank Chambers 'The environmental impact of prehistoric mining at Copa Hill, Cwmystwyth, Wales', *The Holocene*, 3 (1993), 260-264; Tim Mighall, and Frank Chambers 'Early Iron working and its impact on the environment: palaeocological evidence from Bryn y Castell Hillfort, Snowdonia, North Wales', *Proceedings of the Prehistoric Society* 63 (1997), 199-219; Tim Mighall *et al.*, 2002; Tim Mighall *et al.*, 2004; Tim Mighall, Simon Timberlake, Ian Forster, and Surjit Singh, Ancient copper and lead pollution records from a raised bog complex in Central Wales, UK, *Journal of Archaeological Science*, 36 (2009), 1504-1515.

⁸ Chris Carey, and Gill Juleff, G. 'Geochemical survey and metalworking: a case study from Exmoor, southwest Britain' in *The World of Iron*, ed. by Jane Humphris, and Thilo Rehren (Archetype, London, 2013), pp.383-392.

⁹ Paul Budd, Janet Montgomery, Jane Evans, and Mark Trickett, 'Human lead exposure in England from approximately 5500 BP to the 16th century AD', *Science of the Total Environment* 318 (2004), 45-58.

¹⁰ Intergovernmental Panel on Climate Change (IPCC) 2013. *Climate change 2013: The Physical Science Basis. Working Group 1 Contribution to the 5th Assessment Report of the IPCC*. (<http://www.ipcc.ch/>); Intergovernmental Panel on Climate Change (IPCC) 2014. *Climate Change 2014. Implications for adaption and vulnerability. Summary for Policy Makers*. (<http://www.ipcc.ch/>).

¹¹ European Commission 2000. Directive 2000/60/EC. *Establishing a framework for community action in the field of water policy*. European Commission PE-CONS 3639/1/100 REV 1, Luxembourg.

¹² Martin Haigh 'Ground retreat and slope evolution on plateau-type colliery spoil mounds at Blaenavon, Gwent', *Transactions of the Institute of British Geographers* 4 (1979), 321-328; Martin Haigh, 'Slope retreat and gullying on revegetated surface mine dumps, Waun Hoscyn, Gwent', *Earth Surface Processes* 5 (1980), 77-79; B.E. Davies, and H.M. White 'Environmental pollution by wind blown lead mine waste: A case study in Wales, UK', *Science of The Total Environment* 20 (1981), 57-74; Graham Merrington, and Brian Alloway 'The transfer and fate of Cd, Cu, Pb and Zn from two historic metalliferous mine sites in the UK' *Applied Geochemistry* 9 (1994), 677-687.

¹³ B.E. Davies, and John Lewin 'Chronosequences in alluvial soils with special reference to historic lead pollution in Cardiganshire, Wales', *Environmental Pollution* 6 (1974), 49-57; John Lewin, B.E. Davies, and P.J. Wolfenden, Interactions between channel change and historic mining sediments' in *River Channel Changes*, ed. by Ken Gregory (Wiley, Chichester, 1977), pp. 353-367.

¹⁴ Mark Macklin 'Floodplain sedimentation in the Upper Axe valley, Mendip, England', *Transactions of the Institute of British Geographers* 10 (1985), 235-244; Steve Bradley and J.J. Cox, 'The significance of the floodplain to the cycling of metals in the River Derwent catchment, UK', *The Science of the Total Environment* 97/98 (1990), 441-454; Mark Taylor 'The variability of heavy metals in floodplain sediments: a case study from mid Wales', *Catena* 28 (1996), 71-87; Paul Brewer, and Mark Taylor 'The spatial distribution of heavy metal contaminated sediment

across terraced floodplains', *Catena* 30 (1997), 229-249; Karen Hudson Edwards, Mark Macklin, and Mark Taylor, 'Historic mining inputs to Tees river sediment', *The Science of the Total Environment* 194/195 (1997), 437-445; Mark Macklin, 'Fluvial geomorphology of North-east England', in *Fluvial Geomorphology of Great Britain*, ed. By Ken Gregory, (Chapman & Hall, London, 1997), pp202-238; Mark Macklin, Karen Hudson-Edwards, and Jo Dawson, 'The significance of pollution from historic mining in the Pennine orefields on river sediment contaminant fluxes to the North Sea', *Science of the Total Environment*, 194/195 (1997), 391-397; Karen Hudson-Edwards, Mark Macklin, Rhona Finlayson, and Dave Passmore, 'Medieval lead pollution in the River Ouse at York, England'. *Journal of Archaeological Science* 26 (1999a), 809-819; Karen Hudson-Edwards, Mark Macklin, and Mark Taylor, '2000 years of sediment borne heavy metal storage in the Yorkshire Ouse basin, NE England, UK'. *Hydrological Processes* 13 (1999b), 1087-1102; Mark Macklin, Mark Taylor, Karen Hudson-Edwards, and Andy Howard, 'Holocene environmental change in the Yorkshire Ouse Basin and its influence on river dynamics and sediment fluxes to the coastal zone', in *Holocene Land-Ocean Interaction and Environmental Change Around the North Sea*, ed. by Ian Shennan, and John Andrews (Geological Society of London Special Publication 166, 2000), pp.87-96; Karen Hudson Edwards, Mark Macklin, Paul Brewer, and Ian Dennis, 'Assessment of Metal Mining-Contaminated river sediments in England and Wales' (Science Report SC030136/SR4, 2008, Environment Agency, Bristol); Heather Jamieson, 'Geochemistry and mineralogy of solid mine waste: essential knowledge for predicting environmental impact', *Elements* 7 (2011), 381-386.

¹⁵ John Lewin, and Mark Macklin, 'Metal mining and floodplain sedimentation in Britain', in *International Geomorphology 1986 Part 1*, ed. by Vince Gardiner (Wiley, Chichester, 1987), pp. 1009-1027; Mark Macklin, Paul Brewer, Karen Hudson-Edwards, Graham Bird, Tom Coulthard, Ian Dennis, Paul Lechler, Jerry Miller, and Jonathan Turner, 'A geomorphological approach to the management of rivers contaminated by metal mining', *Geomorphology* 79 (2006), 423-447.

¹⁶ Mark Macklin, Paul Brewer, Tom Coulthard, Jonathan Turner, Graham Bird, and Karen Hudson-Edwards, 'The chemical and physical impacts of recent mine tailings dam failures on river systems: key issues for sustainable catchment management in former and present mining areas', Proceedings of a seminar on proposed EU Directive on Mine Waste. Office of the Deputy Prime Minister, London, 2002), pp. 18-24.

¹⁷ Karen Hudson-Edwards *et al.*, 1999b.

¹⁸ Paul Brewer, Ian Dennis, and Mark Macklin, 'The use of geomorphological mapping and modelling for identifying land affected by metal contamination on river floodplains', (DEFRA, London, 2005).

¹⁹ Tom Coulthard, and Mark Macklin, 'Modelling long term contamination in river systems from historical metal mining', *Geology* 31 (2003), 451-454.

²⁰ Karen Hudson-Edwards *et al.*, 1999a.

²¹ D. Kirk Nordstrom, 'Mine waters: acidic to circumneutral', *Elements* 7 (2011), 393-398.

²² William Mayes, Hugh Potter, and Adam Jarvis, A.P. 2010, 'Inventory of aquatic contaminant flux arising from historical metal mining in England and Wales', *Science of the Total Environment* 408 (2010), 3576-3583.

²³ William Mayes *et al.*, 2010.

²⁴ Emma Gozzard, William Mayes, Hugh Potter, and Adrian Jarvis, A.P., 'Seasonal and spatial variation in diffuse (non point) source zinc pollution in a historically metal-mined river catchment, UK', *Environmental Pollution* 159 (2011), 3113-3122.

²⁵ John Lowe, and Mike Walker, *Reconstructing Quaternary Environments* (Longman, Harlow, 1997).

²⁶ Intergovernmental Panel on Climate Change 2013; Intergovernmental Panel on Climate Change 2014.

²⁷ Jan Zalasiewicz, Mark Williams, Allan Haywood, and Mike Ellis, 'The Anthropocene: a new epoch of geological time?' *Phil. Trans. R. Soc. A.* 369 (2011), 835-841.; Antony Brown, Stephen Tooth, Richard Chiverrell, Jim Rose, David Thomas, John Wainwright, Joanne Bullard, J.E., Varyl Thorndycraft, Rolf Aalto, and Peter Downs, The Anthropocene: is there a geomorphological case? *Earth Surface Processes and Landforms* 38 (2013), 431-434.

²⁸ Mike Hulme, John Turnpenny, and Geoff Jenkins, '*Climate Change Scenarios for the United Kingdom. The UKCIP02 Briefing Report*' (Tyndall Centre for Climate Research, Norwich, 2002); '*Measuring Progress: Preparing for Climate Change through the UK Climate Impacts Programme*', ed. by Chris West, and Megan Gawith, (UKCIP, Oxford, 2005).

²⁹ Ian Dennis, Mark Macklin, Tom Coulthard, and Paul Brewer, The impact of the October-November 2000 floods on contaminant metal dispersal in the River Swale catchment, North Yorkshire, UK, *Hydrological Processes* 17 (2003), 1641-1657.

³⁰ European Commission 2000. Directive 2000/60/EC. *Establishing a framework for community action in the field of water policy*. European Commission PE-CONS 3639/1/100 REV 1, Luxembourg.

³¹ Environment Agency 2002. *The Water Framework Directive – Guiding principles on the technical requirement* (Bristol, UK); David Johnston, Hugh Potter, Ceri Jones, Stuart Rolley, Ian Watson, and Jim Pritchard, J. 2008. *Abandoned mines and the Water Environment* (Environment Agency, England and Wales, Science Project SC030136/R41, Bristol).

³² Environment Agency 2007. *The unseen threat to water quality: diffuse pollution in England and Wales report – May 2007* (Environment Agency, Bristol).

³³ A.K. Mackay, Mark Taylor, N.C. Munksgaard, Karen Hudson-Edwards, and L. Burn-Nunes 'Identification of environmental lead sources, pathways and forms in a mining and smelting town: Mount Isa, Australia', *Environmental Pollution*, 180b (2013), 304-311.

³⁴ David Johnston *et al.*, 2008.

³⁵ Adam Jarvis, and William Mayes, 'Prioritisation of abandoned non-coal mine impacts on the environment. *The National Picture*' (Report SC030136/R2, 2012a, The Environment Agency, Bristol); Adam Jarvis, and William Mayes, 'Prioritisation of abandoned non-coal mine impacts on the environment. *Future management of abandoned non-coal mine water discharges*' (Report SC030136/R12, 2012b, The Environment Agency, Bristol); William Mayes, and Adam Jarvis, 'Prioritisation of abandoned non-coal mine impacts on the environment. *Hazards and risk management at abandoned non-coal mine sites*' (Report SC030136/R13, 2012c, The Environment Agency, Bristol).

³⁶ Mark Macklin *et al.*, 2006; Karen Hudson Edwards *et al.*, 2008.

³⁷ Ian Dennis *et al.*, 2003

³⁸ Mark Kincey, Lesley Batty, Henry Chapman, Ben Gearey, and Stuart Ainsworth, 'Assessing the changing condition of industrial archaeological remains on Alston Moor, UK, using multisensory remote sensing', *Journal of Archaeological Science* 45 (2014), 36-51.

³⁹ David Cranstone, 'To hush or not to hush: where, when, and how?', in *Men, Mines and Minerals of the North Pennines*, ed. by B. Chambers (Friends of Killhope), pp. 41-48.

⁴⁰ Mark Kincey *et al.*, 2014

⁴¹ English Heritage 'Science for Historic Industries: Guidelines for the investigation of 17th-19th century industries' (English Heritage Publishing, London, 2006); English Heritage 'Conservation Bulletin. A Bulletin of the Historic Environment' (Issue 67, 2011, English Heritage, London).

⁴² David Johnston *et al.*, 2008

⁴³ David Johnston *et al.*, 2008

⁴⁴ Adam Jarvis, and William Mayes, 2012a.

⁴⁵ Adam Jarvis, and William Mayes, 2012b.

⁴⁶ Karen Hudson Edwards *et al.*, 2011, 375-380; Bernd Lottermoser, 'Recycling, reuse and rehabilitation of mine wastes', *Elements* 7 (2011), 405-410.

⁴⁷ Adam Jarvis, and William Mayes, 2012b.

⁴⁸ William Mayes, and Adam Jarvis, 2012c.

⁴⁹ William Mayes, and Adam Jarvis, 2012c.

⁵⁰ Ant Maddock, 'BAP 2011. *UK Biodiversity Action Plan, Priority Habitat Descriptions*, (JNCC. 2011, http://jncc.defra.gov.uk/PDF/UKBAP_PriorityHabitatDesc-Rev2011.pdf).

⁵¹ John Maskall, Keith Whitehead, and Iain Thornton, 'Heavy metal migration in soils and rocks at historical smelting sites', *Environmental Geochemistry and Health* 17 (1995), 127-138; Andy

Meharg, Kevin Edwards, Edward Schofield, Andrea Raab, Joerg Feldmann, Annette. Moran, Charlotte Bryant, Barry Thornton, and Julian Dawson, First comprehensive peat depositional records for tin, lead and copper associated with the antiquity of Europe's largest cassiterite deposits, *Journal of Archaeological Science* 39 (2012), 717-727; Chris Carey, Helen Wickstead, Gill Juleff, Jens Anderson, and Martyn Barber, 'Geochemical survey and metalworking: analysis of chemical residues derived from experimental non-ferrous metallurgical processes in a reconstructed roundhouse', *Journal of Archaeological Science* 49 (2014), 383-397.

⁵² Karen Hudson Edwards *et al.*, 2011.

⁵³ Bernd Lottermoser 2011.

⁵⁴ Ian Dennis, Tom Coulthard, Paul Brewer, and Mark Macklin, 'The role of floodplains in attenuating contaminated sediment fluxes in formerly mined drainage basins', *Earth Surface Processes and Landforms* 34 (2009), 453-466.

⁵⁵ William Mayes, David Johnston, Hugh Potter, and Adam Jarvis, 'A national strategy for identification, prioritisation and management of pollution from abandoned non-coal mines in England and Wales. I. Methodology, development and initial results', *Science of the Total Environment* 407 (2009), 5435-5447.

⁵⁶ Environment Agency Wales 2002. *Executive Summary: Metal Mine Strategy for Wales* (Environment Agency Wales, Cardiff).

⁵⁷ John Grattan, David Gilbertson, and Chris Hunt, 'The local and global dimensions of metalliferous pollution derived from a reconstruction of an eight thousand year record of copper smelting and mining at a desert-mountain frontier in southern Jordan', *Journal of Archaeological Science*, 34 (2007), 83-110

⁵⁸ Graham Bird, Paul Brewer, and Mark Macklin, 'Management of the Danube drainage basin: implications of contaminant metal dispersal for the implementation of the EU Water Framework Directive', *International Journal of River Basin Management* 8 (2010), 63-78.

Note on contributors

Andy J. Howard was a Senior Lecturer for over a decade at the University of Birmingham before the closure of the Institute of Archaeology & Antiquity in 2013. He now runs his own landscape research consultancy as well as holding a Honorary Fellowship within the Department of Archaeology at the University of Durham.

Mark Kinsey worked for over a decade in field archaeology at the University of Birmingham before leaving to pursue a PhD based within the Department of Geography at Durham, focusing upon the geomorphology of mining landscapes in the North Pennines.

Chris Carey gained a PhD in archaeometallurgy from the University of Exeter and is now a Senior Lecturer in Geoarchaeology within the Department of Environment and Technology at the University of Brighton.

Correspondence to: Andy J. Howard, Landscape Research & Management, 24 Russell Close, Stanmore, Bridgnorth, Shropshire, WV15 5JG, UK. Email: andy.howard@landscape-research-management.co.uk; a.j.howard@durham.ac.uk

LIST OF FIGURES

Figure 1. The major orefields of mainland Britain

Plate 1 a&b. Examples of erosion of mine tailings around Dunfell Hush, river Tees catchment, North Pennines (photographs, M. Kinsey).

Plate 2. Spring sandwort on spoil at Whitesike Mine, river South Tyne catchment, North Pennines. Also known as Vernal sandwort or Leadwort, it is frequently found associated with Calaminarian Grassland habitats (photograph, M. Kinsey).




Figure 1