

## Durham Research Online

---

### Deposited in DRO:

13 March 2017

### Version of attached file:

Published Version

### Peer-review status of attached file:

Peer-reviewed

### Citation for published item:

DeGiuli, E. and McElwaine, J. N. (2016) 'Comment on "What determines the static force chains in stressed granular media?"', *Physical review letters.*, 117 (15). p. 159801.

### Further information on publisher's website:

<https://doi.org/10.1103/physrevlett.117.159801>

### Publisher's copyright statement:

Reprinted with permission from the American Physical Society: *Physical Review Letters* 117, 159801 © 2016 by the American Physical Society. Readers may view, browse, and/or download material for temporary copying purposes only, provided these uses are for noncommercial personal purposes. Except as provided by law, this material may not be further reproduced, distributed, transmitted, modified, adapted, performed, displayed, published, or sold in whole or part, without prior written permission from the American Physical Society.

### Additional information:

## Use policy

---

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

## Comment on “What Determines the Static Force Chains in Stressed Granular Media?”

Generic packings of frictional disks or spheres are hyperstatic: there are more degrees of freedom in the contact forces than are fixed by the conditions of mechanical equilibrium. This implies that the history of packing preparation affects the internal structure and elastic response of frictional granular materials [1,2]. Such studies imply that a measure of packing fabric is necessary to model the solid behavior of granular materials, but the form of relevant internal variables is debated [3–5]. In a recent Letter [6], Gendelman, Pollack, Procaccia, Sengupta, and Zylberg (GPPSZ) claim to solve a strong version of this problem, presenting an equation from which the forces can be determined, given the positions of all particles and their radii. GPPSZ emphasize that their result (i) does not require the transverse force law, and (ii) does not involve the distances between particle centers, since these cannot be precisely determined in experiments. While their analysis, and claim (i), are correct, we show here that claim (ii) is false; in fact, the solution proposed by GPPSZ requires that particle radii and positions are known to the precision of the deformations at contacts, a feat impossible in experiments.

This result can simply be established by a scaling analysis of the equations in Ref. [6]. We take units with the mean grain diameter  $\langle\sigma\rangle = 1$  and rescale applied forces and torques by the pressure,  $p$ , which must also be the scale of the contact forces  $|f\rangle$  to be determined. Then the main equation of Ref. [6] takes the form

$$G|f\rangle = p \left\langle \begin{array}{c} -|F^{\text{ext}}\rangle \\ \left(\frac{\kappa}{p}\right)Q|\sigma\rangle \end{array} \right\rangle, \quad (1)$$

where  $|F^{\text{ext}}\rangle$  is a vector of rescaled external forces and torques,  $|\sigma\rangle$  is a vector of geometrical quantities linearly related to grain radii,  $Q$  and  $G$  are matrices that involve  $\mathcal{O}(1)$  geometrical quantities, and  $\kappa \gg p$  is the grain stiffness. The term involving  $Q|\sigma\rangle$  contains the nontrivial geometrical constraints, one for each loop in the packing. For the linear elastic forces considered by GPPSZ, the quantity  $\Delta \equiv p/\kappa$  is the typical magnitude of grain deformations; for a typical experiment,  $\Delta \lesssim 10^{-5}$  [7]. Thus from (1) one would naively expect that either (i)  $|f\rangle \sim p/\Delta$ , which is far too large, or (ii)  $G^{-1}$  has an anomalously small projection onto  $Q|\sigma\rangle$ , which is impossible since  $G$  is nominally independent of geometry at the scale of the contacts, in particular, the scale  $\kappa$ . In fact neither of these possibilities occurs: the mechanism by which  $|f\rangle \sim p$ , as required, is that the vector  $Q|\sigma\rangle$  must be  $\mathcal{O}(\Delta)$  *everywhere in the packing*. This is equivalent to the statement that all grain radii and grain positions must be specified to a precision  $\mathcal{O}(\Delta)$ , the scale of particle deflections. If one had access to such data, one could determine the normal forces directly, without invoking Eq. (1).




FIG. 1. (a) Relative error in predicted forces versus noise  $\xi$ , for  $\Delta \approx 10^{-6}$ . (b) Collapse of all data  $\langle(f_{\text{pred}} - f)^2\rangle^{1/2}/\langle f\rangle$  versus  $\xi/\Delta$ . Dashed line is  $0.3\xi/\Delta$ .

To quantitatively demonstrate our result, we implemented Eq. (1) and tested its susceptibility to simulated experimental noise. With a standard DEM code we prepared packings over a range of  $\Delta$  from  $10^{-6}$  to  $10^{-1}$  [8]. We then added Gaussian noise of amplitude  $\xi$  to the grain radii, and measured the relative error in predicted contact forces  $f_{\text{pred}}$  from the true ones,  $f$  (Fig. 1). As expected from Eq. (1), only when  $\xi \lesssim \Delta$  is the relative error  $\langle(f_{\text{pred}} - f)^2\rangle^{1/2}/\langle f\rangle$  much smaller than 1; for larger noise the predicted forces are not even of the correct magnitude.

We have shown that Eq. (1) is not useful for obtaining *all* the forces from geometrical information. If, somehow, the normal forces are known, then the equilibrium equations can be used to obtain the transverse forces, showing that the transverse force law is indeed redundant.

E. DeGiuli<sup>1</sup> and J. N. McElwaine<sup>2</sup>

<sup>1</sup>Institute of Theoretical Physics  
Ecole Polytechnique Fédérale de Lausanne (EPFL)  
CH-1015 Lausanne, Switzerland

<sup>2</sup>Department of Earth Sciences, Durham University  
Science Labs, Durham DH1 3LE, United Kingdom

Received 3 May 2016; published 6 October 2016

DOI: 10.1103/PhysRevLett.117.159801

- [1] I. Agnolin and J.-N. Roux, *Phys. Rev. E* **76**, 061302 (2007).
- [2] A. P. Atman, P. Claudin, G. Combe, and R. Mari, *Europhys. Lett.* **101**, 44006 (2013).
- [3] S. F. Edwards and D. V. Grinev, *Physica (Amsterdam)* **302A**, 162 (2001).
- [4] R. C. Ball and R. Blumenfeld, *Phys. Rev. Lett.* **88**, 115505 (2002).
- [5] E. DeGiuli and C. Schoof, *Europhys. Lett.* **105**, 28001 (2014).
- [6] O. Gendelman, Y. G. Pollack, I. Procaccia, S. Sengupta, and J. Zylberg, *Phys. Rev. Lett.* **116**, 078001 (2016).
- [7] For Hertzian contacts,  $\Delta = (p/E)^{2/3}$ , where  $E$  is the grain Young modulus. Within a 20 cm deep sandpile,  $\Delta \approx 10^{-5}$ .
- [8] E. DeGiuli, J. McElwaine, and M. Wyart, *Phys. Rev. E* **94**, 012904 (2016).