

Durham Research Online

Deposited in DRO:

16 August 2018

Version of attached file:

Published Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Khoze, V.A. and Martin, A.D. and Ryskin, M.G. (2018) 'Elastic and diffractive scattering at the LHC.', Physics letters B., 784 . pp. 192-198.

Further information on publisher's website:

<https://doi.org/10.1016/j.physletb.2018.07.054>

Publisher's copyright statement:

© 2018 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP3.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.



Elastic and diffractive scattering at the LHC

V.A. Khoze^{a,b}, A.D. Martin^{a,*}, M.G. Ryskin^{a,b}

^a Institute for Particle Physics Phenomenology, University of Durham, Durham, DH1 3LE, United Kingdom of Great Britain and Northern Ireland

^b Petersburg Nuclear Physics Institute, NRC Kurchatov Institute, Gatchina, St. Petersburg, 188300, Russia

ARTICLE INFO

Article history:

Received 19 June 2018

Received in revised form 17 July 2018

Accepted 29 July 2018

Available online 31 July 2018

Editor: A. Ringwald

ABSTRACT

Inspired by the new TOTEM data on elastic pp scattering at 13 TeV, we study the possibility to describe all the diffractive collider data (σ_{tot} , $d\sigma_{\text{el}}/dt$, $\rho \equiv \text{Re}A/\text{Im}A$, $\sigma_{\text{low}M}^D$) in a wide interval of energy (0.0625 to 13 TeV) in the framework of a two-channel eikonal model. We show that a satisfactory description can be achieved without an odd-signature (Odderon) exchange contribution. We consider the possible role of the QCD Odderon which may improve the description of ρ and discuss the importance of the odd-signature term if the amplitude were to exceed the black disc limit.

© 2018 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Motivation

TOTEM have recently published very detailed data on elastic proton–proton scattering at 13 TeV at the LHC, covering the very low t region¹ and up to the region of the diffractive dip and well beyond [1,2]. The goal of this paper is to describe these data together with elastic and diffractive data at other collider energies. We use a two-channel eikonal model. In addition to the dominant even-signature amplitude, we discuss the role of the odd-signature (Odderon) amplitude. Moreover we study the present situation concerning information on low-mass diffractive dissociation. Recall that the multi-channel eikonal model is written, in the Good–Walker formalism [3], in terms of diffractive eigenstates; and the experimental information on low-mass diffraction, $\sigma_{\text{low}M}^D$, controls the relative contributions of the different diffractive eigenstates.

Finally, we discuss the high-energy behaviour of the elastic amplitude for a central collision. That is, at impact parameter $b = 0$.

2. Description on the model

We use a two-channel eikonal model which is based on the two-particle unitarity equation,

$$2\text{Im}A_{\text{el}}(b) = |A_{\text{el}}(b)|^2 + G_{\text{inel}}(b), \quad (1)$$

* Corresponding author.

E-mail address: a.d.martin@durham.ac.uk (A.D. Martin).

¹ The behaviour of the cross section at very low t samples Coulomb interference and allows a measure of the real part of the amplitude.

in impact parameter, b , space. This accounts for the possibility of proton dissociation $p \rightarrow N^*$, not only in the final state, but also in an intermediate state. That is the model includes rescattering like $p \rightarrow N^* \rightarrow N^* \rightarrow p$ etc. The beautiful and convenient way to accomplish this is to use the Good–Walker formalism [3], and to introduce diffractive eigenstates $|\phi_i\rangle$ which diagonalize the diffractive amplitude

$$\langle \phi_i | A | \phi_k \rangle = A_{ik} = A_i \delta_{ik}. \quad (2)$$

For each individual eigenstate the elastic amplitude is given by the one-channel eikonal expression

$$A(b) = i \left(1 - e^{-\Omega(b)/2} \right). \quad (3)$$

The incoming ‘beam’ proton wave function is written as a superposition of the diffractive eigenstates

$$|p\rangle = \sum a_i |\phi_i\rangle, \quad (4)$$

and similarly for the incoming ‘target’ proton. In this formalism the pp elastic cross section is given by

$$\frac{d\sigma_{\text{el}}}{dt} = \frac{1}{4\pi} \left| \int d^2b e^{i\mathbf{q}_t \cdot \mathbf{b}} \sum_{i,k} |a_i|^2 |a_k|^2 (1 - e^{-\Omega_{ik}(b)/2}) \right|^2, \quad (5)$$

where $-t = q_t^2$, and the opacity $\Omega_{ik}(b)$ corresponds to the interaction between states ϕ_i and ϕ_k . Also the ‘total’ low-mass diffractive cross section is of the form

$$\sigma_{\text{el+SD+DD}} = \int d^2b \sum_{i,k} |a_i|^2 |a_k|^2 \left| (1 - e^{-\Omega_{ik}(b)/2}) \right|^2, \quad (6)$$

where SD includes the single dissociation of one or the other proton, and DD is the cross section for events where both protons dissociate. So the low-mass diffractive dissociation cross section is

$$\sigma_{\text{lowM}}^D = \sigma_{\text{el+SD+DD}} - \sigma_{\text{el}}, \quad (7)$$

where $\sigma_{\text{el+SD+DD}}$ corresponds to all possible low-mass dissociation caused by the dispersion of the Good–Walker eigenstate scattering amplitudes. A more detailed description of the model is given in [4].

As mentioned above, we use a two-channel eikonal, $i, k = 1, 2$. Each eigenstate has its own coupling v_i to the Pomeron, with its own t dependence parametrised in the parametric form

$$F_i(t) = \exp(-(b_i(c_i - t))^{d_i} + (b_i c_i)^{d_i}), \quad (8)$$

where c_i is added to avoid the singularity t^{d_i} in the physical region of $t < 4m_\pi^2$. Note that $F_i(0) = 1$. The six parameters b_i , c_i , d_i , together with the intercept and slope of the Pomeron trajectory are tuned to describe the elastic scattering data, paying particular attention to the observed energy behaviour of σ_{lowM}^D , at all available collider energies, \sqrt{s} .

The opacity Ω_{ik} corresponding to the scattering between eigenstates $\langle \phi_i |$ and $|\phi_k \rangle$ is given by one-Pomeron exchange

$$\Omega_{ik}(b) = \int \frac{d^2q_t}{4\pi^2} e^{iq_t \cdot b} \Omega_{ik}(t = -q_t^2), \quad (9)$$

with

$$\Omega_{ik}(t) = v_i F_i(t) v_k F_k(t) \left(\frac{s}{s_0} \right)^{\alpha_P(t) - 1} \quad (10)$$

and $s_0 = 1 \text{ GeV}^2$.

3. Low-mass proton dissociation

Note that if the amplitudes are identical, $A_i = A$, then the interaction will not destroy the coherence of the original proton wave function (4). Then the final state that we observe will be only the proton, while the probability of dissociation given by σ_{lowM}^D will be zero. That is, a larger value of σ_{lowM}^D indicates a larger dispersion between the amplitudes A_i .

A model with a large number of Good–Walker components may account for different proton excitations and in this way describe $d\sigma/dM_X$, where M_X is the mass of the system after the $p \rightarrow X$ dissociation. In our t -channel eikonal analysis we use only one effective N^* state, assuming that it includes all the excitations up to $M_X = 3.4 \text{ GeV}$, the mass value used by TOTEM collaboration [5] to separate proton dissociations into low- and high-mass states.²

Experimentally the situation for measurements of σ_{lowM}^D is far from clear. At the relatively low [6,7] and ISR energies [8–12]

$$\sigma_{\text{lowM}}^D \sim 0.3 \sigma_{\text{el}} \sim 2.5 \text{ mb}, \quad (11)$$

while at 7 TeV TOTEM [5] reported a much smaller value for the ratio $\sigma_{\text{lowM}}^D/\sigma_{\text{el}}$

$$\sigma_{\text{lowM}}^D \sim 2.6 \pm 2.2 \text{ mb} \sim 0.1 \sigma_{\text{el}}. \quad (12)$$

Recall that stronger absorptive corrections can decrease the ratio.

² High-mass dissociation is described separately in terms of triple-Regge diagrams.

The situation at 13 TeV is not so evident. At the moment there are no TOTEM data for σ_{lowM}^D . However we can compare the values of the inelastic cross sections measured by ATLAS [13] and CMS [14] with the total and elastic cross section given by TOTEM [15]. A small complication is that ATLAS measure $\sigma_{\text{inel}} = 68.1 \pm 0.6 \pm 1.3 \text{ mb}$, using events where at least one particle carries a momentum fraction $\xi > 10^{-6}$. This corresponds to $M_X > 13 \text{ GeV}$. On the other hand, CMS use the CASTOR detector to cover the region down to $\xi > 10^{-7}$ on one side of the interaction point. In other words CMS collect events for all processes except for the possible dissociative $pp \rightarrow X + Y$, with $M_Y < 4.1 \text{ GeV}$ on one side and $M_X < 13 \text{ GeV}$ on the other side [14]. If we compare the two CMS results then we can estimate

$$\frac{d\sigma^D}{d\ln(M_X^2)} = \frac{\sigma_{\text{inel}}(\xi_Y > 10^{-7}, \xi_X > 10^{-6}) - \sigma_{\text{inel}}(\xi_X, \xi_Y > 10^{-6})}{2\ln(M_X/M_Y)} \quad (13)$$

$$= \frac{68.6 - 67.5}{2\ln(13/4.1)} \simeq 0.48 \text{ mb}. \quad (14)$$

This CMS number is in agreement with the ATLAS data [16] at 7 TeV on $d\sigma/d\Delta\eta^F$ for events with rapidity gaps and with the theoretical estimates of [17,4]. Taking (14), we can evaluate the cross section of events with $M_{X,Y} < 3.4 \text{ GeV}$ to be equal to 70.1 mb. Thus the cross section of dissociation up to the canonical $M_X = 3.4 \text{ GeV}$ is

$$\sigma_{\text{lowM}}^D = \sigma_{\text{tot}} - \sigma_{\text{el}} - \sigma_{\text{inel}}(M_X > 3.4) \quad (15)$$

$$= 110.6 - 31 - 70.1 \simeq 9 \text{ mb}. \quad (16)$$

A comparison of (12) and (16) shows that the value of σ_{lowM}^D increases about three times in the relatively small energy interval from 7 to 13 TeV. This is very strange. Within this rather small $\ln s$ interval we expect the variation of σ_{lowM}^D to be of the order of 0.5 mb. Note, however, that the estimate (16) is obtained from the difference of two large numbers coming from different experiments³ with their own normalization uncertainties like $\pm 3.4 \text{ mb}$ for σ_{tot} (TOTEM) and $\pm 1.6 \text{ mb}$ for σ_{inel} (CMS).⁴ The results of the model description that we shall present in Section 6 give $\sigma_{\text{lowM}}^D = 5.0$ and 5.4 mb at 7 and 13 TeV respectively. In view of the uncertainties just discussed above, the model values are consistent with all data.

4. Real part of the elastic pp amplitude

At high energy the elastic scattering amplitude is dominantly imaginary. The ratio $\text{Re}A/\text{Im}A$ is about 0.1 and the real part plays a very small role in the low t region (except of the Coulomb interference). Nevertheless for a detailed description of the present very precise data we must account for this contribution. Therefore in (5) we have to keep the full complex opacity $\Omega_{ik}(b)$ in the formula for the elastic amplitude

$$A_{ik}(b) = \left(1 - e^{-\Omega_{ik}(b)/2} \right). \quad (17)$$

For Pomeron exchange we have the even-signature factor

$$\eta_{\text{even}} = [1 + \exp(-i\pi\alpha_P(t))] \quad (18)$$

³ If we replace $\sigma_{\text{inel}}(\xi > 10^{-6}) = 67.5 \pm 1.6(\text{lumi}) \text{ mb}$ [14] by the ATLAS value of $68.1 \pm 1.3(\text{lumi}) \text{ mb}$ [13] then we find the bit smaller value of about 8 mb in (16).

⁴ It is also possible that the value of $d\sigma^D/d\ln(M_X^2)$ may be a bit larger for a lower M_X value, in particular due to secondary Reggeon contributions, see Fig. 9 of [17]. This would enlarge σ_{inel} and therefore decrease σ_{lowM}^D a little; though, however, less than 0.5 mb.

for the ‘even’ part of the opacity $\Omega_{ik}(b)$, where $\alpha_P(t)$ is the Pomeron trajectory. If we keep only the even-signature contribution, the real part of the elastic amplitude satisfies the usual dispersion relation and its value can be calculated at $t = 0$ from the known total cross section. Indeed, up to collider energies of $\sqrt{s} = 8$ TeV, the experimental results for $\rho \equiv \text{Re}A/\text{Im}A|_{t=0}$ are consistent with those deduced from the dispersion relation for the even-signature amplitude (see, for example, [18]).

However, at 13 TeV the TOTEM collaboration [1] have reported a measurement of $\rho = 0.09 - 0.10$ with an uncertainty ± 0.01 which is significantly lower than $\rho = 0.135$ expected by the conventional COMPETE analysis [19]. If this difference were to be explained in the even-signature approach, it would indicate a slower growth of the total cross section with \sqrt{s} than that given by the COMPETE parametrization, as stated in [15]. On the other hand, the TOTEM [15] measured value of the total cross section at 13 TeV

$$\sigma_{\text{tot}} = 110.6 \pm 3.4 \text{ mb} \quad (19)$$

is even a bit larger than that given by COMPETE [19].

5. Odderon exchange

Another way to obtain a smaller value of ρ is to include the odd-signature (Odderon) contribution in the opacity $\Omega_{ik}(b)$. The odd-signature factor with α_{Odd} close to 1

$$\eta_{\text{odd}} = [1 - \exp(-i\pi\alpha_{\text{Odd}}(t))] \quad (20)$$

gives an almost real contribution to the elastic amplitude. The Odderon is expected in perturbative QCD,⁵ see in particular [20–22]. However the naive estimates show that its contribution is rather small; say, $\Delta\rho_{\text{odd}} \sim 1\text{mb}/\sigma_{\text{tot}} \lesssim 0.01$ [23] at the LHC energies. The discovery of the long-awaited, but experimentally elusive, Odderon would be very welcome news for the theoretical community. Indeed, there have been several attempts to prove its existence experimentally (see, for example, [24–26] for comprehensive reviews and references).

It is important to note that the Odderon contribution must be included in the opacity $\Omega_{ik}(b)$, and not directly in the elastic amplitude, since (17) is the general form of the solution of the two-particle unitarity equation where Ω_{ik} includes the *full* two-particle irreducible component of the interaction amplitude. Provided we include the odd-signature contribution to $\Omega_{ik}(b)$ via (17) we automatically account for the absorptive effect caused by elastic rescattering.⁶

6. Results for the model description of the data

Using the two-channel eikonal model with a small set of parameters, we attempt to describe all the diffractive data (σ_{tot} , $d\sigma_{\text{el}}/dt$, $\rho \equiv \text{Re}A/\text{Im}A$, $\sigma_{\text{low}M}^D$) over a wide range of collider energies (from $\sqrt{s} = 0.0625$ to 13 TeV) and a large interval of t from 0 up to 1 GeV^2 . The data correspond to more than four orders of magnitude variation for $d\sigma_{\text{el}}/dt$.

In the model the proton is described by a superposition of two diffractive eigenstates, (4), with form factors parametrised as in (8) and with coupling to Pomeron exchange given by

$$v_{1,2} = \sqrt{\sigma_0} (1 \pm \gamma). \quad (21)$$

⁵ QCD is the $SU(N=3)$ gauge theory which contains the spin = 1 particle (gluon) and (for $N > 2$) the symmetric colour tensor, d^{abc} . Due to these facts in perturbative QCD there exists a colourless C -odd t -channel state (formed from three gluons) with intercept, α_{Odd} , close to 1.

⁶ More details on the inclusion of the Odderon can be found in [27,28].




Fig. 1. The 2013 description of the pp (and $p\bar{p}$) elastic data up to 7 GeV in model 2 of [4], together with the prediction for 13 TeV. The references for the data are given in [4]; note that the Tevatron experiments cover data at 1.8 and 1.96 TeV. The recent TOTEM 13 TeV data [1,2] are superimposed on the plot; they are hard to distinguish from the prediction, except for an interval about $t = -0.3 \text{ GeV}^2$ where they lie above.

The Pomeron trajectory is parametrised as

$$\alpha_P(t) = 1 + \Delta + \alpha'_P t. \quad (22)$$

In addition to the constant slope, α'_P , of the Pomeron trajectory, we insert the π -loop contribution as proposed in [32], implemented as in [33,34]. The parameter Δ embodies the BFKL effects which give $\Delta \sim 0.2 - 0.3$ and the renormalization⁷ caused by the Pomeron loop insertions which decrease the resulting values of Δ .

The original implementation of this model [4] described all the diffractive data existing up to 2013 in terms of only even-signature exchange. That is only the Pomeron contribution to $\Omega_{ik}(b)$. At that time there were a few local χ^2 minima in parameter space corresponding to equally good descriptions of the data. In fact we found, and presented [4], four versions of the model which gave good descriptions of all the elastic data available up to 7 TeV. In Fig. 1 we reproduce the 2013 description of the elastic data up to 7 TeV given by version 2 of the model, which also showed the prediction made for $d\sigma_{\text{el}}/dt$ at 13 TeV. In addition we have superimposed on this figure the recent TOTEM measurement made at 13 TeV. The agreement at 13 TeV is surprisingly good. So what is the problem? The deficiency is in values of $\sigma_{\text{low}M}^D$. Version 2 of the model has values of $\sigma_{\text{low}M}^D$ which are too small and no longer in agreement with the experimental information on $\sigma_{\text{low}M}^D$ discussed in Section 3. The version 2 values of the cross section for low-mass diffractive dissociation are $\sigma_{\text{low}M}^D = 1, 2.8, 3.1 \text{ mb}$ at 0.0625,

⁷ Note that multi-Pomeron diagrams, in particular fan diagrams, also have an effect on the ‘effective’ Pomeron form factor, (8), tuned to describe the data. We do not explicitly include these multi-Pomeron diagrams in our simplified two-channel eikonal model in order to keep a clear physical structure of the interaction amplitude. The possible effects of these multi-Pomeron diagrams is allowed for by the renormalized parameters of the Pomeron trajectory and the Pomeron Good-Walker eigenstate couplings.




Fig. 2. As for Fig. 1, but now retuning the model to describe also the new TOTEM 13 TeV data [2]. The description at 7 and 13 TeV is shown in more detail in the region of the diffractive dip by the continuous curves in Fig. 3.

7, 13 TeV respectively, which are too small. However, if we retune the values of the parameters of version 2 (in particular, enlarging the value of γ and make small adjustments to the form factors of the diffractive eigenstates) then we obtain the equally good description of the elastic data shown in Fig. 2, together with

$$\sigma_{\text{lowM}}^D = 2.35, 5.0, 5.4 \text{ mb at } 0.0625, 7, 13 \text{ TeV} \quad (23)$$

respectively. Now the values do not contradict the experimental information discussed in Section 3, when we account for the experimental uncertainties. The values of the parameters of the 2013 and 2018 descriptions of the data are shown in Table 1. The observables as a function of energy corresponding to the present description of the data are shown in Table 2. It is informative to show in more detail in Fig. 3 (by continuous curves) the description of Fig. 2 in the region of the diffractive dip for the 7 and 13 TeV data.

6.1. The Odderon contribution

Before we discuss a possible Odderon contribution, we can see from Fig. 4 that, even without an Odderon, the model produces a rather small value of $\rho \equiv \text{Re}A/\text{Im}A = 0.109$ at 13 TeV, more or less compatible with the recent TOTEM result [1]. However what about our model value of σ_{tot} at 13 TeV? The present version of the model (constrained by the experimental information on low-mass proton dissociation, σ_{lowM}^D , of Section 3) has a flatter energy behaviour of the total cross section. We slightly overestimate σ_{tot} at 62.5 GeV and underestimate σ_{tot} at 13 TeV, but are still in agreement with the data to within 1.5σ .

What happens if we include an Odderon contribution? In order not to introduce too many extra parameters, we use the same couplings for the odd-signature terms to the two different diffractive Good–Walker eigenstates. We parametrize the odd-signature amplitude as

$$A_{\text{odd}}(s, t) = s\sigma_{\text{odd}} \exp(B_{\text{odd}}t) \quad (24)$$

Table 1

The values of the parameters in the two-channel eikonal fit to elastic pp scattering data in which particular attention is paid to the value of σ_{lowM}^D and to the behaviour of the GW eigenstates. The values of the parameters in the 2013 column correspond to version 2 of the original 2013 analysis [4] (see Fig. 1), while the last column shows the values corresponding to the present description of the data (see Fig. 2). The first four rows give the values of parameters connected to the Pomeron trajectory and its couplings, and the last seven rows list the parameters which specify the Good–Walker diffractive eigenstates.

	2013	2018
Δ	0.115	0.13
α'_p (GeV^{-2})	0.11	0.052
σ_0 (mb)	33	23
γ	0.4	0.56
$ a_1 ^2$	0.25	0.505
b_1 (GeV^{-2})	8.0	10.0
c_1 (GeV^2)	0.18	0.233
d_1	0.63	0.462
b_2 (GeV^{-2})	6.0	4.9
c_2 (GeV^2)	0.58	0.52
d_2	0.47	0.47

Table 2

The predictions of the elastic and diffractive observables resulting from the description of the presently available data.

\sqrt{s} (TeV)	ρ	σ_{tot} mb	σ_{el} mb	$B_{\text{el}}(t=0)$ (GeV^{-2})	σ_{lowM}^D mb
0.1	0.141	48.3	8.8	12.6	2.6
0.546	0.129	64.6	13.8	14.8	3.5
1.8	0.121	78.1	18.2	16.7	4.2
7	0.113	95.5	24.1	19.1	5.0
8	0.112	97.4	24.7	19.4	5.1
13	0.109	104.2	27.1	20.4	5.4
100	0.099	136.2	38.6	25.4	6.9




Fig. 3. The description of the 7 and 13 TeV data in the region of the diffractive dip enlarged from Fig. 2. In addition, the dashed curves show the effect of the Odderon if its contribution with $\sigma_{\text{odd}} = 1.5$ mb were to be included with the even-signature amplitude corresponding to the model used to obtain Fig. 2.




Fig. 4. The energy dependence of the $\rho = \text{Re}A/\text{Im}A$ ratio. The data are taken from [1,29–31]; the first two data points correspond to $p\bar{p}$ scattering and the last two to pp scattering. The values of ρ given by the model are shown by the solid curve. The dashed ($p\bar{p}$) and dot-dashed (pp) curves correspond to an alternative behaviour of ρ obtained from a ‘global’ description of diffractive data which include a QCD Odderon contribution calculated as described in the text.

where⁸ $B_{\text{Odd}} = 6 \text{ GeV}^{-2}$, and where the normalization corresponds to $\text{Im}A(t=0) = s\sigma_{\text{tot}}$. In other words we consider a QCD Odderon with intercept $\alpha_{\text{Odd}}(0) = 1$, as was obtained in [22], and normalization given by the parameter σ_{Odd} . The simplified lowest α_s calculation leads to $\sigma_{\text{Odd}} = 0.8 \text{ mb}$ [23]. With such a small coupling of $\mathcal{O}(1 \text{ mb})$ the Odderon is almost invisible in Fig. 2.

Recall that, for $\alpha_{\text{Odd}}(0) = 1$, Odderon-exchange is real, see (20). Thus we have essentially no interference term. The Odderon contribution only becomes visible in the dip region (see Fig. 3) where the imaginary part of the even-signature amplitude vanishes; and in the region of very small t where it interferes with the Coulomb (γ -exchange) term.

Note that the Odderon decreases the value of $\rho \equiv \text{Re}A/\text{Im}A$ in pp collisions, while simultaneously enlarging ρ in $p\bar{p}$ collisions, see Fig. 4. Since the Odderon contribution must be added to the opacity and is screened in the full amplitude by Pomeron exchange it affects the value of ρ at 13 TeV less than at 541 GeV where it was measured by the UA4 collaboration [31] for $p\bar{p}$ scattering, again see Fig. 4. In particular, setting the parameter $\sigma_{\text{Odd}} = 1.5 \text{ mb}$ we have $\Delta\rho = -0.005$ at 13 TeV and $\Delta\rho = +0.012$ at 541 GeV.

Recall that we showed the role of the Odderon in the dip region in more detail in Fig. 3. We may conclude that even without the Odderon the model could be tuned to be consistent with the elastic data. However, a small Odderon comparable with the expectations of QCD may improve the agreement with the measurements on ρ , and not spoil the description of $d\sigma_{\text{el}}/dt$ in the dip region, bearing in mind the uncertainties.

⁸ Note that the C-odd and isospin = 0 state does not couple to the pion. Thus the Odderon only feels the centre of the proton, and not the pion cloud. Therefore it is reasonable to assume that the Odderon slope, B_{Odd} , is lower than that for the even-signature (Pomeron) amplitude.

7. Does the pp -amplitude exceed the black disc limit?

Naive predictions based on a Donnachie–Landshoff parametrization [35] show that the black disc limit is exceeded for central ($b=0$) elastic pp collisions at LHC energies. That is, $\text{Im}A(s, b=0) > 1$. It is therefore relevant to ask if the LHC data respect this limit.

Recall that the imaginary part of the high-energy elastic amplitude in impact parameter space is given by

$$\text{Im}A(b) = \frac{1}{(2\pi)^2} \frac{1}{2s} \int \text{Im}A(t) \exp(-ib \cdot q_t) d^2q_t \quad (25)$$

where $q_t = \sqrt{-t}$, and the values of $\text{Im}A(t)$ can be calculated directly from the data for the differential elastic cross section⁹

$$|\text{Im}A(t)|^2 = \frac{16\pi s^2}{1 + \rho^2} \frac{d\sigma_{\text{el}}^N}{|dt|}, \quad (26)$$

with a small contribution ($\sim 1\%$) coming from ρ^2 . In this way we obtain

$$\text{Im}A(b) = \int \kappa \sqrt{\frac{d\sigma_{\text{el}}^N}{|dt|} \frac{16\pi}{1 + \rho^2}} J_0(bq_t) \frac{q_t dq_t}{4\pi} \quad (27)$$

where J_0 is the Bessel function. Noting that $\text{Im}A(t)$ changes sign at the diffractive dip, we have $\kappa = +1$ or -1 for $|t|$ values below or above this point.

There is some uncertainty since we do not know the t behaviour of the Re/Im ratio ρ . On the other hand, the value of $\rho \sim 0.1$ is rather small, and assuming a flat behaviour ($\rho = \text{const}$ within the t interval relevant for the integral (27)) we are able to calculate $A(b)$ with sufficient accuracy. To obtain a rough estimate of $\text{Im}A(b=0)$ we may further simplify (27) by assuming, in the relevant $|t|$ region, that the differential cross section is well described by a simple exponent $d\sigma_{\text{el}}^N/dt \propto \exp(B_{\text{el}}t)$. In such a case we get

$$\text{Im}A(b=0) = \frac{\sigma_{\text{tot}}}{4\pi B_{\text{el}}}, \quad (28)$$

which we evaluate using the published experimental data for σ_{tot} and B_{el} . The results are presented in Table 3, where the errors have been added in quadrature.

It is known that the proton–proton opacity, Ω , increases with energy and correspondingly increases the value of $A(0)$. Moreover it was claimed in [40] that already at $\sqrt{s} = 7 \text{ TeV}$ the value of $\text{Im}A(b=0) > 1$ exceeds the black disc limit $A = i$. The surprising new result is that the value obtained from (28) for the TOTEM data at 13 TeV exceeds the limit by more than 3 standard deviations. If confirmed, what would this mean?

Recall that the expression (3), $A(b) = i(1 - e^{-\Omega(b)/2})$, is the most general solution of the unitarity equation (1). That is, in order to obtain $\text{Im}A(b) > 1$ we need to have $|\text{Im}\Omega(b)| > \pi$. The opacity $\Omega = \Omega_{\text{even}} + \Omega_{\text{odd}}$ contains both the even and odd signature terms. The imaginary contribution to Ω coming from the even-signature part is strongly limited by dispersion relations. It cannot exceed about 0.3–0.5. Such a large value, $|\text{Im}\Omega(b)| > \pi$, can only come from an odd-signature contribution. For an exponential parametrization $A(t) \propto \exp(Bt)$ the value of Ω_{odd} reads

$$\Omega_{\text{odd}}(b) = -i\eta_{\text{odd}} \frac{\sigma_{\text{Odd}}}{4\pi B} e^{-b^2/4B}. \quad (29)$$

⁹ The notation σ^N means that to be precise we have to subtract from the measured elastic cross section the contributions caused by the pure Coulomb interaction and by Coulomb–nuclear interference.

Table 3

The values of central amplitude, $A(b=0)$, obtained from (28) using the total cross sections and elastic slopes measured by UA4 [31], TOTEM [5,15,29,36,37] and ATLAS-ALFA [38,39] collaborations. Note that with present normalization the black disk limit corresponds to $\text{Im}A(b=0) = 1$.

\sqrt{s} TeV	UA4: 0.541	LHC	2.76	7	8	13
$\text{Im}A(0)$	0.84 ± 0.025	TOTEM	1.01 ± 0.043	1.01 ± 0.03	1.045 ± 0.032	1.11 ± 0.032
$\text{Im}A(0)$		ATLAS		0.988 ± 0.02	0.996 ± 0.015	

In order to get $|\text{Im}\Omega_{\text{odd}}| > \pi$ with a reasonable slope $B = 6 \text{ GeV}^{-2}$ we would need the parameter $\sigma_{\text{odd}} > 90 \text{ mb}$! This looks very unlikely.¹⁰ At present there is no model which can produce such a large real amplitude for high-energy pp -scattering.¹¹ If the value of $\text{Im}A(b=0) > 1$ were to be confirmed, it would be an important hint in favour of a completely **new strong** interaction beyond the Standard Model, which has never been observed before (for $\sqrt{s} \lesssim 1 \text{ TeV}$) and reveals itself only in the LHC energy region.¹²

But first we must question the simplified formula (28). This approximation was acceptable at CERN-ISR energies where the position of the diffractive dip was at larger $|t|$, ($|t_{\text{dip}}| \simeq 1.3 \text{ GeV}^2$), and where the maximum value of $d\sigma_{\text{el}}/dt$ after the dip never exceeds $10^{-6} d\sigma_{\text{el}}^N(t=0)/dt$. However, at the LHC the dip occurs at much smaller $|t| \sim 0.5 \text{ GeV}^2$ and the contribution of the negative amplitude, $\text{Im}A(t) < 0$, to the integral (27) after the dip is not negligible. Thus we must calculate the value of $\text{Im}A(b=0)$ more precisely based on (27) and account for the fact that after the diffractive dip (i.e. for $|t| > 0.47 \text{ GeV}^2$ at 13 TeV) the imaginary part of the elastic amplitude changes sign. It turns out that at LHC energies the contribution after the diffractive dip noticeably decreases the value obtained for $\text{Im}A(b=0)$. The improved calculation gives

$$\text{Im}A(b=0) = 1.026. \quad (30)$$

Bearing in mind a normalization uncertainty of about 3% for σ_{tot} , this value is consistent with the statement that the amplitude *does not exceed black disk limit*. In fact we performed the calculation twice. First, we assumed a constant value $\rho = 0.1$ independent of t , and second, we used the values of $\rho(t)$ given by the model described in Sections 2–4. The difference in $\text{Im}A(b=0)$ is negligible (less than 0.002).

8. Conclusions

We have considered the new TOTEM data [2] on elastic pp scattering at 13 TeV. We showed in Fig. 2 that a satisfactory description of the t distribution (and, in Fig. 4, the TOTEM measurement of $\rho \equiv \text{Re}A/\text{Im}A$ [1]) can be obtained in the framework of a two-channel eikonal model, even without the inclusion of an odd-signature (Odderon) contribution. However, the small addition of a QCD Odderon contribution may slightly improve the agreement with the data, especially for the ρ ratio. We emphasized, in Section 7, that if the value of the imaginary part of the amplitude at some impact parameter b calculated from the 13 TeV data were to exceed the black disc limit this would be a strong argument in favour of a large odd-signature contribution. It is impossible to get

¹⁰ Formally such a large value of σ_{odd} in a limited energy interval which includes 13 TeV does not violate unitarity. However, asymptotically as $s \rightarrow \infty$ the ratio $\text{Re}A/\text{Im}A$ must tend to 0, as was shown in [28].

¹¹ Recall that within our approach we are unable to reproduce such a large $\text{Im}A(0) > 1$ and the model would prefer a smaller value of σ_{tot} of about 105 mb at 13 TeV.

¹² Note however, that the ATLAS-ALFA data at $\sqrt{s} = 7$ and 8 GeV [38,39] are a bit below the black disk limit and all the previous results are consistent with $\text{Im}A(0) \leq 1$ within the error bars.

$\text{Im}A(b) > 1$ without a large odd signature term (much larger than that expected from the perturbative QCD Odderon).

On other hand when we improved the calculation of $\text{Im}A(b)$, for the precise TOTEM 13 TeV data accounting in (27) for the contribution from the large $|t|$ region (after the diffractive dip) where the imaginary part of the amplitude changes sign, we find

$$\text{Im}A(b=0) = 1.026. \quad (31)$$

Within the normalization error of about 3% this is consistent with the ‘black disk limit’ $\text{Im}A(b) \leq 1$.

We emphasize that actually the main analysis of this paper was the description of all the diffractive data obtained for pp (and $p\bar{p}$) collisions (σ_{tot} , $d\sigma_{\text{el}}/dt$, ρ , $\sigma_{\text{lowM}}^{\text{D}}$) over a wide range of collider energies (from $\sqrt{s} = 0.0625$ to 13 TeV) and a large interval of $-t$ from 0 up to 1 GeV^2 , in terms of a two-channel eikonal model. In this ‘global’ analysis, an overall satisfactory description of the data could be achieved either without, or with the inclusion of a small contribution from, a QCD Odderon.

Note that the two-channel ‘global’ description depends crucially on the experimental information on low-mass proton dissociation, $\sigma_{\text{lowM}}^{\text{D}}$. The discussion in Section 3 has required us to increase the values of $\sigma_{\text{lowM}}^{\text{D}}$ as compared to the values fitted in our earlier analyses. The consequence is that we have a flatter energy dependence of the total cross section – we slightly overestimate the measured value of σ_{tot} at 62.5 GeV and underestimate the value measured at 13 TeV. That is, the *overall* description prefers a lower value of $\sigma_{\text{tot}} \sim 105 \text{ mb}$ at 13 TeV, instead of the measured value $\sigma_{\text{tot}} \sim 110.6 \pm 3.4 \text{ mb}$ quoted by TOTEM [1]. We await more precise experimental knowledge of $\sigma_{\text{lowM}}^{\text{D}}$ and further measurements of ρ and σ_{tot} .

Acknowledgements

MGR thank the IPPP at the University of Durham for hospitality. VAK acknowledges support from a Royal Society of Edinburgh Auber award.

References

- [1] TOTEM collaboration, G. Antchev, et al., *First determination of the ρ parameter at $\sqrt{s} = 13 \text{ TeV}$ – probing the existence of a colourless three-gluon bound state*, CERN-EP-2017-335, Phys. Rev. (2017), submitted for publication.
- [2] F. Nemes, F. Ravera, Contributions on Behalf of the TOTEM Collaboration, Respectively, in the Proceedings of the 4th Elba Workshop on Forward Physics at LHC Energy, 24–26 May 2018, to be Published in a Special issue of Instruments; and at the 134th LHCC Meeting – Open Session, 30 May 2018.
- [3] M.L. Good, W.D. Walker, *Diffraction dissociation of beam particles*, Phys. Rev. 120 (1960) 1857.
- [4] V.A. Khoze, A.D. Martin, M.G. Ryskin, *Diffraction at the LHC*, Eur. Phys. J. C 73 (2013) 2503, arXiv:1306.2149.
- [5] TOTEM collaboration, G. Antchev, et al., *Measurement of proton–proton inelastic scattering cross-section at $\sqrt{s} = 7 \text{ TeV}$* , Europhys. Lett. 101 (2013) 21003.
- [6] A.B. Kaydalov, *High-energy particle production and regge cuts in elastic $n n$, πn , and $k n$ scattering*, Sov. J. Nucl. Phys. 13 (1971) 226.
- [7] A.B. Kaidalov, *Diffractive production mechanisms*, Phys. Rep. 50 (1979) 157.
- [8] L. Baksay, et al., *Observation of diffraction excitation in $pp \rightarrow (p\pi^+\pi^-) x$ at the CERN – ISR and tests of limiting fragmentation*, Phys. Lett. B 53 (1975) 484.
- [9] R. Webb, et al., *Production of nucleon resonances by single diffraction dissociation at the CERN ISR*, Phys. Lett. B 55 (1975) 331.

- [10] L. Baksay, et al., Diffraction dissociation in the reaction $pp \rightarrow \Lambda K^+ p$ at the CERN ISR, Phys. Lett. B 61 (1976) 405.
- [11] H. de Kerret, et al., Experimental results on diffractive one pion production at the CERN ISR, Phys. Lett. B 63 (1976) 477.
- [12] G.C. Mantovani, M. Cavalli-Sforza, C. Conta, M. Fraternali, G. Goggi, F. Pastore, et al., First results on diffraction dissociation of neutrons at the ISR, Phys. Lett. B 64 (1976) 471.
- [13] ATLAS collaboration, M. Aaboud, et al., Measurement of the inelastic proton–proton cross section at $\sqrt{s} = 13$ TeV with the ATLAS detector at the LHC, Phys. Rev. Lett. 117 (2016) 182002, arXiv:1606.02625.
- [14] CMS collaboration, A.M. Sirunyan, et al., Measurement of the inelastic proton–proton cross section at $\sqrt{s} = 13$ TeV, arXiv:1802.02613.
- [15] TOTEM collaboration, G. Antchev, et al., First measurement of elastic, inelastic and total cross-section at $\sqrt{s} = 13$ TeV by TOTEM and overview of cross-section data at LHC energies, arXiv:1712.06153.
- [16] ATLAS collaboration, G. Aad, et al., Rapidity gap cross sections measured with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV, Eur. Phys. J. C 72 (2012) 1926, arXiv:1201.2808.
- [17] M.G. Ryskin, A.D. Martin, V.A. Khoze, High-energy strong interactions: from 'hard' to 'soft', Eur. Phys. J. C 71 (2011) 1617, arXiv:1102.2844.
- [18] COMPETE collaboration, J.R. Cudell, V.V. Ezhela, P. Gauron, K. Kang, Yu.V. Kuyanov, S.B. Lugovsky, et al., Benchmarks for the forward observables at RHIC, the Tevatron Run II and the LHC, Phys. Rev. Lett. 89 (2002) 201801, arXiv:hep-ph/0206172.
- [19] Particle Data Group collaboration, C. Patrignani, et al., Review of particle physics, Chin. Phys. C 40 (2016) 590.
- [20] J. Kwiecinski, M. Praszalowicz, Three gluon integral equation and odd c singlet regge singularities in QCD, Phys. Lett. B 94 (1980) 413.
- [21] J. Bartels, High-energy behavior in a nonabelian gauge theory (II), Nucl. Phys. B 175 (1980) 365.
- [22] J. Bartels, L.N. Lipatov, G.P. Vacca, A new Odderon solution in perturbative QCD, Phys. Lett. B 477 (2000) 178, arXiv:hep-ph/9912423.
- [23] M.G. Ryskin, Odderon and polarization phenomena in QCD, Sov. J. Nucl. Phys. 46 (1987) 337.
- [24] M.A. Braun, Odderon and QCD, arXiv:hep-ph/9805394.
- [25] C. Ewerz, The Odderon: theoretical status and experimental tests, in: 11th International Conference on Elastic and Diffractive Scattering: Towards High Energy Frontiers: The 20th Anniversary of the Blois Workshops, 17th Rencontre de Blois (EDS 05) Chateau de Blois, Blois, France, May 15–20, 2005, 2005, arXiv:hep-ph/0511196.
- [26] M.M. Block, Hadronic forward scattering: predictions for the large hadron collider and cosmic rays, Phys. Rep. 436 (2006) 71, arXiv:hep-ph/0606215.
- [27] J. Finkelstein, H.M. Fried, K. Kang, C.I. Tan, Forward scattering at collider energies and Eikonal unitarization of Odderon, Phys. Lett. B 232 (1989) 257.
- [28] V.A. Khoze, A.D. Martin, M.G. Ryskin, Black disk, maximal Odderon and unitarity, Phys. Lett. B 780 (2018) 352, arXiv:1801.07065.
- [29] TOTEM collaboration, G. Antchev, et al., Measurement of elastic pp scattering at $\sqrt{s} = 8$ TeV in the Coulomb-nuclear interference region: determination of the ρ -parameter and the total cross-section, Eur. Phys. J. C 76 (2016) 661, arXiv:1610.00603.
- [30] E710 collaboration, N.A. Amos, et al., Measurement of ρ , the ratio of the real to imaginary part of the $\bar{p}p$ forward elastic scattering amplitude, at $\sqrt{s} = 1.8$ -TeV, Phys. Rev. Lett. 68 (1992) 2433.
- [31] UA4/2 collaboration, C. Augier, et al., A Precise measurement of the real part of the elastic scattering amplitude at the S anti-p p S, Phys. Lett. B 316 (1993) 448.
- [32] A.A. Anselm, V.N. Gribov, Zero pion mass limit in interactions at very high-energies, Phys. Lett. B 40 (1972) 487.
- [33] V.A. Khoze, A.D. Martin, M.G. Ryskin, Soft diffraction and the elastic slope at Tevatron and LHC energies: a MultiPomeron approach, Eur. Phys. J. C 18 (2000) 167, arXiv:hep-ph/0007359.
- [34] V.A. Khoze, A.D. Martin, M.G. Ryskin, t dependence of the slope of the high energy elastic pp cross section, J. Phys. G 42 (2015) 025003, arXiv:1410.0508.
- [35] A. Donnachie, P.V. Landshoff, Total cross-sections, Phys. Lett. B 296 (1992) 227, arXiv:hep-ph/9209205.
- [36] TOTEM collaboration, G. Antchev, et al., Luminosity-independent measurement of the proton–proton total cross section at $\sqrt{s} = 8$ TeV, Phys. Rev. Lett. 111 (2013) 012001.
- [37] F.J. Nemes, Elastic and total cross-section measurements by TOTEM: past and future, PoS DIS2017 (2018) 059.
- [38] ATLAS collaboration, G. Aad, et al., Measurement of the total cross section from elastic scattering in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector, Nucl. Phys. B 889 (2014) 486, arXiv:1408.5778.
- [39] ATLAS collaboration, M. Aaboud, et al., Measurement of the total cross section from elastic scattering in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector, Phys. Lett. B 761 (2016) 158, arXiv:1607.06605.
- [40] A. Alkin, E. Martynov, O. Kovalenko, S.M. Troshin, Impact-parameter analysis of TOTEM data at the LHC: black disk limit exceeded, Phys. Rev. D 89 (2014) 091501, arXiv:1403.8036.