

## Durham Research Online

---

### Deposited in DRO:

13 February 2019

### Version of attached file:

Accepted Version

### Peer-review status of attached file:

Peer-reviewed

### Citation for published item:

Hillier, Matthew and Sutton, Louise and James, Lewis and Mojtahedi, Dara and Keay, Nicola and Hind, Karen (2019) 'High prevalence and magnitude of rapid weight loss in mixed martial arts athletes.', *International journal of sports nutrition and exercise metabolism.*, 29 (5). pp. 512-517.

### Further information on publisher's website:

<https://doi.org/10.1123/ijsnem.2018-0393>

### Publisher's copyright statement:

Manuscript has been published in a journal issue: Accepted author manuscript version reprinted, by permission, from Hillier, Matthew, Sutton, Louise, James, Lewis, Mojtahedi, Dara, Keay, Nicola Hind, Karen (2019). High Prevalence and Magnitude of Rapid Weight Loss in Mixed Martial Arts Athletes. *International Journal of Sports Nutrition and Exercise Metabolism* 29(5): 512-517, <https://doi.org/10.1123/ijsnem.2018-0393>. © Human Kinetics, Inc.

### Additional information:

## Use policy

---

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

## ACCEPTED VERSION

*Int J Sport Nutr Exerc Metab*

### Original Investigation

#### High Prevalence and Magnitude of Rapid Weight Loss in Mixed Martial Arts Athletes

M Hillier, L Sutton, L James, D Motjhed, N Keay and K Hind.

### ABSTRACT

The practice of rapid weight loss in mixed martial arts (MMA) is an increasing concern but data remains scarce. The aim of this study was to investigate the prevalence, magnitude, methods and influencers of rapid weight loss in professional and amateur MMA athletes. MMA athletes (n=314; 287 men, 27 women) across nine weight categories (strawweight to heavyweight), completed a validated questionnaire adapted for this sport. Sex-specific data were analysed, and sub-group comparisons were made between athletes competing at professional and amateur levels. Most athletes purposefully reduced body weight for competition (men: 97.2%; women: 100%). The magnitude of rapid weight loss in one week prior to weigh-in was significantly greater for professional athletes compared to those competing at amateur level (men: 5.9% v 4.2%; women: 5.0% v 2.1% of body weight;  $p<0.05$ ). In the 24 h preceding weigh-in, the magnitude of rapid weight loss was greater at professional than amateur level in men (3.7% v 2.5% of body weight;  $p<0.05$ ). Most athletes 'always' or 'sometimes' used water loading (72.9%), restricting fluid intake (71.3%) and sweat suits (55.4%) for rapid weight loss. Coaches were cited as the primary source of influence on rapid weight loss practices (men: 29.3%, women: 48.1%). There is a high reported prevalence of rapid weight loss in MMA, at professional and amateur levels. Our findings, constituting

the largest enquiry to date, call for urgent action from MMA organisations to safeguard the health and wellbeing of athletes competing in this sport.

**Key words:** martial arts; combat sport; dieting; health risks; rapid weight loss; weight loss.

## Introduction

Mixed martial arts (MMA) is a combat sport, with bouts defined by weight divisions (Reale et al., 2017a) with the aim of endorsing balanced and stimulating matches whilst reducing potential injuries that may result from substitutional differences in weight (Mendes et al., 2013). For the athlete, the process of 'making weight' is imperative, because failure to make weight results in bout cancellation or deduction from the athlete's payment.

MMA athletes engage in gradual and rapid weight loss (RWL) prior to competition and then re-gain weight post weigh-in (Jetton et al., 2013; Matthews & Nicholas, 2017; Coswig et al., 2018). Gradual weight loss includes long-term energy restriction and exercise training. Energy and other nutrient deficits can lead to altered hormonal milieu, muscle atrophy, fatigue, altered bone metabolism, depressed immune system and decreased energy metabolism, all of which are particularly problematic for athletes (Zanker and Hind, 2007; Mountjoy et al., 2014). RWL describes the practice of reducing weight over a short period of time. RWL is an ongoing problem in combat sports, indicated by reports that athletes are using controversial and potentially harmful practices to achieve entry into the lowest weight class possible (Artioli et al., 2010a; Crighton et al., 2015). Given that RWL is detrimental to health and performance with risks to cognitive function (Franchini et al., 2012), and a higher risk of injury (Green et al., 2007), calls to action have been published (Crighton et al., 2015). At its worst, RWL has been implicated in the deaths of combat athletes in the days/hours prior to competition. Three collegiate wrestlers died from complications arising from RWL in

1997 (Centers for Disease Control, Prevention, 1998). The first death of an MMA athlete during RWL was reported in 2013 (Crighton et al., 2015). Since then, the number of reported fatalities during RWL have increased in martial arts with similarities to MMA (Matthews & Nicholas, 2017) including the death of a Scottish martial arts athlete prior to a Muay Thai fight in 2017.

Despite growing concerns, published data on the extent of RWL in MMA remains scant. Therefore, the purpose of this study was to provide robust data on the prevalence, magnitude, methods and influencers of gradual and rapid weight loss in professional and amateur MMA athletes.

## Methods

### *Participants*

Professional and amateur MMA athletes over the age of 18 years (strawweight to heavy-weight categories) were recruited from MMA gyms, events, competitions and websites. Participants were categorised by the level at which they competed at the time of completing the questionnaire. Professional level athletes participated under the rule set of Unified Rules of Mixed Martial Arts (URMMA) and received payment to compete. Amateur level athletes did not receive payment for competing and participated under the rule sets of Amateur A, B or C; which have shorter rounds (3 x 3 min) and a more restricted rule set. Of the 318 athletes who replied to the study invitation, 314 were eligible to take part. One athlete was excluded for not specifying their weight category, whilst three athletes responding online were under 18 years and were also excluded from the study. Additionally, 24 athletes submitted incomplete answers to questions on usual weight loss practices. Therefore, the final sample size was 290 (264 men, 26 women) for data relating to weight loss magnitude and 314 (287 men, 27 women) for data relating to the method of weight loss and key influencers. The study

was approved by the University Research Ethics Committee, was conducted in accordance with the Declaration of Helsinki (2013) and all participants provided informed consent.

#### *Validated questionnaire*

With approval from the authors (Artioli et al., 2010b) the validated Rapid Weight Loss Questionnaire (RWLQ) was adapted for the current study to ensure appropriateness to MMA. The questionnaire was divided into 3 sections with a total of 21 questions, incorporating a Likert Scale using a range of 1-5 to measure the prevalence and magnitude of dieting methods (1-Always, 2-Sometimes, 3-Almost never, 4-I don't use anymore, 5-Never used). The questions covered the level and frequency of competition, training, athletic achievements, weight history, diet and RWL. The adapted questionnaire was piloted through completion by local MMA athletes, and feedback was provided on the suitability and clarity of the questions. Following minor amendments arising from the piloting process, the final questionnaire was deemed appropriate for the study objectives and the target population (Appendix).

#### *Statistical analysis*

Statistical analyses were completed using SPSS (Version 22.0, IBM Corp., Armonk, NY). MANCOVA analysis was used to compare levels of weight loss across the independent variables (sex & competition level), whilst controlling for age. A series of Chi-square analyses were conducted to test the relationships between the independent variables, weight loss methods used and key influencers for weight loss. Standardised residual values were observed to identify the responses that were over or under represented within the independent variable groups. Standardised residual values of +/- 1.9 were identified as being significant due to corresponding to an alpha level of  $p < 0.05$ . Cohen's d effect size was calculated where

appropriate and interpreted in accordance to Cohen, (1988). Data are presented as mean (SD), with range where appropriate.

## Results

The descriptive results are given in Tables 1 and 2.

### *Prevalence and magnitude of weight loss and RWL*

One hundred percent of women and 97.2% of men athletes reported engaging in purposeful weight loss. For the purpose of this study, total weight loss encompasses both gradual weight loss and late RWL. The average reported total weight lost was  $7.3 \pm 3.4$  kg (range 0-18.0 kg) in men and  $5.1 \pm 2.7$  kg (range 2–11 kg) in women, representing  $9.0 \pm 3.9\%$  (range 0-20.5%) and  $7.7 \pm 3.3\%$  (range 2.5-13.7%) of body weight in men and women, respectively. The most amount of weight lost for a fight (not specific to RWL) was  $10.2 \pm 4.6$  kg (0-25 kg) in men and  $7.0 \pm 3.6$  kg (2-14 kg) in women, representing  $12.5 \pm 5.0\%$  (0-25.9%) and  $10.7 \pm 4.5\%$  (2.8-20.6%) of body weight in men and women, respectively.

In men, 96.1% (professional: 95.2%, amateur: 96.9%) and 90.2% (professional: 93.4%, amateur: 87.2%) reported engaging in RWL in the one week and 24 hours prior to weigh-in, respectively. Men reported losing an average of  $4.1 \pm 2.5$  kg (0-13 kg) or  $5.0 \pm 3.1\%$  (0-14.0%) of body weight one week before weigh-in and  $2.5 \pm 1.8$  kg (0-10 kg) or  $3.1 \pm 2.2\%$  (0-10.8%) of body weight in the 24 hours before weigh-in. In women, all and 78.3% (professional: 88.9%; amateur: 71.4%) athletes reported engaging in RWL in the one week and 24 hours prior to weigh-in, respectively. Women reported losing an average of  $2.5 \pm 1.4$  kg (1-7 kg) or  $3.8 \pm 2.0\%$  (0.9-9.6%) of body weight one week before weigh-in and  $1.5 \pm 1.3$  kg (0-5 kg) or  $2.3 \pm 1.9\%$  (0-6.9 %) of body weight in the 24 hours before weigh-in. One

third ( $n = 9$ ) of women athletes reported missing 3 or more consecutive menstrual cycles within the last 12 months.

To determine whether sex or competition level effected reported weight lost, controlling for age, %total weight loss, % weight lost in the final week, and % weight lost in the final 24 hours were tested in a 2 (men v women) x 2 (amateur v professional) MANCOVA. There was a significant main effect for competition level,  $F(3,280) = 3.64$ ;  $p < 0.05$ , using the *Wilks' lambda* criterion. Total weight loss was greater in professional athletes ( $10.0 \pm 3.9\%$  v  $7.9 \pm 6.6\%$ ;  $F(1,282) = 5.5$ ;  $p < 0.05$ ;  $d = -.37$ ; small effect). Professional athletes also lost more weight in the final week ( $5.9 \pm 2.9\%$  v  $4.1 \pm 2.5\%$ ;  $F(1,282) = 9.97$ ;  $p < 0.05$ ;  $d = -.69$ ; medium effect) and 24 h ( $3.7 \pm 2.1\%$  v  $2.5 \pm 2.0\%$ ;  $F(1,282) = 9.39$ ;  $p < 0.05$ ;  $d = -.62$ ; medium effect). There were no interaction effects with sex in any models.

#### *Methods of weight loss and RWL*

Methods of weight loss are reported in Table 3. The most common method was gradual dieting, with 81.2% reporting to always use this. Approximately half of athletes (50.6%) reported 'always' using water loading. The least common method was self-induced vomiting, with 94.6% of athletes claiming to never use this method.

Chi-square analysis indicated women athletes (44.4%) were significantly more likely to have 'never used' water loading methods, in comparison to men athletes (15.7%). With regard to restricting fluids, professionals (57.4%) were significantly over represented for the response 'Always'. For water loading, professionals (4.7%) were over represented for the response 'I don't use anymore' and amateur athletes (25.3%) were over represented for the response 'Never use'. For sauna use, amateur athletes (31.3%) were over represented for the

response '*Never use*'. For laxative use, amateur athletes (94.9%) were significantly over represented for the response '*Never use*'. Professionals were over represented for the '*Always*' and '*Sometimes*' responses (7.4% and 8.9%, respectively). With regard to diuretics use, amateur athletes (81.9%) were significantly over represented for the response '*Never use*' and professionals were over represented for the '*Always*' response (14.9%). There was a significant difference in diet pill use, with professional athletes (6.8%) more represented than amateur athletes (2.4%) for the response '*Always*', however the residual score did not correspond to an alpha level of 0.05. For sweat suit use, professionals (39.2%) were significantly over represented for the response '*Always*'. Professional athletes (7.4%) were significantly over represented for the '*I don't use anymore*' response. Amateur athletes were also significantly over represented (39.8%) for the '*Never use*' response.

A chi-square test of independence found a significant difference in the use of gradual dieting between the different weight classes.  $X^2 (32, N = 314) = 67.58, p < .001$ . Inspection of the standardized residual scores indicated that Heavyweight respondents were overrepresented in stating that they '*Never used*' gradual dieting (corresponded to a critical value of .01). A chi-square test of independence found a significant difference in the use of fluid restrictions between the different weight classes.  $X^2 (32, N = 314) = 59.24, p < .01$ . Inspection of the standardized residual scores indicated that Heavyweight respondents were overrepresented in stating that they '*Never used*' fluid restriction (corresponded to a critical value of .01).

#### *Influences on weight loss practices*

There were significant sex differences in reported influencers of weight loss practices,  $\chi^2 (8, N = 314) = 26.45, p < 0.01$  (Table 4). There were no differences in reported influencers between professional and amateur athletes,  $\chi^2 (8, N = 314) = 14.12; p > 0.05$ .



171

172 **Discussion**

173 The purpose of this study was to investigate the prevalence, magnitude and influencers of  
 174 weight loss prior to competition in professional and amateur MMA athletes. In this largest  
 175 enquiry to date, the major finding was that most men and all women MMA athletes reported  
 176 engaging in gradual and rapid weight loss leading up to competition, with coaches being key  
 177 influencers of their weight loss practices. Our findings also indicate that a significant  
 178 proportion of athletes are using strategies that reduce body water stores (e.g. water loading,  
 179 fluid restriction, increasing sweat losses through exercise and/or heat exposure) as a primary  
 180 means of RWL both at professional and amateur levels.

181 In the present study, the prevalence of weight loss in MMA athletes ranged from 97.2 to  
 182 100%. With RWL in the week or final 24 hours before weigh-in ranging from (90.2-100%).  
 183 This is similar or higher than rates reported for other combat sports athletes (Kinningham et  
 184 al., 2001; Jetton et al., 2013; Horswill, 2009; Artioli et al., 2010a; Brito et al., 2012; Barley  
 185 et al., 2017). In judo, 62-90% of athletes report to engage in RWL (Artioli et al., 2010a; Brito  
 186 et al., 2012; Barley et al., 2017) in wrestling, 60-97% (Kinningham et al., 2001; Barley et al.,  
 187 2017), in Brazilian jiu-jitsu, 56.8-88% (Bruto et al., 2012; Barley et al., 2017), in Muay  
 188 Thai/kickboxing, 94% (Barley et al., 2017) and in taekwondo, 63.3-85% (Bruto et al., 2012;  
 189 Barley et al., 2017). The higher prevalence in the current study of MMA athletes may reflect  
 190 the rapid growth in popularity of the sport and prize money, but might also reflect the broad  
 191 weight categories and/or the extended time between weigh-in and competition for most  
 192 events (usually 24-36 hours) compared to that of judo (2.5-5 hours) and amateur boxing (0-8  
 193 hours). Whilst most previous studies of RWL in combat sports have focused on men, we  
 194 found that the prevalence of RWL in women MMA athletes is equivalent to men. While there

were some significant differences between amateur and professional level men athletes for weight loss in the week and 24 h before weigh-in and total weight loss, the absolute differences were small. Together the findings infer that RWL in the final week and 24 hours before weigh-in is not exclusive to professional men athletes - rather it appears to be a significant problem across the board.

The reported range of weight loss varied in the current study. A number of athletes reported losing significant amounts of weight, including one professional (male), who reported losing as much as 25.8% of body weight. The reported average amounts of RWL for professional MMA athletes were 5.9% (men) and 4.2% (women) with the most extreme being 14% of body weight across the week leading up to competition and 3.7% (men) and 2.5% (women) with the most extreme being 10.7% of body weight over the 24 hours before weigh-in. These percentages are similar to those reported by smaller studies (Andreato, et al., 2014; Crighton et al., 2015; Coswig et al., 2015; Matthews & Nicholas, 2017) and similar or higher than rates reported for other combat sports such as Brazilian jiu-jitsu (4%) (Brito et al., 2012), taekwondo (4%) (Brito et al., 2012), wrestling (5%) (16) and kickboxing (4%) (Boguszewski et al., 2010). Our findings are concerning especially given reports of fatalities in combat athletes reducing similar amounts of body weight (Crighton et al., 2015; Centers for Disease Control, Prevention, 1998).

The methods of weight loss reported by athletes were predominantly related to manipulation of body water and inducing energy deficit through increased exercise and reduced dietary intake. The main health risks associated with RWL are mainly cardiocirculatory complications arising from severe dehydration/hyperthermia. A recent study of MMA athletes ( $N = 7$ ) engaging in RWL leading up to competition reported raised urine osmolality indicative of significant/severe dehydration in athletes at weigh-in (Matthews & Nicholas, 2017). Jetton et al. (2013) reported similar findings, with significant dehydration

present immediately before competition following average RWL of ~4% of body weight. Whilst acute dehydration (James et al., 2017) and energy restriction (James et al. 2015) might impair performance, the impairments in thermoregulatory and cardiovascular function caused by hypovolaemia/hyperosmolality (Cheuvront and Kenefick, 2014) are of particular concern, and may exacerbate any negative health effects related to heat exposure used to facilitate weight loss (Casa et al., 2015). This is particularly relevant given that heat exposure has been implicated in a number of the deaths reported in those attempting to make weight (Centers for Disease Control, Prevention, 1998). Furthermore, the severe energy restriction many athletes use in the days prior to weigh-in promotes electrolyte imbalance and plasma volume reduction (James and Shirreffs, 2013; James et al. 2015), that may further exacerbate these heat/cardiovascular-related issues.

Water loading was the second most frequently reported method of RWL for men and women, at both professional and amateur levels. Water loading involves consuming significant volumes of water (~10 litres of water/day for 3-5 days) followed by complete fluid restriction over the 24 hours before weigh-in, a strategy that appears to result in a small increase in weight loss (Reale et al, 2017b). The popularity of water loading is also concerning given the risk of dilutional hyponatraemia if the large volume of water ingested is not spaced appropriately over the day (Mohan et al., 2013). High volume intakes over a short time frame (e.g. >10 litres in 6 hours) has resulted in fatalities (Mohan et al., 2013), but progressive water loading over a 6 day period leading to competition may attenuate the risk of hyponatraemia (Adrogué and Madias, 2000) More research into water loading is needed before recommendations can be made on risks.

Energy deficit through long-term weight cycling in weight class sports is known to effect metabolic, endocrine, and immune function (Turocy et al., 2011), although there are also acute negative metabolic health consequences associated with short-term RWL (Kasper

et al., 2018; Green et al., 2007). Raised cortisol in the short term, will disrupt other endocrine networks and diurnal variation of this hormone. In the longer term, raised cortisol would contribute to adverse effects on bone. In women, energy deficit can suppress menstrual function and in the current study, one third of women athletes reported amenorrhea in the last 12 months. It is possible that this figure is an underestimation because women using oral contraceptives were not excluded from the study. Low energy availability and disruption of menstrual functioning also negatively impacts bone metabolism and cardiovascular function (Artioli et al., 2010a; Mountjoy et al., 2014). Adverse effects on the cardiovascular system include reduction of endothelial reactivity and disruption of lipid profile due to low oestrogen levels, as seen in post-menopausal women where there is marked increase in incidence of cardiovascular disease. Additionally, there is growing concern that RWL might increase the risk for brain injury in MMA compared to other combat sports, given that head trauma can still occur after an athlete has fallen unconscious/fainted (Crighton et al., 2015). In terms of effects on performance, energy deficit can impair judgement, concentration and coordination (Mountjoy et al., 2014).

The key influencer of RWL for both professional and amateur MMA athletes, and for both sexes, was the coach. Professional women athletes were more likely to report consulting a dietician than their men counterparts. The use of the internet as the primary source of information on rapid weight loss was higher in men than in women MMA athletes. The lack of clinical input or support is concerning, as there can be individual variability in physiological response. If any underlying medical conditions have not been excluded, then extreme practices of inducing RWL through energy deficit and/or manipulation of intra and extracellular water could lead to increased risk of serious health consequences.

The authors acknowledge that while this is the largest enquiry of its kind to date, the self-report nature of the study is a limitation. The questionnaire was piloted prior to data

collection to ensure comprehension and whilst responses were anonymously sought, we cannot guarantee full honesty in disclosure. Aside from RWL (weight loss in the 1 week before weigh-in), questions pertaining to gradual weight loss were not time bound.

In conclusion, our study indicates a high prevalence of RWL in MMA regardless of competition level and sex, with the deleterious practice of water loading as one of the most commonly reported methods for weight loss. Our findings, constituting the largest enquiry to date, call for urgent action from MMA organisations to safeguard the health and wellbeing of athletes competing in this sport.

### **Acknowledgements**

The study was designed by MH, KH, LS, and LJ; data were collected and analysed by MH, DM, KH, LJ; data interpretation and manuscript preparation were undertaken by MH, KH, LS, LJ, DM, and NK. All authors inputted to and approved the final version of the paper.

### **Competing interests**

There are no competing influences to declare.

### **Funding**

This study was not supported by external funding.

### **References**

Adrogué, H. J., & Madias, N. E. (2000). Hyponatremia. *New England Journal of Medicine*, 342(21), 1581-1589.

- 294 Andreato, L., Andreato, T., Ferreira da Silva Santos, J., Del Conti Esteves, J., Franzói de  
 295 Moraes, S., & Franchini, E. (2014). Weight loss in mixed martial arts athletes. *J Combat*  
 296 *Sports Martial Arts*, 5(2), 117-23.
- 297
- 298 Artioli, G. G., Gualano, B., Franchini, E., Scagliusi, F. B., Takesian, M., Fuchs, M., &  
 299 Antonio Herbert Lancha, J. R. (2010). Prevalence, magnitude, and methods of rapid weight  
 300 loss among judo competitors. *Medicine & Science in Sports & Exercise*, 42(3), 436-442.
- 301
- 302 Artioli, G. G., Scagliusi, F., Kashiwagura, D., Franchini, E., Gualano, B., & Junior, A. L.  
 303 (2010). Development, validity and reliability of a questionnaire designed to evaluate rapid  
 304 weight loss patterns in judo players. *Scandinavian Journal of Medicine & Science in*  
 305 *Sports*, 20(1), e177-e187.
- 306
- 307 Barley, O. R., Chapman, D. W., & Abbiss, C. R. (2017). Weight Loss Strategies in Combat  
 308 Sports and Concerning Habits in Mixed Martial Arts. *International journal of sports*  
 309 *physiology and performance*, 1-24.
- 310
- 311 Boguszewski, D., & Kwapisz, E. (2010). Sports massage and local cryotherapy as a way to  
 312 reduce negative effects of rapid weight loss among kickboxing contestants. *Archives of*  
 313 *Budo*, 6(1), 1-7.
- 314
- 315 Brito, C. J., Roas, A. F. C. M., Brito, I. S. S., Marins, J. C. B., Córdova, C., & Franchini, E.  
 316 (2012). Methods of body-mass reduction by combat sport athletes. *International journal of*  
 317 *sport nutrition and exercise metabolism*, 22(2), 89-97.
- 318

Casa, D. J., DeMartini, J. K., Bergeron, M. F., Csillan, D., Eichner, E. R., Lopez, R. M., ... & Yeargin, S. W. (2015). National Athletic Trainers' Association position statement: exertional heat illnesses. *Journal of athletic training*, 50(9), 986-1000.

Centers for Disease Control, & Prevention (US). Epidemiology Program Office. (1998). *MMWR.: Recommendations and Reports* (Vol. 47). US Department of Health and

Cheuvront, S. N., & Kenefick, R. W. (2014). Dehydration: physiology, assessment, and performance effects. *Compr Physiol*, 4(1), 257-285.

Cohen, J. (1988). Statistical power analysis for the behavioral sciences. Hillsdale, NJ: Lawrence Erlbaum (2nd ed.)

Coswig, V. S., Fukuda, D. H., & Del Vecchio, F. B. (2015). Rapid weight loss elicits harmful biochemical and hormonal responses in mixed martial arts athletes. *International journal of sport nutrition and exercise metabolism*, 25(5), 480-486.

Coswig, V. S., Miarka, B., Pires, D. A., da Silva, L. M., Bartel, C., & Del Vecchio, F. B. (2018). Weight Regain, But Not Weight Loss, Is Related to Competitive Success in Real-life Mixed Martial Arts Competition. *International journal of sport nutrition and exercise metabolism*, 1-26.

Crighton, B., Close, G. L., & Morton, J. P. (2016). Alarming weight cutting behaviours in mixed martial arts: a cause for concern and a call for action.

Franchini, E., Brito, C. J., & Artioli, G. G. (2012). Weight loss in combat sports: physiological, psychological and performance effects. *Journal of the international society of sports nutrition*, 9(1), 52.

Green, C. M., Petrou, M. J., Fogarty-Hover, M. L., & Rolf, C. G. (2007). Injuries among judokas during competition. *Scandinavian journal of medicine & science in sports*, 17(3), 205-210.

James, L. J., & Shirreffs, S. M. (2013) Fluid and electrolyte balance during 24-hour fluid and/or energy restriction. *International Journal of Sport Nutrition and Exercise Metabolism*, 23(6), 545-553.

James, L. J., Mears, S. A., & Shirreffs, S. M. (2015). Electrolyte supplementation during severe energy restriction increases exercise capacity in the heat. *European Journal of Applied Physiology*, 115(12), 2621-2629.

James, L. J., Moss, J., Henry, J., Papadopoulou, C., & Mears, S. A. (2017). Hypohydration impairs endurance performance: a blinded study. *Physiological reports*, 5(12), e13315.

Jetton, A. M., Lawrence, M. M., Meucci, M., Haines, T. L., Collier, S. R., Morris, D. M., & Utter, A. C. (2013). Dehydration and acute weight gain in mixed martial arts fighters before competition. *The Journal of Strength & Conditioning Research*, 27(5), 1322-1326.

Kasper, A. M., Crighton, B., Langan-Evans, C., Riley, P., Sharma, A., Close, G. L., & Morton, J. P. (2018). Case Study: Extreme Weight Making Causes Relative Energy



Deficiency, Dehydration and Acute Kidney Injury in a Men Mixed Martial Arts  
Athlete. *International journal of sport nutrition and exercise metabolism*, 1-20.

Kinningham, R. B., & Gorenflo, D. W. (2001). Weight loss methods of high school  
wrestlers. *Medicine and science in sports and exercise*, 33(5), 810-813.

Matthews, J. J., & Nicholas, C. (2017). Extreme rapid weight loss and rapid weight gain  
observed in UK mixed martial arts athletes preparing for competition. *International journal  
of sport nutrition and exercise metabolism*, 27(2), 122-129.

Mendes, S. H., Tritto, A. C., Guilherme, J. P. L., Solis, M. Y., Vieira, D. E., Franchini, E., ...  
& Artioli, G. G. (2013). Effect of rapid weight loss on performance in combat sport men  
athletes: does adaptation to chronic weight cycling play a role?. *Br J Sports Med*, bjsports-  
2013.

Mohan, S., Gu, S., Parikh, A., & Radhakrishnan, J. (2013). Prevalence of hyponatremia and  
association with mortality: *The American journal of medicine*, 126(12), 1127-1137.

Mountjoy, M., Sundgot-Borgen, J., Burke, L., Carter, S., Constantini, N., Lebrun, C., ... &  
Ljungqvist, A. (2014). The IOC consensus statement: beyond the women athlete triad—  
Relative Energy Deficiency in Sport (RED-S). *Br J Sports Med*, 48(7), 491-497.

Reale, R., Slater, G., & Burke, L. M. (2017). Acute-weight-loss strategies for combat sports  
and applications to Olympic success. *International Journal of Sports Physiology and  
Performance*, 12(2), 142-151.

Reale, R., Slater, G., Cox, G. R., Dunican, I. C., & Burke, L. M. (2017). The Effect of Water Loading on Acute Weight Loss Following Fluid Restriction in Combat Sports Athletes. *International journal of sport nutrition and exercise metabolism*, 20(XX), 1-9.

Turocy, P. S., DePalma, B. F., Horswill, C. A., Laquale, K. M., Martin, T. J., Perry, A. C., ... & Utter, A. C. (2011). National athletic trainers' association position statement: safe weight loss and maintenance practices in sport and exercise. *Journal of athletic training*, 46(3), 322-336.

Zanker, C., & Hind, K. (2007). The effect of energy balance on endocrine function and bone health in youth. In *Optimizing Bone Mass and Strength* (Vol. 51, pp. 81-101). Karger Publishers.

**Table 1.** Mixed martial arts athlete demographics and weight classes (count),  $N = 314$ .

	Men, n = 287	Women, n = 27
Mean age, years (SD)	27.2 (5.2)	28.8 (5.2)
Mean body mass, kg (SD)	81.8 (12.1)	63.2 (8.1)
Weight class (kg; range from previous category)	0	9
Straw (52.2)		
Fly (56.7; 4.5 kg)	19	4
Bantam (61.2; 4.5 kg)	40	10
Feather (65.8; 4.6 kg)	32	2

---

Light (70.3; 4.5 kg)	80	2
Welter (77.1; 6.8 kg)	52	0
Middle (83.9; 6.8 kg)	38	0
Light-heavy (93.0; 9.1 kg)	16	0
Heavy (120.2; 27.2 kg)	10	0

---

409

410

411

412

413

414

415

416

417

418

419

420 **Table 2:** Reported magnitude (mean  $\pm$  SD, range) of weight cutting in mixed martial arts ( $n = 290$ )

	Men		Women	
	<i>Professional, n = 126</i>	<i>Amateur, n = 138</i>	<i>Professional, n = 9</i>	<i>Amateur, n = 17</i>
Age, years	29.3 (5.1)	25.2 (4.5)	31.9 (2.7)	27 (5.5)
Walking weight, kg	83.6 (12.6)	80.3 (11.2)	62.1 (12.6)	63.9 (8.5)
Average total weight loss, kg	8.4 (3.5)	6.4 (3.1)	5.5 (2.7)	4.8 (2.7)
Average total weight loss, %	10.1 (3.9)	8 (3.6)	8.6 (3.4)	7.2 (3.2)
Weight loss in 1 week before weigh-in, kg	5 (2.6)	3.4 (2.1)	3.2 (1.7)	2.1 (1.2)
Weight loss in 1 week before weigh-in, %	5.9 (2.9)	4.2 (2.5)	5 (2.2)	3.1 (1.6)
Weight loss in 24 h before weigh-in, kg	3.1 (1.9)	2 (1.6)	2.1 (1.4)	1.2 (1.6)
Weight loss in 24 h before weigh-in, %	3.7 (2.1)	2.5 (1.9)	3.2 (1.9)	1.8 (1.6)

421

422 **Table 3.** Methods of weight loss in mixed martial arts (% of group;  $N = 314$ )

	Response (percentage)					<i>Chi-square</i> ( $\chi^2$ )	
	Always	Sometimes	Almost never	I don't use anymore	Never used	Sex <sup>a</sup>	Competition level <sup>a</sup>
Gradual dieting	81.2	12.1	2.5	0.6	3.5	2.99	8.44
Skipping meals	8.6	27.1	19.4	6.7	38.2	7.94	6.62
Fat burners	8	14	4.8	6.1	67.2	2.25	3.47
Diet pills	4.5	8	3.5	3.5	80.6	4.21	11*
Laxatives	3.8	13.1	7.6	3.8	71.7	3.14	32.84***
Increased exercise	51	25.5	8.6	1.6	13.4	.86	2.35
Water loading	50.6	22.3	6.7	2.2	18.2	15.49**	23.9***
Restricting fluid intake	46.8	24.5	10.2	2.2	16.2	2.69	15.16**
Sweat suit	29	26.4	12.1	4.1	28.3	1.78	33.68***
Saunas	25.5	35	11.5	4.8	23.2	7.21	16.41**
Fasting	20.1	19.7	15.6	6.7	37.9	2.99	6.53
Heated training rooms	15.3	26.4	17.2	2.2	38.9	4.12	7.69
Diuretics	8	10.2	7.6	5.4	68.8	1.53	33.54***
Vomiting	0.3	0.6	3.5	1	94.6	5.55	6.3

423 \* =  $p < .05$ , \*\* =  $p < .01$ , \*\*\* =  $p < .001$ . <sup>a</sup>:  $df = 4$ .

424 **Table 4.** Distribution of responses for influential sources on weight loss (percentage within group).

Influencer	Sex		Competitive level	
	Men	Women	Professional	Amateur
Training partner	16.7	3.7	17.6	13.9
Fellow MMA athlete	15.7	3.7	18.9	10.8
Doctor	0.7	0.0	0.7	0.6
Physical trainer	3.1	7.4	3.4	3.6
MMA coach	29.3	48.1	22.3	38.6
Parents	0.3	3.7	0.7	0.6
Dietician	10.1	29.6	14.2	9.6
Internet	15.0	3.7	12.2	15.7

425

