

Durham Research Online

Deposited in DRO:

02 July 2019

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Gaspers, Serge and Huang, Shenwei and Paulusma, Daniël (2019) 'Colouring square-free graphs without long induced paths.', *Journal of computer and system sciences.*, 106 . pp. 60-79.

Further information on publisher's website:

<https://doi.org/10.1016/j.jcss.2019.06.002>

Publisher's copyright statement:

© 2019 This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Accepted Manuscript

Colouring Square-Free Graphs without Long Induced Paths

Serge Gaspers, Shenwei Huang, Daniël Paulusma

PII: S0022-0000(18)30814-6
DOI: <https://doi.org/10.1016/j.jcss.2019.06.002>
Reference: YJCSS 3223

To appear in: *Journal of Computer and System Sciences*

Received date: 24 September 2018
Accepted date: 22 June 2019

Please cite this article in press as: S. Gaspers et al., Colouring Square-Free Graphs without Long Induced Paths, *J. Comput. Syst. Sci.* (2019), <https://doi.org/10.1016/j.jcss.2019.06.002>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Colouring Square-Free Graphs without Long Induced Paths*

Serge Gaspers^{†‡} Shenwei Huang[§] Daniël Paulusma[¶]

June 26, 2019

Abstract

The complexity of COLOURING is fully understood for H -free graphs, but there are still major complexity gaps if two induced subgraphs H_1 and H_2 are forbidden. Let H_1 be the s -vertex cycle C_s and H_2 be the t -vertex path P_t . We show that COLOURING is polynomial-time solvable for $s = 4$ and $t \leq 6$, strengthening several known results. Our main approach is to initiate a study into the boundedness of the clique-width of atoms (graphs with no clique cutset) of a hereditary graph class. As a complementary result we prove that COLOURING is NP-complete for $s = 4$ and $t \geq 9$, which is the first hardness result on COLOURING for (C_4, P_t) -free graphs. Combining our new results with known results leads to an almost complete dichotomy for COLOURING restricted to (C_s, P_t) -free graphs.

1 Introduction

Graph colouring has been a popular and extensively studied concept in computer science and mathematics since its introduction as a map colouring problem more than 150 years ago due to its many application areas crossing disciplinary boundaries and to its use as a benchmark problem in research into computational hardness. The corresponding decision problem, COLOURING, is to decide, for a given graph G and integer k , if G admits a k -colouring, that is, a mapping $c : V(G) \rightarrow \{1, \dots, k\}$ such that $c(u) \neq c(v)$ whenever $uv \in E(G)$. Unless $P = NP$, it is not possible to solve COLOURING in polynomial time for general graphs, not even if the number of colours is limited to 3 [45]. To get a better understanding of the borderline between tractable and intractable instances of COLOURING, it is natural to restrict the input to some special graph class. Hereditary graph classes, which are classes of graphs closed under vertex deletion, provide a unified framework for a large collection of well-known graph classes. It is readily seen that a graph class is hereditary if and only if it can be characterized by a unique set \mathcal{H} of minimal forbidden induced subgraphs. Graphs with no induced subgraph isomorphic to a graph in a set \mathcal{H} are called \mathcal{H} -free.

*An extended abstract of this paper appeared in the proceedings of STACS 2018 [25].

[†]School of Computer Science and Engineering, UNSW Sydney, Sydney 2052, Australia.

[‡]Decision Sciences Group, Data61, CSIRO, Sydney 2052, Australia.

[§]College of Computer Science, Nankai University, Tianjin 300350, China. The corresponding author (shenweihuang@nankai.edu.cn).

[¶]Department of Computer Science, Durham University, Durham DH13LE, UK.

Over the years, the study of COLOURING for hereditary graph classes has evolved into a deep area of research in theoretical computer science and discrete mathematics (see, for example, [8, 29, 38, 53]). One of the best-known results is the classical result of Grötschel, Lovász, and Schrijver [31], who showed that COLOURING is polynomial-time solvable for perfect graphs. Faster, even linear-time, algorithms are known for subclasses of perfect graphs, such as chordal graphs, bipartite graphs, interval graphs, and comparability graphs; see for example [29]. All these classes are characterized by *infinitely* many minimal forbidden induced subgraphs.

Král', Kratochvíl, Tuza, and Woeginger [43] initiated a systematic study into the computational complexity of COLOURING restricted to hereditary graph classes characterized by a *finite* number of minimal forbidden induced subgraphs. In particular they gave a complete classification of the complexity of COLOURING for the case where \mathcal{H} consists of a single graph H .

Theorem 1 ([43]). *If H is an induced subgraph of P_4 or of $P_1 + P_3$, then COLOURING restricted to H -free graphs is polynomial-time solvable, otherwise it is NP-complete.*

Theorem 1 led to two natural directions for further research:

1. Is it possible to obtain a dichotomy for COLOURING on H -free graphs if the number of colours k is fixed (that is, k no longer belongs to the input)?
2. Is it possible to obtain a dichotomy for COLOURING on \mathcal{H} -free graphs if \mathcal{H} has size 2?

We briefly discuss known results for both directions below and refer to [26] for a detailed survey. Let C_s and P_t denote the cycle on s vertices and path on t vertices, respectively. We start with the first question. If k is fixed, then we denote the problem by k -COLOURING. It is known that for every $k \geq 3$, the k -COLOURING problem on H -free graphs is NP-complete whenever H contains a cycle [22] or an induced claw [35, 44]. Therefore, only the case when H is a disjoint union of paths remains. In particular, the situation where $H = P_t$ has been thoroughly studied. On the positive side, 3-COLOURING on P_7 -free graphs [5], 4-COLOURING on P_6 -free graphs [11, 12] and k -COLOURING on P_5 -free graphs for any $k \geq 1$ [33] are polynomial-time solvable. On the negative side, Huang [36] proved NP-completeness for $(k = 5, t = 6)$ and for $(k = 4, t = 7)$. The case $(k = 3, t \geq 8)$ remains open, although some partial results are known [13].

In this paper we focus on the second question, that is, we restrict the input of COLOURING to \mathcal{H} -free graphs for $\mathcal{H} = \{H_1, H_2\}$. For two graphs G and H , we write $G + H = (V(G) \cup V(H), E(G) \cup E(H))$ for the disjoint union of two vertex-disjoint graphs G and H , and rG for the disjoint union of r copies of G . As a starting point, Král', Kratochvíl, Tuza, and Woeginger [43] identified the following three main sources of NP-completeness:

- both H_1 and H_2 contain a claw;
- both H_1 and H_2 contain a cycle; and
- both H_1 and H_2 contain an induced subgraph from the set $\{4P_1, 2P_1 + P_2, 2P_2\}$.

They also showed additional NP-completeness results by mixing the three types. Since then numerous papers [3, 9, 10, 17, 18, 20, 32, 34, 36, 40, 43, 46, 50, 51, 52, 56] have been devoted to this problem, but despite all these efforts the complexity classification

for COLOURING on (H_1, H_2) -free graphs is still far from complete, and even dealing with specific pairs (H_1, H_2) may require substantial work.

One of the “mixed” results obtained in [43] is that COLOURING is NP-complete for (C_s, H) -free graphs when $s \geq 5$ and $H \in \{4P_1, 2P_1 + P_2, 2P_2\}$. This, together with the well-known result that COLOURING can be solved in linear time for P_4 -free graphs (see also Theorem 1) implies the following dichotomy.

Theorem 2 ([43]). *Let $s \geq 5$ be a fixed integer. Then COLOURING for (C_s, P_t) -free graphs is polynomial-time solvable when $t \leq 4$ and NP-complete when $t \geq 5$.*

Theorem 2 raises the natural question: what is the complexity of COLOURING on (C_s, P_t) -free graphs when $s \in \{3, 4\}$?

For $s = 3$, Huang, Johnson and Paulusma [37] proved that 4-COLOURING, and thus COLOURING, is NP-complete for (C_3, P_{22}) -free graphs. A result of Brandstädt, Klemmt and Mahfud [7] implies that COLOURING is polynomial-time solvable for (C_3, P_6) -free graphs.

For $s = 4$, it is only known that COLOURING is polynomial-time solvable for (C_4, P_5) -free graphs [50]. This is unless we fix the number of colours: for every $k \geq 1$ and $t \geq 1$, it is known that k -COLOURING is polynomial-time solvable for $(K_{r,r}, P_t)$ -free graphs for every $s \geq 1$, and thus for (C_4, P_t) -free graphs [28] (take $r = 2$). The underlying reason for this is a result of Atminas, Lozin and Razgon [2], who proved that every P_t -free graph either has small treewidth or contains a large biclique $K_{s,s}$ as a subgraph. Then Ramsey arguments can be used but only if the number of colours k is fixed. The result for $s = 4$ and fixed k is in contrast to the result of [37] that for all $k \geq 4$ and $s \geq 5$, there exists a constant t_k^s such that k -COLOURING is NP-complete even for $(C_3, C_5, \dots, C_s, P_{t_k^s})$ -free graphs.

Our Main Results

We show, in Section 4, that COLOURING is polynomial-time solvable for (C_4, P_6) -free graphs. The class of (C_4, P_6) -free graphs generalizes the classes of split graphs (or equivalently, $(C_4, C_5, 2P_2)$ -free graphs) and pseudosplit graphs (or equivalently, $(C_4, 2P_2)$ -free graphs). The case of (C_4, P_6) -free graphs was explicitly mentioned as a natural case to consider in [26]. Our result unifies several previous results on colouring (C_4, P_t) -free graphs, namely: the polynomial-time solvability of COLOURING for (C_4, P_5) -free graphs [50]; the polynomial-time solvability of k -COLOURING for (C_4, P_6) -free graphs for every $k \geq 1$ [28]; and the recent 3/2-approximation algorithm for COLOURING for (C_4, P_6) -free graphs [24]. It also complements a recent result of Karthick and Maffray [41] who gave tight linear upper bounds of the chromatic number of a (C_4, P_6) -free graph in terms of its clique number and maximum degree that strengthen a similar bound given in [24].

It was not previously known if there exists an integer t such that COLOURING is NP-complete for (C_4, P_t) -free graphs. In Section 5 we complement our positive result of Section 4 by giving an affirmative answer to this question: already the value $t = 9$ makes the problem NP-complete.

Our Methodology

The general research goal of our paper is to increase, in a systematic way, our insights in the computational hardness of COLOURING by developing new techniques. In particular we

aim to narrow the complexity gaps between the hard and easy cases. Clique-width is a well-known width parameter and having bounded clique-width is often the underlying reason for a large collection of NP-complete problems, including COLOURING, to become polynomial-time solvable on a special graph class; this follows from results of [14, 23, 42, 54, 55]. For this reason we want to use clique-width to solve COLOURING for (C_4, P_6) -free graphs. However, the class of (C_4, P_6) -free graphs has unbounded clique-width, as it contains the class of *split graphs*, or equivalently, $(C_4, C_5, 2P_2)$ -free graphs, which may have arbitrarily large clique-width [49].

To overcome this obstacle we first preprocess the (C_4, P_6) -free input graph. An *atom* is a graph with no clique cutset. Clique cutsets were introduced by Dirac [21], who proved that every chordal graph is either complete or has a clique cutset. Later, decomposition into atoms became a very general tool for solving combinatorial problems on chordal graphs and other hereditary graph classes, such as those that forbid some Truemper configuration [4]. For instance, COLOURING and also other problems, such as INDEPENDENT SET and CLIQUE, are polynomial-time solvable on a hereditary graph class \mathcal{G} if they are so on the atoms of \mathcal{G} [58]. Hence, we may restrict ourselves to the subclass of (C_4, P_6) -free atoms in order to solve COLOURING for (C_4, P_6) -free graphs.

Adler et al. [1] proved that (diamond, even-hole)-free atoms have unbounded clique-width. However, so far, (un)boundedness of the cliquewidth of atoms in special graph classes has not been well studied. It is known that a class of H -free graphs has bounded clique-width if and only if H is an induced subgraph of P_4 (see [19]). As a start of a more systematic study, we show in Section 3 that the same result holds for atoms: a class of H -free atoms has bounded clique-width if and only if H is an induced subgraph of P_4 . In contrast, we observe that, although split graphs have unbounded clique-width [49], split atoms are cliques [21] and thus have clique-width at most 2. Recall that split graphs are characterized by three forbidden induced subgraphs. This yields the natural question whether one can prove the same result for a graph class characterized by two forbidden induced subgraphs. In this paper we give an *affirmative* answer to this question by showing that the class of (C_4, P_6) -free atoms has bounded clique-width. As mentioned, this immediately yields a polynomial-time algorithm for COLOURING on (C_4, P_6) -free graphs,

In order to prove that (C_4, P_6) -free atoms have bounded clique-width, we further develop the approach of [24] used to bound the chromatic number of (C_4, P_6) -free graphs as a linear function of their maximum clique size and to obtain a $3/2$ -approximation algorithm for COLOURING for (C_4, P_6) -free graphs. The approach of [24] is based on a decomposition theorem for (C_4, P_6) -free atoms. We derive a new variant of this decomposition theorem for so-called strong atoms, which are atoms that contain no universal vertices and no pairs of twin vertices. We use this decomposition to prove that (C_4, P_6) -free strong atoms have bounded clique-width. To obtain this result we also apply a divide-and-conquer approach for bounding the clique-width of a subclass of C_4 -free graphs. As another novel element of our proof, we show a new bound on the clique-width for (general) graphs in terms of the clique-width of recursively defined subgraphs induced by homogeneous triples and pairs of sets. Our techniques may be of independent interest and can possibly be used to prove polynomial-time solvability of COLOURING on other graph classes.

Remark. The INDEPENDENT SET problem is to decide if a given graph G has an independent set of at least k vertices for some given integer k . Brandstädt and Hoàng [6] proved that INDEPENDENT SET is polynomial-time solvable for (C_4, P_6) -free graphs. As

mentioned, just as for COLOURING, it suffices to consider only the atoms of a hereditary graph class in order to solve INDEPENDENT SET [58]. Brandstädt and Hoàng followed this approach. Although we will use one of their structural results as lemmas, their method does not yield a polynomial-time algorithm for COLOURING on (C_4, P_6) -free graphs.

2 Preliminaries

Let $G = (V, E)$ be a graph. For $S \subseteq V$, the subgraph *induced* by S , is denoted by $G[S] = (S, \{uv \mid u, v \in S\})$. The *complement* of G is the graph \overline{G} with vertex set V and edge set $\{uv \mid uv \notin E\}$. A clique $K \subseteq V$ is a *clique cutset* if $G - K$ has more connected components than G . If G has no clique cutsets, then G is called an *atom*. The *edge subdivision* of an edge $uv \in E$ removes uv from G and replaces it by a new vertex w and two new edges uw and wv .

The *neighbourhood* of a vertex v is denoted by $N(v) = \{u \mid uv \in E\}$ and its degree by $d(v) = |N(v)|$. For a set $X \subseteq V$, we write $N(X) = \bigcup_{v \in X} N(v) \setminus X$. For $x \in V$ and $S \subseteq V$, we let $N_S(x)$ be the set of neighbours of x that are in S , that is, $N_S(x) = N_G(x) \cap S$. A subset $D \subseteq V$ is a *dominating set* of G if every vertex not in D has a neighbour in D . A vertex u is *universal* in G if it is adjacent to all other vertices, that is, $\{u\}$ is a dominating set of G .

For $X, Y \subseteq V$, we say that X is *complete* (resp. *anti-complete*) to Y if every vertex in X is adjacent (resp. non-adjacent) to every vertex in Y . Let $u, v \in V$ be two distinct vertices. We say that a vertex $x \notin \{u, v\}$ *distinguishes* u and v if x is adjacent to exactly one of u and v . A set $H \subseteq V$ is a *homogeneous set* if no vertex in $V \setminus H$ can distinguish two vertices in H . A homogeneous set H is *proper* if $1 < |H| < |V|$. A graph is *prime* if it contains no proper homogeneous set.

We say that u and v are (*true*) *twins* if u and v are adjacent and have the same set of neighbours in $V \setminus \{u, v\}$. Note that the binary relation of being twins is an equivalence relation on V , and so V can be partitioned into equivalence classes T_1, \dots, T_r of twins. The *skeleton* of G is the subgraph induced by a set of r vertices, one from each of T_1, \dots, T_r . A *blow-up* of G is a graph G' obtained by replacing each vertex $u \in V$ by a clique K_u of size at least 1, such that two distinct cliques K_u and K_v are complete in G' if u and v are adjacent in G , and anti-complete otherwise. Since each equivalence class of twins is a clique and any two equivalence classes are either complete or anti-complete, every graph is a blow-up of its skeleton.

Let $\{H_1, \dots, H_p\}$ be a set of p graphs for some integer $p \geq 1$. We say that G is (H_1, \dots, H_p) -*free* if G contains no induced subgraph isomorphic to H_i for some $1 \leq i \leq p$. If $p = 1$, we may write that G is H_1 -free instead. If V can be partitioned into a clique C and an independent set I , then G is a *split graph*.

The *clique-width* of a graph G , denoted by $\text{cw}(G)$, is the minimum number of labels required to construct G using the following four operations:

- $i(v)$: create a new graph consisting of a single vertex v with label i ;
- $G_1 \oplus G_2$: take the disjoint union of two labelled graphs G_1 and G_2 ;
- $\eta_{i,j}$: join each vertex with label i to each vertex with label j (for $i \neq j$);
- $\rho_{i \rightarrow j}$: rename label i to j .




Figure 1: Walls of height 2, 3, and 4, respectively [19].

A *clique-width expression* of G is an algebraic expression that describes how G can be recursively constructed using these operations. An ℓ -*expression* of G is a clique-width expression using at most ℓ distinct labels. For instance, $\eta_{3,2}(3(d) \oplus \rho_{3 \rightarrow 2}(\rho_{2 \rightarrow 1}(\eta_{3,2}(3(c) \oplus \eta_{2,1}(2(b) \oplus 1(a))))))$ is a 3-expression for the path on vertices a, b, c, d in that order. A class of graphs \mathcal{G} has *bounded* clique-width if there is a constant c such that the clique-width of every graph in \mathcal{G} is at most c , and *unbounded* otherwise.

Clique-width is of fundamental importance in computer science, since every problem expressible in monadic second-order logic using quantifiers over vertex subsets but not over edge subsets becomes polynomial-time solvable for graphs of bounded clique-width [14]. Although this meta-theorem does not directly apply to COLOURING, a result of Espelage, Gurski and Wanke [23] (see also [42, 55]) combined with the polynomial-time approximation algorithm of Oum and Seymour [54] for finding an ℓ -expression of a graph, showed that COLOURING can be added to the list of such problems.

Theorem 3 ([23, 54]). COLOURING can be solved in polynomial time for graphs of bounded clique-width.

3 Atoms and Clique-Width

Recall that a graph is an atom if it contains no clique cutset. It is a natural question whether the clique-width of a graph class of unbounded clique-width becomes bounded after restricting to the atoms of the class. For classes of H -free graphs we note that this is not the case though. In order to explain this we need the notion of a wall; see Figure 1 for three examples (for a formal definition we refer to, for example, [16]). A k -*subdivided wall* is a graph obtained from a wall after subdividing each edge exactly k times for some constant $k \geq 0$. The following lemma is well known.

Lemma 1 ([48]). For every constant $k \geq 0$, the class of k -subdivided walls has unbounded clique-width.

We also need the following lemma.

Lemma 2. For every constant $k \geq 0$, every k -subdivided wall and every complement of a k -subdivided wall is an atom.

Proof. Let $k \geq 0$. Let W be a k -subdivided wall. As W is C_3 -free, a largest clique has size 2. It is readily seen that W contains no set of at most two vertices that disconnect W .

Now consider the complement \bar{W} of W . For contradiction, assume that \bar{W} is not an atom. Then \bar{W} has a clique cutset K . Let A and B be two connected components of $\bar{W} - K$. If A and B both have at least two vertices a_1, a_2 and b_1, b_2 , respectively, then $W[\{a_1, a_2, b_1, b_2\}]$ contains a C_4 , which is not possible. Hence, one of A, B , say A , only

contains one vertex a . As the neighbourhood of a in \overline{W} is a clique, the non-neighbourhood of a in W is an independent set. However, no vertex in W has this property. \square

Recall that a class of H -free graphs has bounded clique-width if and only if H is an induced subgraph of P_4 (see [19]). We show that the same classification holds for H -free atoms.

Proposition 1. *Let H be a graph. The class of H -free atoms has bounded clique-width if and only if H is an induced subgraph of P_4 .*

Proof. If H is an induced subgraph of P_4 , then the class of H -free graphs, which contains all H -free atoms, has clique-width at most 2 [15].

Now suppose that H is not an induced subgraph of P_4 . For every $k \geq 0$, every k -subdivided wall is an atom by Lemma 2. First suppose that H contains a cycle. Then the class of k -subdivided walls is contained in the class of H -free atoms for some appropriate value of k . Hence, the class of H -free atoms has unbounded clique-width due to Lemma 1.

Now suppose that H does not contain a cycle. Hence H is a forest. As H is not an induced subgraph of P_4 , we find that H must contain an induced $3P_1$ or an induced $2P_2$. Let \mathcal{G} be the class of H -free atoms, and let $\overline{\mathcal{G}}$ be the class that consists of the complements of H -free atoms. As every wall is (C_3, C_4) -free, the complement of every wall is $(3P_1, 2P_2)$ -free. By Lemma 2, the complement of every wall is an atom as well. Hence, $\overline{\mathcal{G}}$ contains all complements of walls. It is well known that complementing all graphs in a class of unbounded clique-width results in another class of unbounded clique-width [39]. Hence, complements of walls have unbounded clique-width due to Lemma 1. This means that $\overline{\mathcal{G}}$, and thus \mathcal{G} , has unbounded clique-width. \square

In contrast to Proposition 1, we recall that there exist classes of (H_1, H_2, H_3) -free graphs of unbounded clique-width whose atoms have bounded clique-width. Namely, the class of split graphs, or equivalently, the class of $(C_4, C_5, 2P_2)$ -free graphs, has unbounded clique-width [49], whereas split atoms are cliques and thus have clique-width at most 2. In the next section, we will prove that there exist even classes of (H_1, H_2) -free graphs with this property by showing that the property holds even for the class of (C_4, P_6) -free graphs.

4 The Polynomial-Time Result

In this section, we will prove our main result.

Theorem 4. *COLOURING is polynomial-time solvable for (C_4, P_6) -free graphs.*

The main ingredient for proving Theorem 4 is a new structural property of (C_4, P_6) -free atoms, which asserts that (C_4, P_6) -free atoms have bounded clique-width. The following result is due to Tarjan.

Theorem 5 ([58]). *If COLOURING is polynomial-time solvable on atoms in an hereditary class \mathcal{G} , then it is polynomial-time solvable on all graphs in \mathcal{G} .*

As the class of (C_4, P_6) -free graphs is hereditary, we can apply Theorem 5 and may restrict ourselves to (C_4, P_6) -free atoms. Then, due to Theorem 3, it suffices to show the following result in order to prove Theorem 4.

Theorem 6. *The class of (C_4, P_6) -free atoms has bounded clique-width. More precisely, every (C_4, P_6) -free atom has clique-width at most 18.*

Note that (C_4, P_6) -free atoms are an example of a class of (H_1, H_2) -free graphs of unbounded clique-width, whose atoms have bounded clique-width.

The remainder of the section is organised as follows. In Section 4.1, we present the key tools on clique-width that play an important role in the proof of Theorem 6. In Section 4.2, we list structural properties around a 5-cycle in a (C_4, P_6) -free graph that are frequently used in later proofs. We then present the proof of Theorem 6 in Section 4.3.

4.1 Key Tools for Clique-Width

Let $G = (V, E)$ be a graph and H be a proper homogeneous set in G . Then $V \setminus H$ is partitioned into two subsets N and M where N is complete to H and M is anti-complete to H . Let $h \in H$ be an arbitrary vertex and $G_h = G - (H \setminus \{h\})$. We say that H and G_h are *factors* of G with respect to H . Suppose that τ is an ℓ_1 -expression for G_h using labels $1, \dots, \ell_1$ and σ is an ℓ_2 -expression for H using labels $1, \dots, \ell_2$. Then substituting $i(h)$ in τ with $\rho_{1 \rightarrow i} \dots \rho_{\ell_2 \rightarrow i} \sigma$ results in an ℓ -expression for G where $\ell = \max\{\ell_1, \ell_2\}$. Moreover, all vertices in H have the same label in this ℓ -expression for G .

Lemma 3 ([15]). *The clique-width of any graph G is the maximum clique-width of any prime induced subgraph of G .*

A bipartite graph is a *chain graph* if it is $2P_2$ -free. A *co-bipartite chain graph* is the complement of a bipartite chain graph. Let G be a (not necessarily bipartite) graph such that $V(G)$ is partitioned into two subsets A and B . We say that an ℓ -expression for G is *nice* if all vertices in A end up with the same label i and all vertices in B end up with the same label j with $i \neq j$. It is well-known that any co-bipartite chain graph whose vertex set is partitioned into two cliques has a nice 4-expression (see Appendix A for a proof).

Lemma 4 (Folklore). *There is a nice 4-expression for any co-bipartite chain graph.*

We now use a divide-and-conquer approach to show that a special graph class has a nice 4-expression. This plays a crucial role in our proof of the main theorem (Theorem 6).

Lemma 5. *A C_4 -free graph G has a nice 4-expression if $V(G)$ can be partitioned into two (possibly empty) subsets A and B that satisfy the following conditions:*

- (i) $G[A]$ is a clique;
- (ii) $G[B]$ is P_4 -free;
- (iii) no vertex in A has two non-adjacent neighbours in B ;
- (iv) there is no induced P_4 in G that starts with a vertex in A followed by three vertices in B .

Proof. We use induction on $|B|$. If B contains at most one vertex, then G is a co-bipartite chain graph and the lemma follows from Lemma 4. Assume that B contains at least two vertices. Since $G[B]$ is P_4 -free, either B or \overline{B} is disconnected [57]. Suppose first that B is disconnected. Then B can be partitioned into two nonempty subsets B_1 and B_2 that are

We say that (X, Y, Z) is a *homogeneous triple* if no vertex in $V \setminus (X \cup Y \cup Z)$ can distinguish any two vertices in X , Y or Z . A pair (X, Y) of sets is a *homogeneous pair* if (X, Y, \emptyset) is a homogeneous triple. If both X and Y are cliques, then (X, Y) is a *homogeneous pair of cliques*. Note that homogeneous sets are special cases of homogeneous pairs and triples. An ℓ -expression for a homogeneous triple (X, Y, Z) is *nice* if two vertices of $X \cup Y \cup Z$ have the same label if and only if they belong to the same set X , Y or Z . We establish a new bound on the clique-width of a graph G in terms of the number of pairwise disjoint homogenous pairs and triples of G .

Lemma 6. *Let G be a graph. If $V(G)$ can be partitioned into a subset V_0 , with $|V_0| \geq 3$, and p homogeneous pairs and t homogeneous triples such that there is a nice 4-expression for each homogeneous pair and a nice 6-expression for each homogeneous triple, then $\text{cw}(G) \leq |V_0| + 2p + 3t$.*

Proof. We first construct the homogenous pairs and triples and the edges inside these pairs and triples one by one using nice 4-expressions and nice 6-expressions, respectively. So we need at most four different labels for each homogenous pair and at most six different labels for each homogenous triple. As soon as we have constructed a homogenous pair (triple) with its internal edges using a nice 4-expression (6-expression), we introduce a new label for all vertices of each of its two (three) sets before considering the next homogenous pair or triple. We can do so, because all vertices of each set in a homogeneous pair received the same label by the definition of a nice ℓ -expression for homogenous sets and triples. Consequently, we may use the previous labels over and over again as auxiliary labels. Afterwards, we can view each set in a homogeneous pair or triple as a single vertex, each with its own unique label.

So far we used at most $2p + 3t + 6$ different labels. By using the auxiliary labels as unique labels for the sets of the last pair or triple and by considering pairs before triples, we need in fact at most $2p + 3t + 3$ distinct labels if $t \geq 1$ and at most $2p + 2$ labels if $t = 0$. We now assign a unique label to each vertex in V_0 after first using all the remaining auxiliary labels.

So far we only constructed edges of G that are within a homogenous pair or triple. From our labelling procedure and the definitions of homogenous pairs and triples it follows that we can put in all the remaining edges of G using only join and disjoint union operations. Hence, as $|V_0| \geq 3$, the total number of distinct labels is at most $|V_0| + 2p + 3t$. \square

4.2 Structure around a 5-Cycle

Let $G = (V, E)$ be a graph and H be an induced subgraph of G . We partition $V \setminus V(H)$ into subsets with respect to H as follows: for any $X \subseteq V(H)$, we denote by $S(X)$ the set of vertices in $V \setminus V(H)$ that have X as their neighbourhood among $V(H)$, i.e.,

$$S(X) = \{v \in V \setminus V(H) : N_{V(H)}(v) = X\}.$$

For $0 \leq j \leq |V(H)|$, we denote by S_j the set of vertices in $V \setminus V(H)$ that have exactly j neighbours among $V(H)$. Note that $S_j = \bigcup_{X \subseteq V(H): |X|=j} S(X)$. We say that a vertex in S_j is a *j -vertex*. Let G be a (C_4, P_6) -free graph and $C = 1, 2, 3, 4, 5$ be an induced C_5 in G . We partition $V \setminus C$ with respect to C as above. All indices below are modulo 5. Since G is C_4 -free, there is no vertex in $V \setminus C$ that is adjacent to vertices i and $i + 2$ but not to vertex

$i + 1$. In particular, $S(1, 3)$, S_4 , etc. are empty. The following properties **(P1)**-**(P9)** of $S(X)$ were proved in [32] using the fact that G is (C_4, P_6) -free.

- (P1)** $S_5 \cup S(i - 1, i, i + 1)$ is a clique.
- (P2)** $S(i)$ is complete to $S(i + 2)$ and anti-complete to $S(i + 1)$. Moreover, if neither $S(i)$ nor $S(i + 2)$ are empty then both sets are cliques.
- (P3)** $S(i, i + 1)$ is complete to $S(i + 1, i + 2)$ and anti-complete to $S(i + 2, i + 3)$. Moreover, if neither $S(i, i + 1)$ nor $S(i + 1, i + 2)$ are empty then both sets are cliques.
- (P4)** $S(i - 1, i, i + 1)$ is anti-complete to $S(i + 1, i + 2, i + 3)$.
- (P5)** $S(i)$ is anti-complete to $S(j, j + 1)$ if $j \neq i + 2$. Moreover, if a vertex in $S(i + 2, i + 3)$ is not anti-complete to $S(i)$ then it is universal in $S(i + 2, i + 3)$.
- (P6)** $S(i)$ is anti-complete to $S(i + 1, i + 2, i + 3)$.
- (P7)** $S(i - 2, i + 2)$ is anti-complete to $S(i - 1, i, i + 1)$.
- (P8)** Either $S(i)$ or $S(i + 1, i + 2)$ is empty. By symmetry, either $S(i)$ or $S(i - 1, i - 2)$ is empty.
- (P9)** At least one of $S(i - 1, i)$, $S(i, i + 1)$ and $S(i + 2, i - 2)$ is empty.

We now prove some further properties that are used in Lemma 10.

- (P10)** *For each connected component A of $S(i)$, each vertex in $S(i - 2, i - 1, i) \cup S(i, i + 1, i + 2)$ is either complete or anti-complete to A .*

Proof. It suffices to prove the property for $i = 1$. Suppose that some vertex $t \in S(4, 5, 1) \cup S(1, 2, 3)$ is neither complete nor anti-complete to a connected component A of $S(1)$. By symmetry, we may assume that $t \in S(4, 5, 1)$. By the connectivity of A , there exists an edge aa' in A such that t is adjacent to a but not to a' . Then $a', a, t, 4, 3, 2$ induces a P_6 , a contradiction. ■

- (P11)** *No vertex in S_5 can distinguish an edge between $S(i)$ and $S(i - 2, i + 2)$.*

Proof. It suffices to prove the property for $i = 1$. Let $x \in S(1)$ and $y \in S(3, 4)$ be adjacent. If a vertex u is adjacent to exactly one of x and y , then either $x, y, 3, u$ or $x, y, u, 1$ induces a C_4 . ■

- (P12)** *If a vertex $x \in S(i - 2, i + 2)$ has a neighbour in $S(i - 2, i - 1, i) \cup S(i, i + 1, i + 2)$, then x is complete to S_5 .*




Figure 3: The Petersen graph.

Proof. It suffices to prove the property for $i = 1$. Suppose that x is not adjacent to some $u \in S_5$. Since x has a neighbour $s \in S(1, 2, 3) \cup S(4, 5, 1)$, say $S(1, 2, 3)$, it follows that $s, u, 4, x$ induces a C_4 . ■

(P13) *Each vertex in $S(i-2, i+2)$ is anti-complete to either $S(i-2, i-1, i)$ or $S(i, i+1, i+2)$.*

Proof. It suffices to prove the property for $i = 1$. Suppose that $x \in S(3, 4)$ has a neighbour $s \in S(1, 2, 3)$ and $t \in S(4, 5, 1)$. By (P4), s and t are not adjacent. Then $x, s, 1, t$ induces a C_4 . ■

(P14) *Each vertex in $S(i-1, i-2, i+2)$ and $S(i+1, i+2, i-2)$ is either complete or anti-complete to each connected component of $S(i-2, i+2)$.*

Proof. It suffices to prove the property for $i = 1$. Suppose that $s \in S(2, 3, 4) \cup S(3, 4, 5)$ distinguishes an edge xy in $S(3, 4)$, say s is adjacent to x but not to y . By symmetry, we may assume that $s \in S(2, 3, 4)$. Then $y, x, s, 2, 1, 5$ induces a P_6 . ■

(P15) *If both $S(i-1, i-2)$ and $S(i+1, i+2)$ are not empty, then each vertex in $S(i-1, i, i+1)$ is either complete or anti-complete to $S(i-1, i-2) \cup S(i+1, i+2)$.*


Proof. It suffices to prove the property for $i = 1$. Let $x \in S(2, 3)$ and $y \in S(4, 5)$ be two arbitrary vertices. If $s \in S(5, 1, 2)$ distinguishes x and y , say s is adjacent to x but not to y , then $1, s, x, 3, 4, y$ induces a P_6 , since y is not adjacent to x by (P3). ■

4.3 Proof of Theorem 6

In this section, we give a proof of Theorem 6, which states that (C_4, P_6) -free graphs have clique-width at most 18.

A graph is *chordal* if it does not contain any induced cycle of length at least 4. The following structure of (C_4, P_6) -free graphs discovered by Brandstädt and Hoàng [6] is of particular importance in our proofs below.

Theorem 7 ([6]). *Let G be a (C_4, P_6) -free atom. Then the following statements hold: (i) every induced C_5 is dominating; (ii) if G contains an induced C_6 which is not dominating, then G is the join of a blow-up of the Petersen graph (Figure 3) and a (possibly empty) clique.*

Figure 4: Two special graphs F_1 and F_2 .

We say that an atom is *strong* if it has no pair of twin vertices or universal vertices. Note that a pair of twin vertices and a universal vertex in a graph give rise to two special kinds of proper homogeneous sets such that one of the factors decomposed by these homogeneous sets is a clique. Therefore, removing twin vertices and universal vertices does not change the clique-width of the graph by Lemma 3. So, to prove Theorem 6 it suffices to prove the theorem for strong atoms.

We follow the approach of [24]. In [24], the first and second author showed how to derive a useful decomposition theorem for (C_4, P_6) -free atoms by eliminating a sequence F_1 , C_6 , F_2 and C_5 (see Figure 4 for the graphs F_1 and F_2) of induced subgraphs and then employing Dirac's classical theorem [21] on chordal graphs. Here we adopt the same strategy and show in Lemma 7–Lemma 10 below that if a (C_4, P_6) -free strong atom G contains an induced C_5 or C_6 , then it has clique-width at most 18. The remaining case is therefore that G is a chordal atom, and so G is a clique by Dirac's theorem [21]. Since cliques have clique-width 2, Theorem 6 follows. It turns out that we can easily prove Lemma 7 and Lemma 8 via the framework formulated in Lemma 6 using the structure of the graphs discovered in [24]. The difficulty is, however, that we have to extend the structural analysis in [24] extensively for Lemma 9 and Lemma 10 and provide new insights on bounding the clique-width of certain special graphs using divide-and-conquer (see Lemma 5).

Lemma 7. *If a (C_4, P_6) -free strong atom G contains an induced F_1 , then G has clique-width at most 13.*

Proof. Let G be a (C_4, P_6) -free strong atom that contains an induced subgraph H that is isomorphic to F_1 with $V(H) = \{1, 2, 3, 4, 5, x, y, z\}$ where 1, 2, 3, 4, 5, 1 induces the underlying 5-cycle C of F_1 and x is adjacent to 3 and 4, y is adjacent to 2 and 3, z is adjacent to 4 and 5, and x is adjacent to y and z , see Figure 4. We partition $V(G)$ with respect to C . We choose H such that $|S_2|$ maximized. Note that $x \in S(3, 4)$, $y \in S(2, 3)$ and $z \in S(4, 5)$. All indices below are modulo 5. Since G is an atom, it follows from Theorem 7 that $S_0 = \emptyset$. Moreover, it follows immediately from the (C_4, P_6) -freeness of G that $V(G) = C \cup S_1 \cup \bigcup_{i=1}^5 S(i, i+1) \cup \bigcup_{i=1}^5 S(i-1, i, i+1) \cup S_5$. If $S_5 \neq \emptyset$, then G is a blow-up of the graph F_3 (see Figure 5) [24]. Since G contains no twin vertices, G is isomorphic to F_3 and so has clique-width at most 9. If $S_5 = \emptyset$ then G has the structure prescribed in Figure 6 [24]. Note that $S(5, 1, 2) \cup \{1\}$ is a homogeneous clique in G and so $S(5, 1, 2) = \emptyset$. We partition $S(3, 4)$ into two subsets $X = \{x \in S(3, 4) : x \text{ has a neighbour in } S(4, 5, 1)\}$ and $Y = S(3, 4) \setminus X$. Note that Y is anti-complete to $S(4, 5, 1)$. In addition, X is anti-complete to $S(1, 2, 3)$ since G is C_4 -free. It is routine to check that each of $(X, S(4, 5, 1))$,

$(Y, S(1, 2, 3))$, $(S(2, 3), S(3, 4, 5))$ and $(S(4, 5), S(2, 3, 4))$ is a homogeneous pair of cliques in G . Now $V(G)$ is partitioned into a subset C of size 5 and four homogeneous pairs of cliques. Since each pair of homogeneous cliques induces a co-bipartite chain graph and so has a nice 4-expression by Lemma 4. So, $\text{cw}(G) \leq |V_0| + 2 \times 4 = 13$ by Lemma 6. \square




Figure 5: The graph F_3 .




Figure 6: The structure of G . A thick line between two sets represents that the two sets are complete, and a thin line represents that the edges between the two sets can be arbitrary. Two sets are anti-complete if there is no line between them.

Lemma 8. *If a (C_4, F_1, P_6) -free strong atom G contains an induced C_6 , then G has clique-width at most 13.*

Proof. Let $C = 1, 2, 3, 4, 5, 6, 1$ be an induced six-cycle of G . We partition $V(G)$ with respect to C . If C is not dominating, then it follows from Theorem 7 that G is the join of a blow-up of the Petersen graph and a (possibly empty) clique. Since G has no twin vertices or universal vertices, G is isomorphic to the Petersen graph and so G has clique-width at most 10. In the following, we assume that C is dominating, i.e., $S_0 = \emptyset$. It was shown in [24] that G is a blow-up of a graph of order at most 13. Therefore, G has clique-width at most 13. \square

Lemma 9. *If a (C_4, C_6, F_1, P_6) -free strong atom G contains an induced F_2 , then G has clique-width at most 14.*

Proof. Let G be a (C_4, C_6, F_1, P_6) -free strong atom that contains an induced subgraph H that is isomorphic to F_2 with $V(H) = \{1, 2, 3, 4, 5, t, x, y\}$ such that 1, 2, 3, 4, 5, 1 induces the underlying 5-cycle C , and t is adjacent to 5, 1 and 2, x is adjacent to 4, 5 and y is adjacent to 2 and 3. Moreover, t is adjacent to both x and y , see Figure 4. We partition $V(G)$ with respect to C . We choose H such that C has $|S_2|$ maximized. Note that $x \in S(4, 5)$, $y \in S(2, 3)$ and $t \in S(5, 1, 2)$.

The overall strategy is to first decompose G into a subset V_0 of constant size and constant number of homogeneous pairs of sets, and then finish off the proof via Lemma 6 by showing that each homogeneous pair of sets has a nice 4-expression using Lemma 5.

We start with the decomposition. Since $S(2, 3)$ and $S(4, 5)$ are not empty, it follows from **(P8)** that $S_1 = S(2) \cup S(5)$. If both $S(2)$ and $S(5)$ are not empty, say $u \in S(2)$ and $v \in S(5)$, then $u, 2, 3, 4, 5, v$ induces either a P_6 or a C_6 , depending on whether u and v are adjacent. This shows that $S_1 = S(i)$ for some $i \in \{2, 5\}$. Now we argue that $S_2 = S(2, 3) \cup S(4, 5)$. If $S(3, 4)$ contains a vertex z , then z is adjacent to x and y by **(P3)** but not adjacent to t by **(P7)**. This implies that t, x, z, y induces a C_4 . So, $S(3, 4) = \emptyset$. If $S(1, 2)$ contains a vertex z , then z is adjacent to y by **(P3)** and so $1, z, y, 3, 4, 5, 1$ induces a C_6 , a contradiction. This shows that $S(1, 2) = \emptyset$. By symmetry, $S(5, 1) = \emptyset$. Therefore, $S_2 = S(2, 3) \cup S(4, 5)$. The following properties among subsets of G were proved in [24].

- (a) Each vertex in $S(5, 1, 2)$ is either complete or anti-complete to S_2 .
- (b) $S(2, 3)$ and $S(4, 5)$ are cliques.
- (c) Each vertex in $S(3, 4, 5) \cup S(4, 5, 1)$ is either complete or anti-complete to $S(4, 5)$. By symmetry, each vertex in $S(1, 2, 3) \cup S(2, 3, 4)$ is either complete or anti-complete to $S(2, 3)$.
- (d) $S(4, 5)$ is anti-complete to $S(2, 3, 4)$. By symmetry, $S(2, 3)$ is anti-complete to $S(3, 4, 5)$.
- (e) $S(1, 2, 3)$ is complete to $S(5, 1, 2)$. By symmetry, $S(5, 1, 2)$ is complete to $S(4, 5, 1)$.
- (f) $S(4, 5)$ is complete to $S(4, 5, 1)$. By symmetry, $S(2, 3)$ is complete to $S(1, 2, 3)$.
- (g) $S(1, 2, 3)$ is complete to $S(2, 3, 4)$. By symmetry, $S(3, 4, 5)$ is complete to $S(4, 5, 1)$.
- (h) S_5 is complete to S_2 .

Recall that $S_1 = S(i)$ for some $i \in \{2, 5\}$. By symmetry, we may assume that $S_1 = S(5)$. Note that $S(5)$ is complete to $S(4, 5, 1)$ by Theorem 7 and anti-complete to $S(1, 2, 3)$ by **(P6)**. It follows from **(P1)**, **(P4)**, **(P7)**, **(e)**, **(f)** and **(g)** that $S(i-1, i, i+1) \cup \{i\}$ is a homogeneous clique in G and therefore $S(i-1, i, i+1) = \emptyset$ for $i = 2, 5$. Similarly, $S(4, 5)$ is a homogeneous clique in G by **(P7)**, **(a)-(d)**, **(f)** and **(h)** and so $S(4, 5) = \{x\}$. Let $T = \{t \in S(5, 1, 2) : t \text{ is complete to } S_2\}$.

(1) $S(5)$ is anti-complete to $S(5, 1, 2) \setminus T$.

Let $u \in S(5)$ and $t' \in S(5, 1, 2) \setminus T$. If u and t' are adjacent, then $u, t', 2, 3, 4, x$ induces either a P_6 or a C_6 , depending on whether u and x are adjacent. ■

By (1) and (d), $(S(5, 1, 2) \setminus T) \cup \{1\}$ is a homogeneous set in G and so $S(5, 1, 2) \setminus T = \emptyset$. In other words, $S(5, 1, 2)$ is complete to S_2 . We now partition $S(5)$ into $X = \{v \in S(5) : v \text{ has a neighbour in } S(2, 3)\}$ and $Y = S(5) \setminus X$.

(2) X is anti-complete to $S(3, 4, 5)$.

Let $v \in X$ and $s \in S(3, 4, 5)$ be adjacent. By the definition of X , v has a neighbour $y' \in S(2, 3)$. By (d), y' is not adjacent to s and so $v, y', 3, s$ induces a C_4 . ■

(3) X is complete to $S(5, 1, 2)$.

Assume, by contradiction, that $v \in X$ and $t' \in T$ are not adjacent. By the definition of X , v has a neighbour $y' \in S(2, 3)$. Since t' is adjacent to y' , it follows that $v, 5, t', y'$ induces a C_4 . ■

(4) X is anti-complete to Y .

Suppose that $u \in X$ and $v \in Y$ are adjacent. Let $y' \in S(2, 3)$ be a neighbour of u . Note that x is adjacent to neither u nor v by (P5). But now $x, 4, 3, y', u, v$ induces a P_6 . ■

(5) X is complete to S_5 .

Suppose that $v \in X$ and $u \in S_5$ are not adjacent. Let $y' \in S(2, 3)$ be a neighbour of v . By (h), y' and u are adjacent. Then $u, 5, v, y'$ induces a C_4 . ■

It follows from (P1)-(P7), (a)-(d), (f), (h) and (2)-(5) that $(X, S(2, 3))$ is a homogeneous pair of sets in G .

(6) For each connected component A of Y , each vertex in $S(5, 1, 2) \cup S(3, 4, 5)$ is either complete or anti-complete to A .

Let A be an arbitrary connected component of Y . Suppose that $s \in S(5, 1, 2) \cup S(3, 4, 5)$ distinguishes an edge aa' in A , say s is adjacent to a but not adjacent to a' . We may assume by symmetry that $s \in S(5, 1, 2)$. Then $a', a, s, 2, 3, 4$ induces a P_6 , a contradiction. ■

(7) Each connected component of Y has a neighbour in both $S(5, 1, 2)$ and $S(3, 4, 5)$.

Suppose that a connected component A of Y does not have a neighbour in one of $S(5, 1, 2)$ and $S(3, 4, 5)$, say $S(5, 1, 2)$. Then $S_5 \cup S(3, 4, 5) \cup \{5\}$ is a clique cutset of G by (4). ■

(8) Each connected component of Y is a clique.

Let A be an arbitrary connected component of Y . By (7), A has a neighbour $s \in S(5, 1, 2)$ and $r \in S(3, 4, 5)$. Note that s and r are not adjacent. Moreover, $\{s, r\}$ is complete to A by (6). Now (8) follows from the fact that G is C_4 -free. ■

(9) Y is complete to S_5 .

Suppose, by contradiction, that $v \in Y$ and $u \in S_5$ are not adjacent. By (7), v has a neighbour $s \in S(5, 1, 2)$ and $r \in S(3, 4, 5)$. Then v, s, u, r induces a C_4 . ■

It follows from **(P1)**, **(h)**, **(5)** and **(9)** that each vertex in S_5 is a universal vertex in G and so $S_5 = \emptyset$. Let $S'(3, 4, 5) = \{s \in S(3, 4, 5) : s \text{ has a neighbour in } Y\}$ and $S''(3, 4, 5) = S(3, 4, 5) \setminus S'(3, 4, 5)$. Note that $S''(3, 4, 5)$ is anti-complete to Y . We now show further properties of Y and $S'(3, 4, 5)$.

(10) $S'(3, 4, 5)$ is complete to $S(2, 3, 4)$.

Suppose, by contradiction, that $r' \in S'(3, 4, 5)$ is not adjacent to $s \in S(2, 3, 4)$. By the definition of $S'(3, 4, 5)$, r' has a neighbour $v \in Y$. Then $v, r, 4, s, 2, 1$ induces a P_6 , a contradiction. ■

(11) Each vertex in $S(5, 1, 2)$ is either complete or anti-complete to Y .

Let $t' \in S(5, 1, 2)$ be an arbitrary vertex. Suppose that t' has a neighbour $u \in Y$. Let A be the connected component of Y containing u . Then t' is complete to A by **(6)**. It remains to show that t' is adjacent to each vertex $u' \in Y \setminus A$. By **(7)**, u has a neighbour $s \in S(3, 4, 5)$. Note that $C' = u, t', y, 3, s$ induces a C_5 . Moreover, x and s are not adjacent for otherwise x, s, u, t' induces a C_4 . This implies that x is adjacent only to t' on C' . On the other hand, u' is not adjacent to any of $u, 3$ and y . This implies that u' is adjacent to either s or t' by Theorem 7. If u' is not adjacent to t' , then u' is adjacent to s . This implies that $u', s, 3, y, t', x$ induces a P_6 or C_6 , depending on whether u' and x are adjacent. Therefore, u' is adjacent to t' . Since u' is an arbitrary vertex in $Y \setminus A$, this proves **(11)**. ■

(12) $S'(3, 4, 5)$ is anti-complete to x .

Suppose not. Let $s \in S'(3, 4, 5)$ be adjacent to x . By definition, s has a neighbour $y' \in Y$. Note that x and y' are not adjacent by **(P5)**. By **(6)** and **(7)**, y' has a neighbour $t \in T = S(5, 1, 2)$. So, t is adjacent to x . But now s, y', t, x induces a C_4 , a contradiction. ■

It follows from **(P1)**-**(P7)**, **(d)**, **(2)**, **(4)**, **(10)**, **(11)** and **(12)** that $(Y, S'(3, 4, 5))$ is a homogeneous pair of sets in G .

Let $S'(5, 1, 2) = \{s \in S(5, 1, 2) : s \text{ is complete to } Y\}$. Then $S(5, 1, 2) \setminus S'(5, 1, 2)$ is anti-complete to Y by **(11)**. It follows from **(3)** that both $S'(5, 1, 2)$ and $S(5, 1, 2) \setminus S'(5, 1, 2)$ are homogeneous cliques in G . So, $|S(5, 1, 2)| \leq 2$. We now show that $|S''(3, 4, 5)| \leq 1$ and $|S(2, 3, 4)| \leq 1$. First, we note that if $r \in S(3, 4, 5) \cap N(x)$, then r is complete to $S(2, 3, 4)$ for otherwise any non-neighbour $s \in S(2, 3, 4)$ of r would start an induced $P_6 = s, 3, r, x, t, 1$. By symmetry, each vertex in $S(2, 3, 4) \cap N(y)$ is complete to $S(3, 4, 5)$. By **(c)**, each vertex in $S(2, 3, 4) \cap N(y)$ is also complete to $S(2, 3)$. Thus, $(S''(3, 4, 5) \cap N(x)) \cup \{4\}$ and $(S(2, 3, 4) \cap N(y)) \cup \{3\}$ are homogeneous cliques in G . So, $S''(3, 4, 5) \cap N(x) = S(2, 3, 4) \cap N(y) = \emptyset$. Namely, x and y are anti-complete to $S''(3, 4, 5)$ and $S(2, 3, 4)$, respectively. By **(c)**, it follows that $S(2, 3)$ is anti-complete to $S(2, 3, 4)$. Secondly, $S''(3, 4, 5)$ and $S(2, 3, 4)$ are anti-complete to each other. If $r \in S''(3, 4, 5)$ and $s \in S(2, 3, 4)$ are adjacent, then $x, 5, r, s, 2, y$ induces a P_6 by **(d)** and the fact that x and y are not adjacent to r and s , respectively. These two properties imply that each of $S''(3, 4, 5)$ and $S(2, 3, 4)$ is a homogeneous clique in G . Hence, $|S''(3, 4, 5)| \leq 1$ and $|S(2, 3, 4)| \leq 1$. Now G is partitioned into a subset $V(C) \cup S(5, 1, 2) \cup S''(3, 4, 5) \cup S(2, 3, 4) \cup \{x\}$ of size at most 10 and two homogeneous pairs of sets $(X, S(2, 3))$ and $(Y, S'(3, 4, 5))$.

We now show that each of $G[X \cup S(2, 3)]$ and $G[Y \cup S'(3, 4, 5)]$ has a nice 4-expression. First, note that no vertex in $S(1, 2)$ can have two non-adjacent neighbours in X since G is C_4 -free. If there is an induced $P_4 = y', x_1, x_2, x_3$ such that $y' \in S(2, 3)$ and $x_i \in X$, then $x_3, x_2, x_1, y', 3, 4$ induces a P_6 in G . Now if $P = x_1, x_2, x_3, x_4$ is an induced P_4 in $G[X]$, any neighbour y_1 of x_1 is not adjacent to x_3 and x_4 . But then $P \cup \{y_1\}$ contains such a labelled P_4 in $G[X \cup S(2, 3)]$. Therefore, $G[X \cup S(2, 3)]$ with the partition $(X, S(2, 3))$ satisfies all the conditions of Lemma 5 and so has a nice 4-expression. Finally, note that each vertex in $S(3, 4, 5)$ can have neighbours in at most one connected component of Y due to (7), (11) and the fact that G is C_4 -free. It then follows from (6)-(8) that $G[Y \cup S'(3, 4, 5)]$ with the partition $(Y, S'(3, 4, 5))$ satisfies all the condition in Lemma 5 (where $A = S'(3, 4, 5)$ and $B = Y$) and so has a nice 4-expression. By Lemma 6, it follows that $\text{cw}(G) \leq 10 + 2 \times 2 = 14$. This completes our proof. \square

Lemma 10. *If a $(C_4, C_6, F_1, F_2, P_6)$ -free strong atom G contains an induced C_5 , then G has clique-width at most 18.*

Proof. Let $C = 1, 2, 3, 4, 5, 1$ be an induced C_5 of G . We partition $V(G) \setminus C$ with respect to C . We choose C such that

- $|S_5|$ is maximized, and
- $|S_3|$ is minimized subject to the above.

We first prove the following claim which makes use of the choice of C .

Claim 1. *For each $1 \leq i \leq 5$, $S(i-1, i, i+1)$ is complete to $S(i, i+1, i+2)$.*

Proof of Claim 1. By symmetry, it suffices to prove the claim for $i = 1$. Suppose by contradiction that $S(5, 1, 2)$ is not complete to $S(1, 2, 3)$. Then there exist vertices $s \in S(5, 1, 2)$ and $t \in S(1, 2, 3)$ that are not adjacent. Consider the induced 5-cycle $C' = C \setminus \{1\} \cup \{s\}$. Note that t is not a 3-vertex with respect to C' . By the choice of C , there must exist a vertex $r \in V(G)$ that is a 3-vertex for C' but not for C . By the definition of 3-vertices, it follows that $r \in S(2, 3) \cup S(4, 5)$ and r is adjacent to s . Similarly, by considering the induced 5-cycle $C'' = C \setminus \{2\} \cup \{t\}$ we conclude that there exist a vertex $q \in S(1, 5) \cup S(3, 4)$ that is adjacent to t . Note that r is not adjacent to t for otherwise $r, t, 1, s$ induces a C_4 . By symmetry, q is not adjacent to s . If $r \in S(2, 3)$, then $4, q, t, 1, s, r$ induces a P_6 if $q \in S(3, 4)$ and $3, r, s, 5, q, t$ induces a C_6 if $q \in S(5, 1)$. This shows that $r \in S(4, 5)$. By symmetry, $q \in S(3, 4)$. But now $5, 1, 2, 3, r, q$ induces a C_6 since r and q are adjacent by (P3). This contradicts the assumption that G is C_6 -free. \blacksquare

Claim 2. *If $S(i)$ contains a vertex that is anti-complete to $S(i-2, i+2)$, then G contains an induced F_2 .*

Proof of Claim 2. By symmetry, we may assume that $i = 1$. First note that $S(1)$ is anti-complete to $S_1 \setminus S(1)$ by (P2) and the C_6 -freeness of G . So, the neighbours of the vertices in $S(1)$ are in $\{1\} \cup S(3, 4) \cup S(4, 5, 1) \cup S(5, 1, 2) \cup S(1, 2, 3) \cup S_5$ by (P5) and (P6). Let $X \subseteq S(1)$ be the set of vertices that have a neighbour in $S(3, 4)$, and $S'(1) = S(1) \setminus X$. Note that $S'(1) \neq \emptyset$ due to our assumption. Let $a \in S'(1)$ and A be the connected component of $S'(1)$ containing a . If A has a neighbour in both $S(1, 2, 3)$ and $S(4, 5, 1)$, it follows from (P10) that G contains an induced F_2 . Therefore, we may assume that A is anti-complete to $S(4, 5, 1)$. This implies that $N(A) \subseteq \{1\} \cup S(5, 1, 2) \cup S(1, 2, 3) \cup S_5 \cup X$. Note that

$S(5, 1, 2)$ is complete to $S(1, 2, 3)$ by Claim 1. So, $\{1\} \cup S(1, 2, 3) \cup S(5, 1, 2) \cup S_5$ is a clique by **(P1)**. If $x \in X$ distinguishes an edge aa' of A , say x is adjacent to a but not to a' , then $a', a, s, y, 4, 5$ induces a P_6 , where $y \in S(3, 4)$ is a neighbour of x . So, each vertex in X is either complete or anti-complete to A . Recall that each vertex in $S(1, 2, 3)$ is either complete or anti-complete to A by **(P10)**. Let X' and $S'(1, 2, 3)$ be the set of vertices in X and $S(1, 2, 3)$ that are complete to A , respectively. If $x' \in X'$ and $s' \in S'(1, 2, 3)$ are not adjacent, let $y \in S(3, 4)$ be a neighbour of x' . Then either s', a, x', y induces a C_4 or $s', a, x', y, 4, 5$ induces a P_6 depending on whether s' and y are adjacent. So, X' and $S'(1, 2, 3)$ are complete. Suppose that X' contains two non-adjacent vertices x_1 and x_2 . Let $y_i \in S(3, 4)$ be a neighbour of x_i for $i = 1, 2$. If $y_1 = y_2$, then $y_1, x_1, 1, x_2$ induces a C_4 . So, $y_1 \neq y_2$. This means that x_1 is not adjacent to y_2 and x_2 is not adjacent to y_1 . By **(P5)**, y_1 is adjacent to y_2 and so y_1, x_1, a, x_2, y_2 induces a C_5 not dominating 2, which contradicts Theorem 7. This shows that X' is a clique. We have proved that $N(A) \setminus S_5$ is a clique. Since G contains no clique cutset, A must have a pair of non-adjacent neighbours $u \in S_5$ and $x \in X'$. By the definition of X' , x has a neighbour $y \in S(3, 4)$. Note that u is not adjacent to y by **(P11)**. We may choose $a \in A$ to be a neighbour of u . But now $\{1, 2, 3, 4, a, x, y, u\}$ induces an F_2 (where the underlying 5-cycle is $1, 2, 3, y, x$). ■

Claim 3. *The set $S(i)$ is anti-complete to $S(i-2, i-1, i) \cup S(i, i+1, i+2)$.*

Proof of Claim 3. By symmetry, we assume that $i = 1$. Suppose, by contradiction, that $x \in S(1)$ is adjacent to some vertex $s \in S(1, 2, 3) \cup S(4, 5, 1)$. We may assume by symmetry that $s \in S(4, 5, 1)$. By Claim 2, x has a neighbour $y \in S(3, 4)$. Note that $C' = 1, x, y, 4, 5$ induces a 5-cycle and s is complete to C' . By the choice of C' , there exists $u \in S_5$ such that u is not complete to C' . By **(P11)**, u is anti-complete to $\{x, y\}$. However, this contradicts **(P12)**. ■

Claim 4. *If $S(i-2, i+2) \neq \emptyset$ and is anti-complete to S_1 , then each vertex in $S(i-2, i+2)$ has a neighbour in $S(i-2, i-1, i) \cup S(i, i+1, i+2)$, and $S(i-2, i+2)$ is a clique and it is anti-complete to exactly one of $S(i-2, i-1, i)$ and $S(i, i+1, i+2)$.*

Proof of Claim 4. By symmetry, assume that $S(3, 4) \neq \emptyset$. It follows from **(P3)**, **(P9)** and the C_6 -freeness of G that $S_2 = S(3, 4) \cup S(i, i+1)$ for some $i \in \{5, 1\}$, and $S(3, 4)$ is anti-complete to $S_2 \setminus S(3, 4)$. We first show that each vertex in $S(3, 4)$ has a neighbour in $S(4, 5, 1) \cup S(1, 2, 3)$. Let $Y \subseteq S(3, 4)$ be the set of vertices that have a neighbour in $S(4, 5, 1) \cup S(1, 2, 3)$, and $S'(3, 4) = S(3, 4) \setminus Y$. Suppose that the claim does not hold, namely $S'(3, 4) \neq \emptyset$. We shall show that G contains a clique cutset and this is a contradiction. Let $b \in S'(3, 4)$ and B be the connected component of $S'(3, 4)$ containing b . First, if $y \in Y$ distinguishes an edge bb' in B , say y is adjacent to b but not to b' , then $b', b, y, s, 1, 2$ or $b', b, y, s, 1, 5$ induces a P_6 where $s \in S(1, 2, 3) \cup S(4, 5, 1)$ is a neighbour of y . So, any vertex Y is either complete or anti-complete to B . Let $Y', S'(2, 3, 4)$ and $S'(3, 4, 5)$ be the subsets of $Y, S(2, 3, 4)$ and $S(3, 4, 5)$, respectively that are complete to B . Note that $N(B) = \{3, 4\} \cup S'(2, 3, 4) \cup S'(3, 4, 5) \cup S_5 \cup Y'$ by **(P7)**. If $p \in S'(2, 3, 4)$ is not adjacent to $q \in S'(3, 4, 5)$, then $p, b, q, 5, 1, 2, p$ induces a C_6 , a contradiction. This shows that $\{3, 4\} \cup S'(2, 3, 4) \cup S'(3, 4, 5) \cup S_5$ is a clique by **(P1)**.

Now let $y_1, y_2 \in Y'$ be two arbitrary and distinct vertices. Suppose that y_1 and y_2 are not adjacent. Let $x_i \in S(4, 5, 1) \cup S(1, 2, 3)$ be a neighbour of y_i for $i = 1, 2$. If $x_1 = x_2$, then b, y_1, x_1, y_2 induces a C_4 . So, $x_1 \neq x_2$ and this implies that x_1 (resp. x_2) is not

adjacent to y_2 (resp. y_1). If x_1 and x_2 are in $S(4, 5, 1)$, then b, y_1, x_1, x_2, y_2 induces a C_5 not dominating 2, which contradicts Theorem 7; if $x_1 \in S(4, 5, 1)$ and $x_2 \in S(1, 2, 3)$, then $b, y_1, x_1, 1, x_2, y_2, b$ induces a C_6 , which contradicts our assumption. So, Y' is a clique. Moreover, Y' is complete to $S'(2, 3, 4) \cup S'(3, 4, 5)$ by **(P14)** and to S_5 by **(P12)**. Therefore, $N(B)$ is a clique. This proves the first statement of the claim.

Let $X_2 = \{y \in S(3, 4) : y \text{ has a neighbour in } S(1, 2, 3)\}$ and $X_5 = \{y \in S(3, 4) : y \text{ has a neighbour in } S(4, 5, 1)\}$. The by **(P13)**, X_2 and X_5 form a partition of $S(3, 4)$, and X_2 is anti-complete to $S(4, 5, 1)$ and X_5 is anti-complete to $S(1, 2, 3)$. If $y \in X_2$ and $z \in X_5$ are adjacent, let $t_2 \in S(1, 2, 3)$ and $t_5 \in S(4, 5, 1)$ be neighbours of y and z , respectively. Then $5, t_5, z, y, t_2, 2$ induces a P_6 , a contradiction. This shows that X_2 and X_5 are anti-complete to each other. Suppose now that $x \in X_5$ and $y \in X_2$. Then x has a neighbour $t \in S(4, 5, 1)$ and y has a neighbour $s \in S(1, 2, 3)$. Note also that x and y are not adjacent. Now $(C \setminus \{5\}) \cup \{x, y, s, t\}$ induces a F_2 (whose underlying 5-cycle is $y, s, 1, t, 4$). Therefore, we may assume by symmetry that $X_5 = \emptyset$. In other words, every vertex in $S(3, 4)$ has a neighbour in $S(1, 2, 3)$. It remains to show that $S(3, 4)$ is a clique. Suppose that xs and $x's'$ induce a $2P_2$ in the bipartite graph between $S(1, 2, 3)$ and $S(3, 4)$ where $x, x' \in S(3, 4)$ and $s, s' \in S(1, 2, 3)$. Note that s and s' are adjacent since $S(1, 2, 3)$ is a clique. Thus, x and x' are not adjacent for otherwise x, s, s', x' induces a C_4 . Now $C' = s, x, 4, x', s'$ induces a 5-cycle. By **(P12)**, $\{x, x'\}$ is complete to S_5 and this implies that S_5 is complete to C' . Moreover, 3 is complete to C' and this contradicts the choice of C . Now we can order the vertices in $S(3, 4)$ as x_1, \dots, x_r such that

$$N_{S(1,2,3)}(x_1) \subseteq N_{S(1,2,3)}(x_2) \dots \subseteq N_{S(1,2,3)}(x_r).$$

Recall that x_1 has a neighbour $t \in S(1, 2, 3)$ and thus t is complete to $S(3, 4)$. This implies that $S(3, 4)$ is a clique since G is C_4 -free. \blacksquare

We now consider two cases.

Case 1. $S_1 = \emptyset$. By Claim 4 and **(P12)**, S_5 is complete to S_2 and so each vertex in S_5 is a universal vertex in G by **(P1)**. Therefore, $S_5 = \emptyset$. Recall that $S(i-1, i, i+1)$ and $S(i, i+1, i+2)$ are complete for each $1 \leq i \leq 5$ by Claim 1. Suppose first that $S_2 = \emptyset$. Then $S(i-1, i, i+1) \cup \{i\}$ is a homogeneous clique in G by **(P1)** and **(P4)**. So, $S(i-1, i, i+1) = \emptyset$. Now G is the 5-cycle and has clique-width at most 4.

Suppose now that $S_2 \neq \emptyset$, say $S(3, 4) \neq \emptyset$. By Claim 4, we may assume that $S(3, 4)$ is anti-complete to $S(4, 5, 1)$, each vertex in $S(3, 4)$ has a neighbour in $S(1, 2, 3)$, and $(S(3, 4), S(1, 2, 3))$ is a co-bipartite chain graph. Let $x' \in S(3, 4)$ and $s' \in S(1, 2, 3)$ be a neighbour of x' . If $S(1, 5)$ contains a vertex y , then either $2, s', x', 4, 5, y$ induces a P_6 or $C \cup \{x', y, s'\}$ induces a subgraph isomorphic to F_2 , depending on whether s' and y are adjacent. This shows that $S(1, 5) = \emptyset$. In addition, $S(2, 3) = S(4, 5) = \emptyset$ by **(P3)** and the C_6 -freeness of G . So, $S_2 = S(3, 4) \cup S(1, 2)$. Suppose that $x \in S(3, 4)$ and $t \in S(3, 4, 5)$ are not adjacent. Let $s \in S(1, 2, 3)$ be a neighbour of x . Then $C \cup \{x, s, t\}$ induces a subgraph isomorphic to F_2 (where the underlying 5-cycle is $x, s, 1, 5, 4$). This shows that $S(3, 4)$ is complete to $S(3, 4, 5)$.

Note that $S(1, 2)$ may not be empty. If $S(1, 2) \neq \emptyset$, then it is anti-complete to $S(4, 5, 1)$ by **(P15)** and the fact that $S(4, 5, 1)$ is anti-complete to $S(3, 4)$. It then follows from Claim 4 that each vertex in $S(1, 2)$ has a neighbour in $S(2, 3, 4)$, and $(S(1, 2), S(2, 3, 4))$ is a co-bipartite chain graph. In addition, the above argument for $S(3, 4)$ works symmetrically for $S(1, 2)$. In particular, $S(1, 2)$ is complete to $S(5, 1, 2)$. This implies that

$S(i-1, i, i+1) \cup \{i\}$ for each $i = 1, 4, 5$ is a homogeneous clique in G by **(P4)** and **(P7)**. Therefore, $S(i-1, i, i+1) = \emptyset$ for $i = 1, 4, 5$. Let $S'(2, 3, 4)$ be the set of vertices in $S(2, 3, 4)$ that are complete to $S(3, 4)$ and $S''(2, 3, 4) = S(2, 3, 4) \setminus S'(2, 3, 4)$. Similarly, if $S(1, 2) \neq \emptyset$, let $S'(1, 2, 3)$ be the set of vertices in $S(1, 2, 3)$ that are complete to $S(1, 2)$ and $S''(1, 2, 3) = S(1, 2, 3) \setminus S'(1, 2, 3)$. In case that $S(1, 2) = \emptyset$, we define $S'(1, 2, 3) = S(1, 2, 3)$ and $S''(1, 2, 3) = \emptyset$. By **(P14)** and the way we define $S'(1, 2, 3)$ and $S''(1, 2, 3)$, it follows that $S''(2, 3, 4)$ and $S''(1, 2, 3)$ are anti-complete to $S(3, 4)$ and $S(1, 2)$, respectively. Let $x \in S(3, 4)$. If $s \in S''(2, 3, 4)$ is adjacent to some $y \in S(1, 2)$, then $x, 3, s, y, 1, 5$ induces a P_6 . So, $S''(2, 3, 4)$ is anti-complete to $S(1, 2)$. By symmetry, $S''(1, 2, 3)$ is anti-complete to $S(3, 4)$ (note that this is vacuously true if $S(1, 2) = \emptyset$ since we define $S''(1, 2, 3) = \emptyset$). This implies that $S''(2, 3, 4)$ and $S''(1, 2, 3)$ are homogeneous cliques in G and so both have size at most 1. Furthermore, $(S'(2, 3, 4), S(1, 2))$ and $(S'(1, 2, 3), S(3, 4))$ are homogeneous pairs of cliques in G . Now $V(G)$ is partitioned into a subset $C \cup S''(2, 3, 4) \cup S''(1, 2, 3)$ of size at most 7 and two homogeneous pairs of cliques each of which has a nice 4-expression by Lemma 4. Therefore, $\text{cw}(G) \leq 7 + 2 \times 2 = 11$ by Lemma 6. This completes the proof of Case 1.

Case 2. $S_1 \neq \emptyset$. By symmetry, we assume that $S(4) \neq \emptyset$. It follows from Claim 2 and Claim 3 that each vertex in $S(4)$ has a neighbour in $S(1, 2)$ and $S(4)$ is anti-complete to $S(2, 3, 4) \cup S(4, 5, 1)$. In particular, $S(1, 2)$ contains a vertex that has a neighbour in $S(4)$. By **(P5)**, this vertex is universal in $S(1, 2)$ and so $S(1, 2)$ is connected. Note that $S_1 = S(4)$ by **(P2)**, **(P8)** and the C_6 -freeness of G , and $S_2 = S(1, 2) \cup S(i, i+1)$ for some $i = 3, 4$ by **(P3)**, **(P9)** and the C_6 -freeness of G . By symmetry, we can assume that $S_2 = S(1, 2) \cup S(3, 4)$. Moreover, $S(i-1, i, i+1)$ is complete to $S(i, i+1, i+2)$ for each $1 \leq i \leq 5$ by Claim 1. If some vertex $x \in S(3, 4)$ has a neighbour $s \in S(4, 5, 1)$, then s is complete to $S(1, 2)$ by **(P15)**. Let $v \in S(4)$ and $y \in S(1, 2)$ be a neighbour of v . Then $4, v, y, s$ induces a C_4 . This shows that $S(3, 4)$ is anti-complete to $S(4, 5, 1)$. Recall that $S(3, 4)$ is anti-complete to $S(4)$ by **(P5)**. Thus, it follows from Claim 4 that each vertex in $S(3, 4)$ has a neighbour in $S(1, 2, 3)$ and $(S(3, 4), S(1, 2, 3))$ is a co-bipartite chain graph if $S(3, 4) \neq \emptyset$.

Let $S'(3, 4, 5)$ be the set of vertices in $S(3, 4, 5)$ that are complete to $S(3, 4)$ and $S''(3, 4, 5) = S(3, 4, 5) \setminus S'(3, 4, 5)$. Then $S''(3, 4, 5)$ is anti-complete to $S(3, 4)$ by the fact that $S(3, 4)$ is a clique and **(P14)**. Then $S'(3, 4, 5) \cup \{4\}$ and $S''(3, 4, 5)$ are homogeneous cliques in G by **(P1)**, **(P4)**, **(P7)** and Theorem 7. So, $S'(3, 4, 5) = \emptyset$ and $|S''(3, 4, 5)| \leq 1$. This implies that $|S(3, 4, 5)| \leq 1$.

(1) $S(1, 2, 3)$ is complete to $S(1, 2)$. By symmetry $S(5, 1, 2)$ is complete to $S(1, 2)$.

Let $s \in S(1, 2, 3)$. Note that there is an edge yz between $S(4)$ and $S(1, 2)$ where $y \in S(1, 2)$ and $z \in S(4)$. We note that s and y are adjacent for otherwise $1, y, z, 4, 3, s$ induces a C_6 . On the other hand, since $S(1, 2)$ is connected, it follows from Claim 3 s is complete to $S(1, 2)$. ■

By **(1)**, $S(5, 1, 2) \cup \{1\}$ is a homogeneous clique in G and so $S(5, 1, 2) = \emptyset$. Recall that $S(3, 4)$ is a clique if it is not empty and so no vertex in $S(2, 3, 4) \cup S(3, 4, 5)$ can distinguish two vertices in $S(3, 4)$ by **(P14)**. It then follows from **(P1)**, **(P3)**, **(P5)**, **(P6)**, **(P7)** and **(1)** that $(S(1, 2, 3), S(3, 4))$ is a homogeneous pair of cliques in G (if $S(3, 4) = \emptyset$ this means that $S(1, 2, 3)$ is a homogeneous clique).

(2) *Each vertex in $S(1, 2)$ is anti-complete to either $S(4)$ or $S(2, 3, 4) \cup S(4, 5, 1)$.*

Suppose that $d \in S(1, 2)$ has a neighbour $v \in S(4)$ and $s \in S(2, 3, 4) \cup S(4, 5, 1)$. By Claim 3, v and s are not adjacent. But now $4, v, d, s$ induces a C_4 , a contradiction. ■

We let $X = \{v \in S(1, 2) : v \text{ has a neighbour in } S(4)\}$. We let $Y = \{v \in S(1, 2) : v \text{ has a neighbour in } S(2, 3, 4) \cup S(4, 5, 1)\}$, and $Z = S(1, 2) \setminus (X \cup Y)$. By (2), X , Y and Z form a partition of $S(1, 2)$. Note that X is anti-complete to $S(2, 3, 4) \cup S(4, 5, 1)$, Y is anti-complete to $S(4)$ and Z is anti-complete to $S(2, 3, 4) \cup S(4, 5, 1) \cup S(4)$. By (P5), each vertex in X is universal in $S(1, 2)$. By (P12), Y is complete to S_5 . Suppose that $y \in Y$ is not adjacent to $z \in Z$. Let $s \in S(2, 3, 4) \cup S(4, 5, 1)$ be a neighbour of y . By symmetry, we assume that $s \in S(2, 3, 4)$. Recall that $S(4)$ contains a vertex, say v . Then $z, 1, y, s, 4, v$ induces a P_6 . This shows that Y is complete to Z . Suppose that Y contains a vertex y' that has a neighbour $s \in S(4, 5, 1)$ and a vertex y'' that has a neighbour $t \in S(2, 3, 4)$. Then s (resp. t) is not adjacent to y'' (resp. y') by (P13). Then $C \cup \{s, t, y', y''\}$ contains an induced P_6 or an induced F_2 , depending on whether y' and y'' are adjacent. Thus, either each vertex in Y is anti-complete to $S(4, 5, 1)$ and has a neighbour in $S(2, 3, 4)$ or each vertex in Y is anti-complete to $S(2, 3, 4)$ and has a neighbour in $S(4, 5, 1)$. Then Y is a clique by a similar argument in Claim 4 asserting that there is a vertex in $S(2, 3, 4)$ or $S(4, 5, 1)$ that is complete to Y .

Recall that $S(3, 4)$ is anti-complete to $S(4, 5, 1)$. Let $R \subseteq S(2, 3, 4)$ be the set of vertices that are complete to $S(3, 4)$ and $T = S(2, 3, 4) \setminus R$. By (P10), T is anti-complete to $S(3, 4)$. If Y is anti-complete to $S(2, 3, 4)$, then $R \cup \{3\}$ and T are homogeneous cliques in G . Hence, $|S(2, 3, 4)| \leq 1$. Moreover, $(Y, S(4, 5, 1))$ is a homogeneous pair of cliques in G by (1). If Y is anti-complete to $S(4, 5, 1)$, then $S(4, 5, 1) \cup \{5\}$ is a homogeneous clique in G and so $S(4, 5, 1) = \emptyset$. If $S(3, 4) = \emptyset$, then $(Y, S(2, 3, 4))$ is a homogeneous pair of cliques in G . Otherwise let $d \in S(3, 4)$. We claim that T is anti-complete to Y . Suppose by contradiction that $y \in Y$ and $t \in T$ are adjacent. Then d and t are not adjacent by the definition of T . Note that $S(4)$ contain a vertex z and z has a neighbour $x \in X \subseteq S(1, 2)$. Note that x and y are adjacent and y is not adjacent to z and t is not adjacent to x by (2). But now $d, 3, t, y, x, z$ induces a P_6 . So, T is anti-complete to Y . This implies that (Y, R) is a homogeneous pair of cliques in G , and T is a homogeneous clique in G and so $|T| \leq 1$. In either case, $Y \cup S(2, 3, 4) \cup S(4, 5, 1)$ is partitioned into a subset of size at most 1 and a homogeneous pair of cliques in G .

We next deal with $S_5 \cup S(4) \cup X \cup Z$. Let $S_5'' = \{u \in S_5 : u \text{ has a non-neighbour in } X\}$ and $S_5' = S_5 \setminus S_5''$. Then S_5' is complete to X .

(3) *Each vertex in Z has a neighbour in S_5'' .*

Let $Z' = \{z \in Z : z \text{ has a neighbour in } S_5''\}$ and $Z'' = Z \setminus Z'$. Suppose that Z'' contains a vertex a and let A be the connected component of Z'' containing a . Thus, $N(A) \subseteq \{1, 2\} \cup X \cup Y \cup S(1, 2, 3) \cup S_5' \cup Z'$. Recall that Y is complete to S_5 and so $\{1, 2\} \cup X \cup Y \cup S(1, 2, 3) \cup S_5'$ is a clique by (1). Moreover, Y is complete to Z and so Z' is complete to $\{1, 2\} \cup X \cup Y \cup S(1, 2, 3)$. We now show that $N(A)$ is a clique. First, each vertex in Z' is either complete or anti-complete to A . Suppose not. Let $z' \in Z'$ distinguish an edge aa' in A , say z' is adjacent to a but not to a' . Let $u \in S_5''$ be a neighbour of z' and $x \in X$ be a non-neighbour of u . Then x has a neighbour $v \in S(4)$. By (P11), u is not adjacent to v . So, $a', a, z', u, 4, v$ induces a P_6 . Second, let $u \in S_5'$ and $z' \in Z'$ be in $N(A)$. We may choose $a \in A$ to be a

neighbour of u . By the definition of Z' , z' has a neighbour $u'' \in S_5''$. Moreover, u'' is not adjacent to a by the definition of Z'' . Thus, u and z' must be adjacent for otherwise a, z', u'', u induces a C_4 . This shows that Z' and S_5' are complete. Thirdly, let $z_1, z_2 \in N(A) \cap Z'$. Suppose that z_1 and z_2 are not adjacent. Let $u_i \in S_5''$ be a neighbour of z_i and $x_i \in X$ be a non-neighbour of u_i . By the definition of X , x_i has a neighbour $v_i \in S(4)$. By **(P11)**, u_i is not adjacent to v_i . Note that $u_1 \neq u_2$ for otherwise a, z_1, u_1, z_2 induces a C_4 . This means that u_1 (resp. u_2) is not adjacent to z_2 (resp. z_1). Now z_2, a, z_1, u_1, v_1 induces a P_6 . This shows that $N(A)$ is a clique and so a clique cutset of G . Since G has no clique cutsets, $Z'' = \emptyset$. ■

(4) Z is complete to S_5' .

Let $z \in Z$ be an arbitrary vertex. Then z has a neighbour $u \in S_5''$ which has a non-neighbour $x \in X$. If $u' \in S_5'$ is not adjacent to z , then u, u', x, z induces a C_4 . ■

Since S_5' is complete to X and each vertex in $S(4)$ has a neighbour in X , S_5' is complete to $S(4)$ by **(P11)**. So, (4) implies that each vertex in S_5' is a universal vertex in G (note that if $S(3, 4) \neq \emptyset$ then it is also complete to S_5). Therefore, $S_5' = \emptyset$ and $S_5 = S_5''$. We have shown that each vertex in S_5 has a non-neighbour in X and each vertex in Z has a neighbour in S_5 . We proceed with partitioning X , S_5'' and Z into subsets so that we can decompose G into homogeneous pairs of sets. Let $X_0 = \{x \in X : x \text{ is anti-complete to } S_5''\}$ and $X_1 = X \setminus X_0$. We then partition S_5'' into two subsets $S_5''' = \{u \in S_5'' : u \text{ is complete to } X_1\}$ and $R = S_5'' \setminus S_5'''$. Let $Z' = N(R) \cap Z$ and $Z'' = Z \setminus Z'$.

(5) $N(X_0) \cap S(4)$ is anti-complete to X_1 .

Suppose not. Let $v \in S(4) \cap N(X_0)$ have a neighbour $x_1 \in X_1$. Let $x_0 \in X_0$ be a neighbour of v . By the definition of X_1 , x_1 has a neighbour $u \in S_5''$. Note that u is not adjacent to x_0 and so not adjacent to v by **(P11)**. But now $x_1, v, 4, u$ induces a C_4 , a contradiction. ■

Recall that X is a clique. Therefore, $S(4) \setminus N(X_0)$ and $S(4) \cap N(X_0)$ are anti-complete by (5) and the facts that G is C_4 -free and each vertex in $S(4)$ has a neighbour in X . In addition, $S(4) \cap N(X_0)$ is anti-complete to S_5'' by **(P11)**. Therefore, $(X_0, N(X_0) \cap S(4))$ is a homogeneous pair of sets in G .

(6) No vertex in S_5'' can have two non-adjacent neighbours in Z .

Let $u \in S_5''$ have two non-adjacent neighbours z_1 and z_2 in Z . Then u has a non-neighbour $x \in X$. Recall that x is universal in $S(1, 2)$ and so complete to $\{z_1, z_2\}$. Now u, z_1, x, z_2 induces a C_4 . ■

(7) Z' and Z'' are complete.

Suppose not. Let $z' \in Z'$ and $z'' \in Z''$ be non-adjacent. By the definition of Z' , z' has a neighbour $r \in R \subseteq S_5''$. Let $u \in S_5'''$ be a neighbour of z'' . By the definition of R , r is not adjacent to some vertex $x_1 \in X_1$. Since X is complete to Z , x_1 is adjacent to z' and z'' . Moreover, x_1 is adjacent to u by the definition of X_1 . By (6), u is not adjacent to z' . But now x_1, z', r, u induces a C_4 . ■

(8) Z' is complete to S_5''' .

Suppose not. Let $z' \in Z'$ and $u \in S_5'''$ be non-adjacent. By the definition of Z' ,

z' has a neighbour $r \in R$, and r is not adjacent to some vertex $x_1 \in X_1$. Since X is complete to Z , x_1 is adjacent to z' . Moreover, u is adjacent to x_1 and r . So, u, r, z', x_1 induces a C_4 , a contradiction. ■

Since each vertex in $S(4) \setminus N(X_0)$ has a neighbour in X_1 , S_5''' is complete to $S(4) \setminus N(X_0)$ by **(P11)**. It follows from **(7)** and **(8)** that (S_5''', Z'') is a homogeneous pair of sets in G . Moreover, $(R, X_1 \cup Z', S(4) \setminus N(X_0))$ is a homogeneous triple in G by the structural properties we have proved so far. Therefore, $V(G)$ is partitioned into a subset V_0 consisting of vertices in $C \cup S(3, 4, 5)$ and at most one vertex in $Y \cup S(2, 3, 4) \cup S(4, 5, 1)$ (and so of size at most 7), four homogeneous pairs of sets $(S(1, 2, 3), S(3, 4))$, one contained in $Y \cup S(2, 3, 4) \cup S(4, 5, 1)$, $(X_0, N(X_0) \cap S(4))$ and (S_5''', Z'') , and a homogeneous triple $R \cup (X_1 \cup Z') \cup (S(4) \setminus N(X_0))$.

We now show that each of the four homogenous pairs and the homogenous triple has a nice 4-expression and 6-expression, respectively. First of all, $(S(1, 2, 3), S(3, 4))$ and the homogeneous pair of sets contained in $Y \cup S(2, 3, 4) \cup S(4, 5, 1)$ are homogeneous pairs of cliques and so have a nice 4-expression by Lemma 4.

Recall that each vertex in Z has a neighbour in S_5'' . Suppose that $P = u, z_1, z_2, z_3$ is an induced P_4 where $u \in S_5''$ and $z_1, z_2, z_3 \in Z$. Then u has a non-neighbour $x \in X$, and x has a neighbour $v \in S(4)$. By **(P11)**, u and v are not adjacent. Now $v, 4, P$ induces a P_6 . Suppose now that $P = z_1 z_2 z_3 z_4$ be an induced P_4 in Z . Then z_1 has a neighbour $u \in S_5''$ which has a non-neighbour $x \in X$. By **(6)**, u is not adjacent to z_3 or z_4 . Now $P \cup \{u\}$ contains such a labeled P_4 . Therefore, $G[S_5'' \cup Z]$ with the partition (S_5'', Z) satisfies all the conditions in Lemma 5 and so has a nice 4-expression. Since $G[S_5''' \cup Z'']$ is an induced subgraph of $G[S_5'' \cup Z]$, it follows that $G[S_5''' \cup Z'']$ has a nice 4-expression.

(9) Z' is a clique.

Suppose that $z_1, z_2 \in Z'$ are not adjacent. Then z_i has a neighbour $r_i \in R$ for $i = 1, 2$. By **(6)**, $r_1 \neq r_2$ and r_1 (resp. r_2) is not adjacent to z_2 (resp. z_1). Recall that r_1 has a non-neighbour $x \in X_1$. Since x is adjacent to z_1 and z_2 , x is not adjacent to r_2 . Now by the definition of X_1 , x has a neighbour $u \in S_5''$. Then u is adjacent to z_i for otherwise x, z_i, r_i, u induces a C_4 for $i = 1, 2$. This, however, contradicts **(6)**. ■

Claim 5. $G[S_5 \cup (X \cup Z') \cup S(4)]$ has a nice 6-expression.

Proof of Claim 5. Let $X' = X \cup Z'$. By **(9)**, (S_5, X') is a co-bipartite chain graph. Thus, we can order the vertices in S_5 as u_1, \dots, u_r and the vertices in X' as x_1, \dots, x_s such that $N_{X'}(u_i) = \{x_1, \dots, x_j\}$ for some $0 \leq j \leq s$ and $N_{X'}(u_1) \subseteq N_{X'}(u_2) \subseteq \dots \subseteq N_{X'}(u_r)$. Let U_0 be the set of vertices in S_5 that are anti-complete to X' . Let U_1 be the set of vertices in $S_5 \setminus U_0$ that have the smallest neighbourhood in X' , and U_2 be the set of vertices in $S_5 \setminus U_0$ that have the second smallest neighbourhood in X' , and so on. Then S_5 is partitioned into U_0, U_1, \dots, U_q where $q+1$ is the number of different neighbourhoods of vertices in S_5 . Let $X_1 = N_{X'}(U_1)$ and $X_i = N_{X'}(U_i) \setminus N_{X'}(U_{i-1})$ for $2 \leq i \leq q$. Let X_{q+1} be the set of vertices in X' that are anti-complete to S_5 . Note that X_1, \dots, X_q, X_{q+1} partition X' . Let $M_i = N(X_i) \cap S(4)$ for each $1 \leq i \leq q+1$. Since $S(4)$ is anti-complete to Z , some M_i may be empty. Since each vertex in $S(4)$ has a neighbour in X , $S(4) = M_1 \cup \dots \cup M_{q+1}$. We say that $G[X_j \cup M_j]$ is a *piece* for each $1 \leq i \leq q+1$. Note that any vertex in M_j has a neighbour in $X_j \cap X$ since Z is anti-complete to $S(4)$.

(i) U_0 is anti-complete to $X' \cup S(4)$.

By the definition of U_0 , U_0 is anti-complete to X' . Since any vertex in $S(4)$ has a neighbour in X , U_0 is anti-complete to $S(4)$ by **(P11)**. ■

(ii) For $1 \leq i \leq q$, U_i is complete to $X_j \cup M_j$ for $1 \leq j \leq i$.

Let $u \in U_i$ and $1 \leq j \leq i$. Then u is complete to X_j by the definition of X_j . If M_j is empty, then the proof is complete. So, we assume that $M_j \neq \emptyset$. For each $v \in M_j$, v has a neighbour in $X_j \cap X$. Since u is complete to X_j , u is adjacent to v by **(P11)**. This completes the proof. ■

(iii) For each $1 \leq i \leq q$, U_i is anti-complete to $X_j \cup M_j$ for $i < j \leq q + 1$.

Let $u \in U_i$ and let j be such that $i < j \leq q + 1$. Then u is anti-complete to X_j by the definition of X_j . If M_j is empty, then the proof is complete. So, we assume that $M_j \neq \emptyset$. For each $v \in M_j$, v has a neighbour in $X_j \cap X$. Since u is anti-complete to X_j , u is not adjacent to v by **(P11)**. ■

(iv) For $1 \leq i, j \leq q + 1$ with $i \neq j$, $M_i \cap M_j = \emptyset$.

Suppose that $i < j$ and $v \in M_i \cap M_j$. Then v has a neighbour $x_i \in X_i$ and a neighbour $x_j \in X_j$. Let $u_i \in U_i$. By **(ii)**, u_i is adjacent to v , and by **(iii)** u_i is not adjacent to v . This is a contradiction. ■

(v) For $1 \leq i, j \leq q + 1$ with $i \neq j$, M_i and M_j are anti-complete.

Suppose that $v_i \in M_i$ and $v_j \in M_j$ are adjacent. Then v_i has a neighbour $x_i \in X_i$ and v_j has a neighbour $x_j \in X_j$. By **(iv)**, v_i is not adjacent to x_j and v_j is not adjacent to x_i . Then v_i, x_i, x_j, v_j induces a C_4 . ■

Since no vertex in $S(1, 2)$ can have two non-adjacent neighbours in $S(4)$, each piece $G[X_i \cup M_i]$ with the partition (X_i, M_i) satisfies all the conditions in Lemma 5 and so there is a nice 4-expression τ_i for it where all vertices in X_i have label 2 and all vertices in M_i have label 4. For $0 \leq i \leq q$, let ϵ_i be a 2-expression for U_i in which all vertices in U_i have label 5. We now construct a nice 6-expression for $G[S_5 \cup (X \cup Z') \cup S(4)]$. Let $\sigma_1 = \rho_{5 \rightarrow 6}(\eta_{5,4}(\eta_{5,2}(\epsilon_1 \oplus \tau_1)))$. For each $2 \leq i \leq q$, let

$$\sigma_i = \rho_{5 \rightarrow 6}(\eta_{5,6}(\eta_{5,4}(\eta_{5,2}(\epsilon_i \oplus \rho_{1 \rightarrow 2}(\eta_{1,2}(\sigma_{i-1} \oplus \rho_{2 \rightarrow 1}(\tau_i))))))).$$

Then $\sigma = \eta_{5,6}(\epsilon_0 \oplus (\rho_{1 \rightarrow 2}(\eta_{1,2}(\sigma_q \oplus \rho_{2 \rightarrow 1}(\tau_{q+1}))))))$ is a nice 6-expression for $G[S_5 \cup (X \cup Z') \cup S(4)]$ (the correctness of the construction follows from **(i)-(v)**). This completes the proof of the claim. ■

Since $(X_0, N(X_0) \cap S(4))$ and $R \cup (X_1 \cup Z') \cup (S(4) \setminus N(X_0))$ induce subgraphs of $G[S_5 \cup (X \cup Z') \cup S(4)]$, the pair has a nice 4-expression and the triple has a nice 6-expression, by Claim 5. Finally, $\text{cw}(G) \leq |V_0| + 2 \times 4 + 3 = 18$. This completes our proof of the lemma. □

We are now ready to prove our main theorem.

Proof of Theorem 6. Let G be a (C_4, P_6) -free atom. Let G' be the graph obtained from G by removing twin vertices and universal vertices. It follows from Lemma 7–Lemma 10

that if G' contains an induced C_5 or C_6 , then G' has clique-width at most 18. Therefore, we can assume that G' is also (C_5, C_6) -free and therefore G' is chordal. It then follows from a well-known result of Dirac [21] that G' is a clique whose clique-width is 2. Finally, $\text{cw}(G) = \text{cw}(G')$ by Lemma 3 and this completes the proof. \square

5 The Hardness Result

In this section, we prove that COLOURING is NP-complete on the class of $(C_4, 3P_3, P_3 + P_6, 2P_5, P_9)$ -free graphs. A graph is a *split graph* if its vertex set can be partitioned into two disjoint sets C and I such that C is a clique and I is an independent set. The pair (C, I) is called a *split partition* of G . A split graph is *complete* if it has a *complete* split partition, that is, a partition (C, I) such that C and I are complete to each other. It is straightforward to check that a complete split graph is P_4 -free. A *list assignment* of a graph $G = (V, E)$ is a function L that prescribes, for each $u \in V$, a finite list $L(u) \subseteq \{1, 2, \dots\}$ of colours for u . The *size* of a list assignment L is the maximum list size $|L(u)|$ over all vertices $u \in V$. A colouring c *respects* L if $c(u) \in L(u)$ for all $u \in V$. The LIST COLOURING problem is to decide whether a given graph G has a colouring c that respects a given list assignment L .

For our hardness reduction we need the following result.

Lemma 11 ([27]). *LIST COLOURING is NP-complete even for complete split graphs with a list assignment of size at most 3.*

We are now ready to prove our theorem. In its proof we construct a C_4 -free graph G' that is neither $(sP_2 + P_8)$ -free nor $(sP_2 + P_4 + P_5)$ -free for any $s \geq 0$. Hence, a different construction is needed for tightening our hardness result (if possible).

Theorem 8. *COLOURING is NP-complete for $(C_4, 3P_3, P_3 + P_6, 2P_5, P_9)$ -free graphs.*

Proof. We reduce from LIST COLOURING on complete split graphs with a list assignment of size at most 3. This problem is NP-complete due to Lemma 11. Let G be a complete split graph with a list assignment L of size at most 3. From (G, L) we construct an instance (G', k) of COLOURING as follows. First, let $k \leq 3|V(G)|$ be the size of the union of all lists $L(u)$. Let (C, I) be a complete split partition of $V(G)$. We say that the vertices of C and I are of c -type and i -type, respectively. Let G' be the graph of size $O(|V(G)|k)$ obtained from G by adding the following sets of vertices and edges (see Figure 7 for an illustration):

- Take a clique X on k vertices x_1, \dots, x_k . We say that these vertices are of x -type.
- For each $u \in V(G)$, introduce a clique Y_u of size $k - |L(u)|$ such that every vertex of Y_u is adjacent to u and also to every x_i with $i \in L(u)$ (so, each vertex in every Y_u is adjacent to exactly one vertex of $V(G)$, namely vertex u). We say that the vertices in every Y_u are of y -type.

We now prove two claims that together with NP-membership of COLOURING imply the theorem.

Claim 1. *The graph G has a colouring that respects L if and only if G' has a k -colouring.*




Figure 7: Illustration of the NP-hardness construction for $k = 4$.

Proof of Claim 1. First suppose that G has a colouring c that respects L . We give each x_i colour i . Consider a clique Y_u . By construction, every vertex in every Y_u can only be assigned a colour from $\{1, \dots, k\} \setminus L(u)$. As $|Y_u| = k - |L(u)|$ and Y_u is a clique, we need all of these colours. As every vertex in Y_u has u as its only neighbour in G and $c(u)$ belongs to L , this is possible. Hence, we can extend c to a k -colouring c' of G' . Now suppose that G' has a k -colouring c' . As X is a clique, we may assume without loss of generality that $c'(x_i) = i$ for $i = 1, \dots, k$. Then c colours the vertices of each Y_u with colours from $\{1, \dots, k\} \setminus L(u)$. As Y_u is a clique of size $k - |L(u)|$, all these colours appear as a colour of a vertex in Y_u . This means that u must get a colour from $L(u)$. Hence the restriction of c' to G yields a colouring c that respects L . This completes the proof of Claim 1. ■

Claim 2. *The graph G' is $(C_4, 3P_3, P_3 + P_6, 2P_5, P_9)$ -free.*

Proof of Claim 2. We first prove that G' is C_4 -free. For contradiction, suppose that G' contains an induced C_4 on vertices a_1, a_2, a_3, a_4 in that order. Then at least one of a_1, a_2, a_3, a_4 , say a_1 , is of y -type, since the y -type vertices separate K from G , and both K and G are C_4 -free. If a_2 and a_4 both belong to $X \cup Y_u$ or both $\{u\} \cup Y_u$, then they are adjacent, which is not possible. Hence, one of them, say a_2 , belongs to X and the other one, a_4 , is equal to u . This is not possible either, as in that case a_3 must be of y -type and any two y -type neighbours of a u -type vertex are adjacent. We conclude that G' is C_4 -free.

We now prove that G' is $3P_3$ -free. For contradiction, suppose that G' contains an induced $3P_3$. At most one of the three connected components of the induced $3P_3$ can contain an x -type vertex. Then the other two connected components contain no x -type vertex. As the y -type neighbours of an y -type vertex form a clique together with their neighbour of G , both these other connected components contain at least two vertices from G . Then all these vertices must be of i -type, as c -type vertices are adjacent to every other vertex of G . However, i -type vertices form an independent set and only share c -type vertices as common neighbours, a contradiction. We conclude that G' is $3P_3$ -free.

We now prove that G' is $(P_3 + P_6)$ -free. For contradiction, suppose that G' contains an induced $P_3 + P_6$. Let F_1 be the P_3 -component and F_2 be the P_6 -component. Suppose

F_1 contains an x -type vertex. Then F_2 contains no x -type vertex. As every $\{u\} \cup Y_u$ is a clique and i -type vertices form an independent set, F_2 must contain at least one c -type vertex. As c -type vertices are adjacent to all i -type vertices and all other c -type vertices, F_2 contains at most three vertices from G which form a subpath of F_2 . As F_2 contains six vertices and every $\{u\} \cup Y_u$ is a clique, this is not possible. Hence F_1 contains no x -type vertex. As every $\{u\} \cup Y_u$ is a clique, this means that F_1 contains at least two adjacent vertices of G . As i -type vertices form an independent set, one of these vertices is of c -type. This means that F_2 contains no vertices of G . This is not possible as the x -type and y -type vertices induce a P_5 -free graph. We conclude that G' is $(P_3 + P_6)$ -free.

We now prove that G' is $2P_5$ -free. For contradiction, suppose that G' contains an induced $2P_5$ with connected components F_1 and F_2 . At most one of F_1, F_2 may contain an x -type vertex. Hence we may assume that F_2 contains no x -vertex. This means that F_2 must be of the form $y - i - c - i - y$. As a consequence, F_1 only contains vertices of x -type or y -type. This is not possible as those vertices induce a P_5 -free subgraph. We conclude that G' is $2P_5$ -free.

Finally we prove that G' is P_9 -free. Let P be a maximal induced path of G' . First suppose that P contains at least two i -type vertices. Then P contains a subpath of the form $i - c - i$ or $i - y - x - x - y - i$. We can extend $i - c - i$ to at most an 8-vertex path, which is of the form $y - i - c - i - y - x - x - y$, but we cannot extend $i - y - x - x - y - i$ any further. Now suppose that P contains exactly one i -type vertex. If P contains no c -type vertex, then P is a 5-vertex path of the form $i - y - k - k - y$. Otherwise P contains exactly one c -type vertex. In that case P is a 7-vertex path of the form $y - c - i - y - x - x - y$ or $y - x - x - y - c - i - y$. Now suppose that P has no i -type vertex. If P has no c -type vertex either, then P is of the form $y - x - y$ or $y - x - x - y$, so P has at most five vertices. If P has exactly one c -type vertex, then P is a 5-vertex path of the form $c - y - k - k - y$. Otherwise, P has exactly two c -type vertices. In that case P is a 7-vertex path of the form $y - c - c - y - k - k - y$. We conclude that G' is P_9 -free. ■

The result now follows from Claim 1 and Claim 2. □

6 Conclusions

We proved that COLOURING restricted to (C_4, P_t) -free graphs is polynomial-time solvable for $t \leq 6$ and NP-complete for $t \geq 9$. Combined with the aforementioned known results from [7, 37, 43], we can replace Theorem 2 by the following almost complete dichotomy for COLOURING restricted to (C_s, P_t) -free graphs.

Theorem 9. *Let $s \geq 3$ and $t \geq 1$ be two fixed integers. Then COLOURING for (C_s, P_t) -free graphs is polynomial-time solvable if $s = 3, t \leq 6$, or $s = 4, t \leq 6$, or $s \geq 5, t \leq 4$, and NP-complete if $s = 3, t \geq 22$, or $s = 4, t \geq 9$, or $s \geq 5, t \geq 5$.*

We proved that, in contrast to (C_4, P_6) -free graphs, (C_4, P_6) -free atoms have bounded clique-width. As we also showed that the classification of boundedness of clique-width of H -free graphs and H -free atoms coincides, this result was not expected beforehand. As such, we believe that a systematic study in the applicability of this technique, together with the other techniques developed in our paper, can be used to prove further polynomial-time results for COLOURING. For future work we aim to complete the classification of Theorem 9.

The natural candidate class for a polynomial-time result of COLOURING is the class of (C_4, P_7) -free graphs. However, this may require significant efforts for the following reason. Lozin and Malyshev [46] determined the complexity of COLOURING for \mathcal{H} -free graphs for every finite set of graphs \mathcal{H} consisting only of 4-vertex graphs except when \mathcal{H} is $\{K_{1,3}, 4P_1\}$, $\{K_{1,3}, 2P_1 + P_2\}$, $\{K_{1,3}, 2P_1 + P_2, 4P_1\}$ or $\{C_4, 4P_1\}$. Solving any of these open cases would be considered as a major advancement in the area. Since $(C_4, 4P_1)$ -free graphs are (C_4, P_7) -free, polynomial-time solvability of COLOURING on (C_4, P_7) -free graphs implies polynomial-time solvability for COLOURING on $(C_4, 4P_1)$ -free graphs. As a first step, we aim to apply the techniques of this paper to $(C_4, 4P_1)$ -free graphs.

The class of (C_3, P_7) -free graphs is also a natural class to consider. Interestingly, every (C_3, P_7) -free graph is 5-colourable. This follows from a result of Gravier, Hoàng and Maffray [30] who proved that for any two integers $r, t \geq 1$, every (K_r, P_t) -free graph can be coloured with at most $(t-2)^{r-2}$ colours. On the other hand, 3-COLOURING is polynomial-time solvable for P_7 -free graphs [5]. Hence, in order to solve COLOURING for (C_3, P_7) -free graphs we may instead consider 4-COLOURING for (C_3, P_7) -free graphs. This problem also seems highly nontrivial.

Acknowledgments. Initially we proved NP-hardness of COLOURING for (C_4, P_{16}) -free graphs. Afterwards we were able to improve this result to (C_4, P_9) -free graphs via a simplification of our construction. We would like to thank an anonymous reviewer of the conference version of our paper for pointing out this simplification as well.

Serge Gaspers is the recipient of an Australian Research Council (ARC) Future Fellowship (FT140100048) and acknowledges support under the ARC's Discovery Projects funding scheme (DP150101134). Shenwei Huang is partially supported by the National Natural Science Foundation of China (11801284). Daniël Paulusma is supported by Leverhulme Trust Research Project Grant RPG-2016-258.

References

- [1] Isolde Adler, Ngoc-Khang Le, Haiko Müller, Marko Radovanovic, Nicolas Trotignon, and Kristina Vušković. On rank-width of (diamond, even hole)-free graphs *Discrete Mathematics and Theoretical Computer Science*, 19(1):#24, 2017.
- [2] Aistis Atminas, Vadim V. Lozin and Igor Razgon. Linear time algorithm for computing a small biclique in graphs without long induced paths. *Proc. SWAT 2012, LNCS*, 7357:142–152, 2012
- [3] Alexandre Blanché, Konrad K. Dabrowski, Matthew Johnson, and Daniël Paulusma. Hereditary graph classes: When the complexities of Colouring and Clique Cover coincide. *CoRR*, abs/1607.06757, 2016.
- [4] Valerio Boncompagni, Irena Penev, and Kristina Vušković. Clique-cutsets beyond chordal graphs. *CoRR*, abs/1707.03252, 2017.
- [5] Flavio Bonomo, Maria Chudnovsky, Peter Maceli, Oliver Schaudt, Maya Stein, and Mingxian Zhong. Three-coloring and list three-coloring of graphs without induced paths on seven vertices. *Combinatorica*, (in press).

- [6] Andreas Brandstädt and Chinh T. Hoàng. On clique separators, nearly chordal graphs, and the maximum weight stable set problem. *Theoretical Computer Science*, 389:295–306, 2007.
- [7] Andreas Brandstädt, Tilo Klemmt, and Suhail Mahfud. P_6 - and triangle-free graphs revisited: structure and bounded clique-width. *Discrete Mathematics and Theoretical Computer Science*, 8:173–188, 2006.
- [8] Andreas Brandstädt, Van Bang Le, and Jeremy P. Spinrad. *Graph Classes: A Survey*, volume 3 of *SIAM Monographs on Discrete Mathematics and Applications*. Philadelphia, 1999.
- [9] Hajo Broersma, Petr A. Golovach, Daniël Paulusma, and Jian Song. Determining the chromatic number of triangle-free $2P_3$ -free graphs in polynomial time. *Theoretical Computer Science*, 423:1–10, 2012.
- [10] Kathie Cameron and Chinh T. Hoàng. Solving the clique cover problem on (bull, C_4) -free graphs. *CoRR*, abs/1704.00316, 2017.
- [11] Maria Chudnovsky, Sophie Spirkl, and Mingxian Zhong. Four-coloring P_6 -free graphs. I. Extending an excellent precoloring. *arXiv*, abs/1802.02282, 2018.
- [12] Maria Chudnovsky, Sophie Spirkl, and Mingxian Zhong. Four-coloring P_6 -free graphs. II. Finding an excellent precoloring. *arXiv*, abs/1802.02283, 2018.
- [13] Maria Chudnovsky and Jura Stacho. 3-colorable subclasses of P_8 -free graphs. *SIAM J. Discrete Math.*, 32:1111–1138, 2018.
- [14] Bruno Courcelle, Johann A. Makowsky, and Udi Rotics. Linear time solvable optimization problems on graphs of bounded clique-width. *Theory of Computing Systems*, 33:125–150, 2000.
- [15] Bruno Courcelle and Stephan Olariu. Upper bounds to the clique width of graphs. *Discrete Applied Mathematics*, 101:77–114, 2000.
- [16] Julia Chuzhoy, Improved bounds for the flat wall theorem, *Proc. of SODA 2015* 256–275, 2015.
- [17] Konrad K. Dabrowski, François Dross, and Daniël Paulusma. Colouring diamond-free graphs. *Journal of Computer and System Sciences*, 89:410–431, 2017.
- [18] Konrad K. Dabrowski, Petr A. Golovach, and Daniël Paulusma. Colouring of graphs with Ramsey-type forbidden subgraphs. *Theoretical Computer Science*, 522:34–43, 2014.
- [19] Konrad K. Dabrowski and D. Paulusma, Clique-width of graph classes defined by two forbidden induced subgraphs, *The Computer Journal* 59 (2016) 650–666.
- [20] Konrad K. Dabrowski and Daniël Paulusma. On colouring $(2P_2, H)$ -free and (P_5, H) -free graphs. *Information Processing Letters*, to appear, 134:35–41, 2018.

- [21] Gabriel A. Dirac. On rigid circuit graphs. *Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*, 25:71–76, 1961.
- [22] Thomas Emden-Weinert, Stefan Hougardy, and Bernd Kreuter. Uniquely colourable graphs and the hardness of colouring graphs of large girth. *Combinatorics, Probability and Computing*, 7:375–386, 1998.
- [23] Wolfgang Espelage, Frank Gurski, and Egon Wanke. How to solve NP-hard graph problems on clique-width bounded graphs in polynomial time. *Proc. WG 2001, LNCS*, 2204:117–128, 2001.
- [24] Serge Gaspers and Shenwei Huang. Linearly χ -bounding (P_6, C_4) -free graphs. *Proc. WG 2017, LNCS*, 10520:263–274, 2017.
- [25] Serge Gaspers, Shenwei Huang, and Daniël Paulusma. Colouring square-free graphs without long induced paths, Proc. STACS 2018, Leibniz International Proceedings in Informatics 96, 35:1–35:15, 2018.
- [26] Petr A. Golovach, Matthew Johnson, Daniël Paulusma, and Jian Song. A survey on the computational complexity of coloring graphs with forbidden subgraphs. *Journal of Graph Theory*, 84:331–363, 2017.
- [27] Petr A. Golovach and Daniël Paulusma. List coloring in the absence of two subgraphs. *Discrete Applied Mathematics*, 166:123–130, 2014.
- [28] Petr A. Golovach, Daniël Paulusma, and Jian Song. Coloring graphs without short cycles and long induced paths. *Discrete Applied Mathematics*, 167:107–120, 2014.
- [29] Martin C. Golumbic. *Algorithmic Graph Theory and Perfect Graphs*, volume 57 of *Annals of Discrete Mathematics*. North-Holland Publishing Co., 2004.
- [30] Sylvain Gravier, Chinh T. Hoàng, and Frédéric Maffray. Coloring the hypergraph of maximal cliques of a graph with no long path. *Discrete Mathematics*, 272:285–290, 2003.
- [31] Martin Grötschel, László Lovász, and Alexander Schrijver. Polynomial algorithms for perfect graphs. *Annals of Discrete Mathematics*, 21:325–356, 1984.
- [32] Pavol Hell and Shenwei Huang. Complexity of coloring graphs without paths and cycles. *Discrete Applied Mathematics*, 216, Part 1:211–232, 2017.
- [33] Chinh T. Hoàng, Marcin Kamiński, Vadim V. Lozin, Joe Sawada, and Xiao Shu. Deciding k -colorability of P_5 -free graphs in polynomial time. *Algorithmica*, 57:74–81, 2010.
- [34] Chinh T. Hoàng and D. Adam Lazzarato. Polynomial-time algorithms for minimum weighted colorings of $(P_5, \overline{P_5})$ -free graphs and similar graph classes. *Discrete Applied Mathematics*, 186:106–111, 2015.
- [35] Ian Holyer. The NP-Completeness of edge-coloring. *SIAM Journal on Computing*, 10:718–720, 1981.

- [36] Shenwei Huang. Improved complexity results on k -coloring P_t -free graphs. *European Journal of Combinatorics*, 51:336–346, 2016.
- [37] Shenwei Huang, Matthew Johnson, and Daniël Paulusma. Narrowing the complexity gap for colouring (C_s, P_t) -free graphs. *The Computer Journal*, 58:3074–3088, 2015.
- [38] Tommy R. Jensen and Bjarne Toft. *Graph Coloring Problems*. Wiley Interscience, 1995.
- [39] Marcin Kamiński, Vadim V. Lozin, and Martin Milanič. Recent developments on graphs of bounded clique-width. *Discrete Applied Mathematics*, 157:2747–2761, 2009.
- [40] Thiagarajan Karthick, Frédéric Maffray, and Lucas Pastor. Polynomial cases for the vertex coloring problem. *CoRR*, abs/1709.07712, 2017.
- [41] Thiagarajan Karthick and Frédéric Maffray. Square-free graphs with no six-vertex induced path. *CoRR*, abs/1805.05007, 2018.
- [42] Daniel Kobler and Udi Rotics. Edge dominating set and colorings on graphs with fixed clique-width. *Discrete Applied Mathematics*, 126:197–221, 2003.
- [43] Daniel Král', Jan Kratochvíl, Zsolt Tuza, and Gerhard J. Woeginger. Complexity of coloring graphs without forbidden induced subgraphs. *Proc. WG 2001, LNCS*, 2204:254–262, 2001.
- [44] Daniel Leven and Zvi Galil. NP completeness of finding the chromatic index of regular graphs. *Journal of Algorithms*, 4:35–44, 1983.
- [45] László Lovász. Coverings and coloring of hypergraphs. *Congressus Numerantium*, VIII:3–12, 1973.
- [46] Vadim V. Lozin and Dmitriy S. Malyshev. Vertex coloring of graphs with few obstructions. *Discrete Applied Mathematics*, 216, Part 1:273–280, 2017.
- [47] Vadim V. Lozin and Dieter Rautenbach. On the band-, tree-, and clique-width of graphs with bounded vertex degree. *SIAM Journal on Discrete Mathematics*, 18:195–206, 2004.
- [48] Vadim V. Lozin and Dieter Rautenbach. The tree- and clique-width of bipartite graphs in special classes. *Australasian Journal of Combinatorics*, 34:57–67, 2006.
- [49] Johann A. Makowsky and Udi Rotics. On the clique-width of graphs with few P_4 's. *International Journal of Foundations of Computer Science*, 10:329–348, 1999.
- [50] Dmitriy S. Malyshev. The coloring problem for classes with two small obstructions. *Optimization Letters*, 8:2261–2270, 2014.
- [51] Dmitriy S. Malyshev. Two cases of polynomial-time solvability for the coloring problem. *Journal of Combinatorial Optimization*, 31:833–845, 2016.
- [52] Dmitriy S. Malyshev and O. O. Lobanova. Two complexity results for the vertex coloring problem. *Discrete Applied Mathematics*, 219:158–166, 2017.

- [53] Michael Molloy and Bruce Reed. *Graph Colouring and the Probabilistic Method*. Springer-Verlag Berlin Heidelberg, 2002.
- [54] Sang-Il Oum and Paul D. Seymour. Approximating clique-width and branch-width. *Journal of Combinatorial Theory, Series B*, 96:514–528, 2006.
- [55] Michaël Rao. MSOL partitioning problems on graphs of bounded treewidth and clique-width. *Theoretical Computer Science*, 377:260–267, 2007.
- [56] David Schindl. Some new hereditary classes where graph coloring remains NP-hard. *Discrete Mathematics*, 295:197–202, 2005.
- [57] D. Seinsche. On a property of n -colorable graphs. *Journal of Combinatorial Theory, Series B*, 16:191–193, 1974.
- [58] Robert E. Tarjan. Decomposition by clique separators. *Discrete Mathematics*, 55:221–232, 1985.

A The Proof of Lemma 4

Here is a proof for Lemma 4, which we restate below.

Lemma 4 [Folklore]. *There is a nice 4-expression for any co-bipartite chain graph.*

Proof. Let $G = (A \uplus B, E)$ be a co-bipartite chain graph where A and B are cliques. Since G is a co-bipartite chain graph, we can order the vertices in A as a_0, a_1, \dots, a_s and the vertices in B as b_1, \dots, b_t such that for each $0 \leq i \leq s$, $N_B(a_i) = \{b_1, \dots, b_j\}$ for some $0 \leq j \leq t$ ($j = 0$ means that $N_B(a_i) = \emptyset$) and $N_B(a_0) \subseteq N_B(a_1) \subseteq \dots \subseteq N_B(a_t)$. Note that two vertices in A are twins in G if and only if they have the same neighbours in B . It follows from Lemma 3 that twin vertices do not change the clique-width. Neither do they change the niceness of the clique-width expression. Therefore, we may assume that for each $0 \leq j \leq t$ there is at most one a_i with $N_B(a_i) = \{b_1, \dots, b_j\}$. Moreover, we can assume that for each $0 \leq j \leq t$ there is exactly one a_i with $N_B(a_i) = \{b_1, \dots, b_j\}$ for otherwise G would be an induced subgraph of this graph. In other words, $s = t + 1$ and $N_B(a_i) = \{b_1, \dots, b_i\}$ for each $0 \leq i \leq t$. Let $\tau_1 = \rho_{3 \rightarrow 4}(\rho_{1 \rightarrow 2}(\eta_{1,3}(1(a_1) \oplus 3(b_1))))$. For each $2 \leq i \leq t$, note that

$$\tau_i = \rho_{1 \rightarrow 2}(\eta_{1,4}(\rho_{3 \rightarrow 4}(\eta_{3,4}(\eta_{1,2}((1(a_i) \oplus 3(b_i)) \oplus \tau_{i-1}))))))$$

is a nice 4-expression for $G[\{a_1, \dots, a_i, b_1, \dots, b_i\}]$. Now $\tau = \rho_{1 \rightarrow 2}(\eta_{1,2}(1(a_0) \oplus \tau_t))$ is a nice 4-expression for G . \square