

Durham Research Online

Deposited in DRO:

17 December 2019

Version of attached file:

Published Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Boegli, Sabine and Brown, B. Malcolm and Marletta, Marco and Tretter, Christiane and Wagenhofer, Markus (2014) 'Guaranteed resonance enclosures and exclosures for atoms and molecules.', *Proceedings of the Royal Society A : mathematical, physical and engineering sciences.*, 470 (2171). p. 20140488.

Further information on publisher's website:

<https://doi.org/10.1098/rspa.2014.0488>

Publisher's copyright statement:

© 2014 The Authors. Published by the Royal Society under the terms of the Creative Commons Attribution License <http://creativecommons.org/licenses/by/4.0/>, which permits unrestricted use, provided the original author and source are credited.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Research



Cite this article: Bögli S, Brown MB, Marletta M, Tretter C, Wagenhofer M. 2014 Guaranteed resonance enclosures and exclosures for atoms and molecules. *Proc. R. Soc. A* **470**: 20140488. <http://dx.doi.org/10.1098/rspa.2014.0488>

Received: 20 June 2014

Accepted: 1 September 2014

Subject Areas:

computational mathematics, differential equations, quantum physics

Keywords:

resonance, eigenvalue, complex potential, interval arithmetic

Author for correspondence:

Marco Marletta

e-mail: marlettam@cardiff.ac.uk

Guaranteed resonance enclosures and exclosures for atoms and molecules

Sabine Bögli¹, B. Malcolm Brown², Marco Marletta³,
Christiane Tretter^{4,5} and Markus Wagenhofer⁶

¹Mathematisches Institut, Universität Bern, Alpeneggstrasse 22, 3012 Bern, Switzerland

²School of Computer Science, Cardiff University, 5 The Parade, Cardiff CF24 3AA, UK

³School of Mathematics, Cardiff University, 21–23 Senghennydd Road, Cardiff CF24 4AG, UK

⁴Mathematisches Institut, Universität Bern, Sidlerstrasse 5, 3012 Bern, Switzerland

⁵Matematiska institutionen, Stockholms universitet, 10691 Stockholm, Sweden

⁶Im Gallmoos 11, 93128 Regenstein, Germany

In this paper, we confirm, with absolute certainty, a conjecture on a certain oscillatory behaviour of higher auto-ionizing resonances of atoms and molecules beyond a threshold. These results not only definitely settle a more than 30 year old controversy in Rittby *et al.* (1981 *Phys. Rev. A* **24**, 1636–1639 (doi:10.1103/PhysRevA.24.1636)) and Korsch *et al.* (1982 *Phys. Rev. A* **26**, 1802–1803 (doi:10.1103/PhysRevA.26.1802)), but also provide new and reliable information on the threshold. Our interval-arithmetic-based method allows one, for the first time, to *enclose* and to *exclude* resonances with guaranteed certainty. The efficiency of our approach is demonstrated by the fact that we are able to show that the approximations in Rittby *et al.* (1981 *Phys. Rev. A* **24**, 1636–1639 (doi:10.1103/PhysRevA.24.1636)) *do* lie near true resonances, whereas the approximations of higher resonances in Korsch *et al.* (1982 *Phys. Rev. A* **26**, 1802–1803 (doi:10.1103/PhysRevA.26.1802)) *do not*, and further that there exist two new pairs of resonances as suggested in Abramov *et al.* (2001 *J. Phys. A* **34**, 57–72 (doi:10.1088/0305-4470/34/1/304)).

© 2014 The Authors. Published by the Royal Society under the terms of the Creative Commons Attribution License <http://creativecommons.org/licenses/by/4.0/>, which permits unrestricted use, provided the original author and source are credited.

1. Introduction

Reliable and precise information on the location of resonances is very hard to obtain. While numerical approximations are widely used in physics, so far there has been no way to show that they produce results *near*, or *not near*, true resonances. The reason is that computations of complex eigenvalues in the presence of continuous spectrum are not backed up by any convergence results. This paper presents a new method that, for the first time, permits one to locate resonances with absolute certainty and high accuracy and, at the same time, to show that numerical approximations *fail* to lie near true resonances. We provide new and reliable information on the oscillatory behaviour of the real parts of certain resonance strings and on the threshold beyond which it occurs.

The key ingredient in our method is interval arithmetic. It allows us to carry out every computational step with absolute accuracy by operating on intervals rather than on numbers. Remarkably, this theoretical idea has had convincing impact in different practical physical applications recently: to control the stability of difficult nonlinear systems in robotics to navigate a sailboat autonomously over a distance of 100 km (see [1]); to perform rigorous global optimization of impulsive planet-to-planet transfer (see [2]) or to rigorously govern the long-term stability in particle accelerators (see [3]).

In this paper, we demonstrate the efficacy of interval approaches for the computation of resonance enclosures and enclosures with absolute certainty. The power of our method is substantiated by the fact that it can be applied to definitely settle a more than 30 year old controversy in [4,5] which could not be resolved by any other method before.

In connection with auto-ionizing resonances of atoms and molecules lying above a ionization threshold, Moiseyev *et al.* [6] studied resonances of the Sturm–Liouville problem

$$-y''(x) + \frac{2\mu}{h^2}((0.5x^2 - 0.8)e^{-x^2/10} + 0.8 - \epsilon)y(x) = 0, \quad x \in \mathbb{R}. \quad (1.1)$$

A first resonance was suggested to lie near $2.124 - 0.0185i$ (with μ/h^2 set to 1); moreover, one bound state was proposed to lie near 0.5. The resonance was found by complex scaling of the self-adjoint Hamiltonian and approximation using a variational principle with 10 real Gauss-type basis functions for the scaled Hamiltonian. Because the latter is no longer self-adjoint, the authors pointed out that further exploration is needed to obtain information on the true position of a nearby resonance.

In the two subsequent papers [4] and the more detailed version [7], Rittby *et al.* combined complex scaling with some Weyl-type analysis and numerical integration methods to compute 44 resonance approximations, including approximations for the first resonance and bound state suggested in [6]; the second resonance therein was further studied by Engdahl & Brändas in [8] by computing lower bounds for norms of Riesz projections. The main conclusion of [4,7] is that there exists a complex threshold ϵ_{thresh} with $\text{Re}(\epsilon_{\text{thresh}}) > 0$, $\text{Im}(\epsilon_{\text{thresh}}) < 0$ such that all resonance approximations of (1.1) satisfy $\text{Re}(\epsilon) \leq \text{Re}(\epsilon_{\text{thresh}}) \sim 4.68$ and beyond this threshold, i.e. for $\text{Im}(\epsilon) < \text{Im}(\epsilon_{\text{thresh}}) < 0$, their real parts exhibit a certain oscillatory behaviour.

Shortly after the publication of [4] and the submission of [7], Korsch *et al.* announced in [5, comment] that they had computed a different set of resonance approximations beyond the threshold which did not exhibit any oscillatory behaviour, whereas their earlier computations of lower resonances in [9] had not shown such a disagreement. They used a complex-rotated Milne method and they believed to have backed up their computations by some WKB approximations. Korsch *et al.* concluded that the results of Rittby *et al.* for higher resonances were incorrect; they conjectured this might be due to numerical instabilities or to the too limited range $0 < \theta < \pi/4$ of angles in the complex scaling method in [4,7].

In an immediate reply (see [10, reply to comment]), Rittby *et al.* [5] defended their results and attributed the discrepancies of the results to wrongly chosen outgoing boundary conditions. They argued that the asymptotic solutions of the complex Riccati equation associated with (1.1) undergo a dramatic change when θ passes the critical value $\theta_{\text{crit}} = \pi/4$ of the potential in (1.1)

and hence the rotation angle $\theta = 50^\circ$ used by Korsch *et al.* was too large. Because of this and the stability of the computations in [4,7] against variations of the rotation angle θ , Rittby *et al.* [4,7] believed to have found approximations to true resonances. About 10 years later, Andersson corroborated the arguments and conclusions of Rittby *et al.* by a careful multiple-transition point WKB analysis and explained the failure of the complex-rotated Milne method of Korsch *et al.* by semi-classical theory in [11].

Almost 20 years after the 1982 dispute, the resonance problem (1.1) was studied as an example in two papers in the mathematical literature. In [12], for more general classes of exponentially decaying potentials, Brown *et al.* developed a resonance-finding procedure for resonances close to points of spectral concentration on the real axis. This method relies on analytic continuation of the Weyl–Titchmarsh function rather than on complex scaling and was first used by Hehenberger *et al.* [13] in numerical computations for the Stark effect. As an example, Brown *et al.* computed approximations to the first three resonances of (1.1) which were very close to the ones found in [7]; note that $\mu/\hbar^2 = 1$ in [7] and that the potential q and spectral parameter λ in [12] are related to the potential V and spectral parameter ϵ in (1.1) by

$$q(x) = (x^2 - 1.6)e^{-x^2/10} = \frac{\hbar^2}{\mu} V(x) - 1.6, \quad \lambda = \frac{2\mu}{\hbar^2}(\epsilon - 0.8).$$

Not long after, Abramov *et al.* [14] proved some global analytical bounds for resonances for various classes of potentials. They combined complex scaling with operator theoretic techniques such as numerical ranges and Birman–Schwinger type arguments. Moreover, for the particular case of (1.1), they also performed numerical computations. The analytical results in [14] supported the conjecture of Rittby *et al.* that a wrong asymptotic boundary condition was used by Korsch *et al.* [5]. The numerical results of [14] reproduced the resonances found in [4,7] and they suggested three pairs of additional resonances. Each pair consists of an even and an odd resonance so close to each other that they could not be computed accurately. These new resonance pairs may be related to the oscillatory behaviour of the real parts; because two of these pairs satisfy $-9.57 \sim \text{Im}(\epsilon_{\text{thresh}}) < \text{Im}(\epsilon) < 0$.

As it was rightly put in [14], none of the above methods for finding resonances can be used to locate them accurately, but there is clear evidence that they exist. Moreover, none of these methods allows for a proof that a numerically computed candidate for a resonance is *not* near any true resonance.

The method presented here permits us to settle both questions definitely and adds new information on the threshold beyond which oscillatory behaviour of the real parts of resonances occurs. We prove that the 44 numerical approximations of resonances from [4,7] do lie near true resonances and that the numerical approximations labelled 16–28 in [5] do *not* lie near true resonances. Moreover, we prove that two of the additional pairs of resonances conjectured in [14] do exist. Our provably correct computations are based on a combination of two key tools, the argument principle on the analytic side and interval arithmetic on the computational side.

Briefly, our approach is as follows. By means of complex scaling $x \rightarrow e^{i\theta} x$ with $\theta \in [0, \pi/4)$, the resonances ϵ of (1.1) are given in terms of the eigenvalues $z = e^{2i\theta}(2\epsilon - 1.6)$ of a Sturm–Liouville problem on \mathbb{R} with complex potential. These eigenvalues can be characterized as the zeros of an analytic function Δ . Hence, their number in a rectangle \mathcal{R}_0 can be counted by means of the argument principle. On the other hand, we can compute the contour integral in the argument principle in interval arithmetic, using a code based on the software library VNODE developed by Nedialkov *et al.* [15]. Roughly speaking, this means that all computations, from adding numbers up to integration, amount to working with two-sided estimates; e.g. the sum of two real numbers $a \in [a_1, a_2]$ and $b \in [b_1, b_2]$ is the interval $[a_1 + b_1, a_2 + b_2]$ which is guaranteed to contain $a + b$ (see [16, §2] for a more detailed description). If we obtain that

$$\frac{1}{2\pi i} \int_{\mathcal{R}_0} \frac{\Delta'(z)}{\Delta(z)} dz \in [c_1, c_2] \quad \text{and} \quad [c_1, c_2] \cap \mathbb{N}_0 = \{n_0\}, \quad (1.2)$$

then there are precisely n_0 eigenvalues of the complex-scaled Hamiltonian in the rectangle \mathcal{R}_0 and hence precisely n_0 resonances in the rotated rectangle $e^{-2i\theta}\mathcal{R}_0$.

Our method is the first, in both physical and mathematical literature, that accomplishes the following three different tasks:

1. Enclose resonances with prescribed accuracy, by choosing the size of the rectangle accordingly small and achieving $n_0 = 1$.
2. Exclude resonances in certain rectangles by achieving $n_0 = 0$.
3. Check if the number of resonances in a rectangle of arbitrary size computed with non-reliable methods is correct by checking if it coincides with n_0 .

2. Complex scaling and lack of analytic information

There are various mathematical definitions of resonances and different methods to study them; for details, we refer to the comprehensive review articles by Simon [17], Siedentop [18] and Harrell [19]. Here, we use the method of complex scaling where resonances are characterized as eigenvalues of certain non-self-adjoint Schrödinger operators.

As an example, we consider the spectral problem (1.1), with $\mu/\hbar^2 = 1$ for the sake of simplicity. If we set $\lambda := 2\epsilon - 1.6$, it is easy to see that (1.1) is equivalent to the spectral problem

$$-y''(x) + (x^2 - 1.6)e^{-x^2/10}y(x) - \lambda y(x) = 0, \quad x \in \mathbb{R}, \quad (2.1)$$

for the linear operator L in the Hilbert space $L_2(\mathbb{R})$ given by

$$\begin{aligned} D(L) &:= W_2^2(\mathbb{R}) := \{y \in L_2(\mathbb{R}) : y', y'' \in L_2(\mathbb{R})\}, \\ (Ly)(x) &:= -y''(x) + (x^2 - 1.6)e^{-x^2/10}y(x); \end{aligned}$$

note that $W_2^2(\mathbb{R})$ is the closure of $C_0^\infty(\mathbb{R})$ with respect to the norm of $W_2^2(\mathbb{R})$ given by $\|y\|_{2,2} := (\|y\|_2^2 + \|y'\|_2^2 + \|y''\|_2^2)^{1/2}$, where y', y'' denote the weak derivatives and $\|\cdot\|_2$ denotes the norm of $L_2(\mathbb{R})$ ([20, ch. V]).

According to the method of complex scaling ([21,22], [23, §5] and also [24]), for every $\theta \in [0, \pi/4)$, the spectral problem (2.1) is equivalent to the spectral problem for the operator H_θ in $L_2(\mathbb{R})$ given by $D(H_\theta) = W_2^2(\mathbb{R})$ and

$$(H_\theta y)(x) := -y''(x) + q_\theta(x)y(x) = zy(x), \quad x \in \mathbb{R}, \quad z := e^{2i\theta}\lambda, \quad (2.2)$$

with complex-valued potential

$$q_\theta(x) := e^{2i\theta}(e^{2i\theta}x^2 - 1.6)e^{-e^{2i\theta}x^2/10}, \quad x \in \mathbb{R}. \quad (2.3)$$

Hence, z is an eigenvalue of (2.2) if and only if $\lambda = e^{-2i\theta}z$ is a resonance of (2.1) or, equivalently, if $\epsilon = (\lambda + 1.6)/2 = (e^{-2i\theta}z + 1.6)/2$ is a resonance of (1.1).

Because q_θ is even, the spectral problem (2.2) for the operator H_θ is equivalent to the two spectral problems

$$-y''(x) + q_\theta(x)y(x) = zy(x), \quad x \in [0, \infty), \quad y(0) = 0 \quad (2.4)$$

and

$$-y''(x) + q_\theta(x)y(x) = zy(x), \quad x \in [0, \infty), \quad y'(0) = 0, \quad (2.5)$$

for the operators H_θ^D and H_θ^N induced by the differential expression $\tau_\theta y := -y'' + q_\theta y$ in $L_2([0, \infty))$ with Dirichlet and with Neumann boundary condition, respectively. This was proved in [12, §5] using the Weyl–Titchmarsh function. For eigenvalues, this follows from the following elementary argument.

If $z_0 \in \mathbb{C}$ is an eigenvalue of (2.2) with eigenfunction $y_0 \in D(H_\theta) \subset L_2(\mathbb{R})$, then, by (2.3), the function \tilde{y}_0 given by $\tilde{y}_0(x) := y_0(-x)$ is an eigenfunction as well. Because $y_0(0) = \tilde{y}_0(0)$, the functions y_0, \tilde{y}_0 must be linearly dependent. The particular form of \tilde{y}_0 implies that $\tilde{y}_0 = \gamma y_0$ with $\gamma = \pm 1$. Because $y_0 \in W_2^2(\mathbb{R}) \subset C^1(\mathbb{R})$, the continuity of y_0 and y_0' in 0 yields that $y_0'(0) = 0$

if $\gamma = 1$ and $y_0(0) = 0$ if $\gamma = -1$. Hence, $y_0|_{[0,\infty)}$ is either an eigenfunction of (2.4) or of (2.5). Vice versa, if $z_0 \in \mathbb{C}$ is an eigenvalue of (2.4) with eigenfunction $y_0 \in D(H_\theta^D) \subset L_2([0, \infty))$, we obtain an eigenfunction $y_0 \in D(H_\theta) \subset L_2(\mathbb{R})$ of (2.2) at λ_0 by setting $y_0(x) := -y_0(-x)$, $x \in (-\infty, 0)$; if $z_0 \in \mathbb{C}$ is an eigenvalue of (2.5), we set $y_0(x) := y_0(-x)$, $x \in (-\infty, 0)$.

Because the potential q_θ is complex-valued and hence all the above operators H_θ along with H_θ^D , H_θ^N are no longer self-adjoint, numerical approximations of eigenvalues—and hence of resonances—are prone to be unstable. Examples for such instabilities may be found in [23] for resonances, but they occur already for eigenvalues of matrices (see e.g. [16] for the famous Godunov matrix).

Analytic bounds for resonances are commonly based on numerical range estimates for each complex-scaled problem (2.2) with $\theta \in [0, \pi/4)$ (comp. [14]). For the set of resonances of (1.1), we obtain the following result.

Theorem 2.1. *The resonances of (1.1) in the sector $-\pi/2 < \arg \epsilon \leq 0$ are contained in the closed convex set*

$$\mathcal{C} := \bigcap_{\theta \in [0, \pi/4)} \{ \epsilon \in \mathbb{C} : \operatorname{Re}(\epsilon) \sin(2\theta) + \operatorname{Im}(\epsilon) \cos(2\theta) \leq 0.5a_+(\theta) + 0.8 \sin(2\theta) \}$$

where $a_+(\theta) := \sup_{x \in [0, \infty)} \operatorname{Im}(q_\theta(x))$ for $\theta \in [0, \pi/4)$ with

$$\operatorname{Im}(q_\theta(x)) = e^{-\cos(2\theta)x^2/10} \left(x^2 \sin \left(4\theta - \sin(2\theta) \frac{x^2}{10} \right) - 1.6 \sin \left(2\theta - \sin(2\theta) \frac{x^2}{10} \right) \right).$$

Proof. The set \mathcal{C} is closed and convex being the intersection of closed half-planes. Because $a_+(\theta) \geq 0$ and hence $a_+(\theta)/(2 \sin(2\theta)) + 0.8 \geq 0.8$, it follows that \mathcal{C} contains all $\epsilon \in \mathbb{C}$ with $0 < \operatorname{Re}(\epsilon) \leq 0.8$ and $\operatorname{Im}(\epsilon) \leq 0$.

Thus, it is sufficient to show that every resonance $\epsilon_0 \in \mathbb{C}$ with $\operatorname{Re}(\epsilon_0) > 0.8$, $\operatorname{Im}(\epsilon_0) \leq 0$ belongs to \mathcal{C} . For every $\theta \in [0, \pi/4]$, the point $\lambda_0 := 2\epsilon_0 - 1.6$ lies in the sector $-\pi/2 < \arg \lambda \leq 0$ and is an eigenvalue of the operator $\tilde{H}_\theta := e^{-2i\theta} H_\theta$ given by

$$D(\tilde{H}_\theta) = W_2^2(\mathbb{R}), \quad \tilde{H}_\theta y = e^{-2i\theta} (-y'' + q_\theta y).$$

Because the numerical range of a linear operator contains all eigenvalues, we obtain

$$\lambda_0 \in W(\tilde{H}_\theta) := \{ (\tilde{H}_\theta y, y) : y \in D(\tilde{H}_\theta), \|y\| = 1 \}, \quad \theta \in \left[0, \frac{\pi}{4} \right).$$

If we note that $q_\theta(\mathbb{R}) = q_\theta([0, \infty))$ and, in addition to $a_+(\theta)$, we define

$$a_-(\theta) := \inf_{x \in [0, \infty)} \operatorname{Im}(q_\theta(x)), \quad b_-(\theta) := \inf_{x \in [0, \infty)} \operatorname{Re}(q_\theta(x)),$$

then it is easy to see that, for $\theta \in [0, \pi/4)$,

$$W(\tilde{H}_\theta) \subset e^{-2i\theta} \{ z \in \mathbb{C} : a_-(\theta) \leq \operatorname{Im}(z) \leq a_+(\theta), b_-(\theta) \leq \operatorname{Re}(z) \}.$$

In particular, every resonance λ_0 of L with $-\pi/2 < \arg \lambda \leq 0$ satisfies

$$\lambda_0 \in \bigcap_{\theta \in [0, \pi/4)} \{ \lambda \in \mathbb{C} : \operatorname{Re}(\lambda) \sin(2\theta) + \operatorname{Im}(\lambda) \cos(2\theta) \leq a_+(\theta) \}. \quad \blacksquare$$

Figure 1 shows that the only available analytic information is much too coarse to judge the validity or non-validity of resonance approximations. Therefore, it is necessary to employ a method yielding both guaranteed and much more accurate enclosures and exclosures for eigenvalues of non-self-adjoint eigenvalue problems.




Figure 1. Resonance-free region in theorem 2.1 (grey-shaded), approximate resonance strings of Rittby *et al.* (circles) and Abramov *et al.* (squares and star). (Online version in colour.)

3. Eigenvalue enclosures for complex-valued potentials

The algorithm we use to establish guaranteed eigenvalue enclosures was developed and described in detail in [16,25]. Briefly, it consists of the following two steps. For the sake of simplicity, we consider the Dirichlet problem (2.4); the approach to the Neumann problem (2.5) is completely analogous.

Step A. Solving a truncated problem with guaranteed error bounds. In order to truncate problem (2.4), we restrict the potential q_θ to an interval $[0, X]$ and set it equal to 0 on (X, ∞) . The unique (up to scalar multiplication) solution of $-y'' = zy$ in $L_2([X, \infty))$ is $\exp(-\sqrt{-z}x)$ for $\text{Re}\sqrt{-z} > 0$. Hence, the problem on $[0, X]$, we have to solve is

$$\text{and } \left. \begin{aligned} -y''(x) + q_\theta(x)y(x) &= zy(x), & x \in [0, X] \\ y(0) = 0, \quad y'(X) &= -\sqrt{-z}y(X). \end{aligned} \right\} \quad (3.1)$$

The eigenvalues of this regular boundary value problem can be characterized as the zeros of an analytic function Δ and may thus be counted and found by means of the argument principle.

The algorithms for the calculation of the analytic function Δ and for the contour integral over a chosen starting box $\mathcal{R}_0 \subset \mathbb{C}$ are performed in interval arithmetic, i.e. with guaranteed error bounds. Having achieved (1.2), we obtain a box that contains a certain number n_0 of eigenvalues of the truncated problem (3.1). Repeating this procedure by suitably subdividing the box \mathcal{R}_0 , we may finally arrive at a box \mathcal{R}_Z of desired precision ε_Z that contains exactly one eigenvalue z_{trunc} .

Step B. Use Levinson asymptotics to enclose the eigenvalues of problem (2.4). If $y_2(\cdot, z)$ is the unique (suitably normalized) solution of the differential equation in (2.4) belonging to $L_2([0, \infty))$ for $z \in \mathbb{C} \setminus [0, \infty)$, then z_{true} is an eigenvalue of (2.4) if and only if $y_2(0, z_{\text{true}}) = 0$. Levinson's theorem (see e.g. [25, theorem 3.3]) shows that

$$y_2(x, z) = \exp(-\sqrt{-z}x)(1 + \eta(x)), \quad |\eta(x)| \leq \frac{\alpha_{X,\theta}}{1 - \alpha_{X,\theta}}, \quad \alpha_{X,\theta} := \int_X^\infty |q_\theta(x)| dx, \quad (3.2)$$

for all $X \geq 0$ such that $\alpha_{X,\theta} < 1$. Hence, if $[E] \subset \mathbb{R}$ is an interval with

$$\left[1 - \frac{\alpha_{X,\theta}}{1 - \alpha_{X,\theta}}, 1 + \frac{\alpha_{X,\theta}}{1 - \alpha_{X,\theta}} \right] \subset [E],$$

and $[y_2(\cdot, z)]$ is an interval-valued solution of the truncated problem on $[0, X]$ satisfying the interval-valued initial conditions

$$y(X, z) \in [E] \exp(-\sqrt{-z}X), \quad y'(X, z) \in -[E]\sqrt{-z} \exp(-\sqrt{-z}X),$$

then $y_2(0, z) \in [y_2(0, z)]$. By means of the interval arithmetic argument principle already used in step A, we now obtain enclosures for the zeros of $y_2(0, z)$, and hence for the eigenvalues z_{true} of (2.4) of desired precision.

For the above-described method, several parameters have to be provided; in particular, the length X of the truncated interval has to be determined such that $\alpha_{X,\theta} < 1$. To this end, we note that

$$|q_\theta(x)| = |e^{2i\theta}x^2 - 1.6| e^{-\cos(2\theta)x^2/10} \leq x^2 e^{-\cos(2\theta)x^2/10} \quad \text{if } \cos(2\theta) \geq \frac{0.8}{x^2}$$

and that [26, 7.1.13]

$$\int_{x_0}^{\infty} e^{-x^2} dx \leq e^{-x_0^2} \frac{1}{x_0 + \sqrt{x_0^2 + 4/\pi}}, \quad x_0 \geq 0.$$

Integrating by parts and substituting $t = \sqrt{ax}$, we obtain, for $a \geq 0$,

$$\begin{aligned} \int_X^{\infty} x^2 e^{-ax^2} dx &= \frac{1}{a} X e^{-aX^2} + \frac{1}{a\sqrt{a}} \int_{\sqrt{a}X}^{\infty} e^{-t^2} dt \\ &\leq \frac{1}{a} e^{-aX^2} \left(X + \frac{1}{aX + \sqrt{a^2X^2 + 4a/\pi}} \right) \\ &\leq \frac{1}{a} e^{-aX^2} \left(X + \frac{1}{2aX} \right). \end{aligned}$$

Hence, for all $X \in (0, \infty)$, $\theta \in [0, \pi/4)$ with $\cos(2\theta) \geq 0.8/x^2$, we can estimate

$$\alpha_{X,\theta} \leq \int_X^{\infty} x^2 e^{-\cos(2\theta)x^2/10} dx \leq \frac{10}{\cos(2\theta)} e^{-\cos(2\theta)X^2/10} \left(X + \frac{5}{\cos(2\theta)X} \right) =: A_{X,\theta} \quad (3.3)$$

and we use the analytic expression $A_{X,\theta}$ to obtain a rigorous computable upper bound $A_{X,\theta}^0$ for $A_{X,\theta}$ and hence for $\alpha_{X,\theta}$,

$$\alpha_{X,\theta} \leq A_{X,\theta} \leq A_{X,\theta}^0.$$

To this end, we first expand $\cos(2\theta)$ and use Taylor's theorem with remainder in Lagrange form to see that, for every $m \in \mathbb{N}$,

$$\cos(2\theta) \geq \sum_{j=0}^{4m} \frac{(-1)^j}{(2j)!} (2\theta)^{2j} =: T_{X,\theta}(m); \quad (3.4)$$

note that $\cos^{(4m+1)}(x) = -\sin(x) \leq 0$ for every $x \in [0, 2\theta] \subset [0, \pi/2]$. If θ is a decimal fraction whose fractional part has three digits, the sum $T_{X,\theta}(m)$ is rational and can be evaluated exactly. We choose a rigorous computable lower bound $T_{X,\theta}^0(m)$ of $T_{X,\theta}(m)$ as the unique decimal number whose fractional part has six digits and $T_{X,\theta}^0(m) + 10^{-6} > T_{X,\theta}(m) \geq T_{X,\theta}^0(m)$ (table 1). The function $f(t) := (10/t)e^{-tX^2/10}(X + 5/tX)$, $t \in (0, 1)$, is decreasing, hence, again by Taylor's theorem with remainder in Lagrange form, we obtain that, for all $m, n \in \mathbb{N}$,

$$\begin{aligned} A_{X,\theta} &= f(\cos(2\theta)) \leq f(T_{X,\theta}^0(m)) \\ &\leq \frac{10}{T_{X,\theta}^0(m)} \left(\sum_{k=0}^{2n+1} \frac{(-1)^k X^{2k}}{10^k k!} (T_{X,\theta}^0(m))^k \right) \left(X + \frac{5}{T_{X,\theta}^0(m)X} \right) =: A_{X,\theta}(m, n). \end{aligned}$$

Now, we fix $m, n \in \mathbb{N}$ and proceed in the same way as for $T_{X,\theta}(m)$ to obtain a rigorous computable upper bound $A_{X,\theta}^0(m, n)$ for $A_{X,\theta}(m, n)$ with $A_{X,\theta}^0(m, n) - 10^{-6} < A_{X,\theta}(m, n) \leq A_{X,\theta}^0(m, n)$ (table 1). Because $\theta \mapsto A_{X,\theta}(m, n)$, $\theta \in [0, \pi/4)$, is increasing, the rigorous computable upper bound

Table 1. Rigorous computable bounds for $X = 50$ and various $\theta \in [0, \pi/4]$.

for $\theta \in (0, 0.75]$:	$T_{X,\theta}^0 = 0.070737$	$A_{X,\theta}^0 = 0.000152$
for $\theta = 0.755$:	$T_{X,\theta}^0 = 0.060758$	$A_{X,\theta}^0 = 0.00216$

$A_{X,\theta_0}^0(m, n)$ for $\theta_0 := 0.75 < \pi/4$ is also an upper bound of $A_{X,\theta}(m, n)$ for $\theta \in (0, \theta_0)$. Only in two of our computations (for the resonances numbered 37^- and 44^+), we needed parameter values θ that are larger than $\theta_0 = 0.75$; their upper bound $A_{X,\theta}^0(m, n)$ is computed separately. We use $X = 50$, $m = 2$, $n = 32$ and obtain the rigorous computable lower bounds $T_{X,\theta}^0(m) =: T_{X,\theta}^0$ and upper bounds $A_{X,\theta}^0(m, n) =: A_{X,\theta}^0$ displayed in table 1; note that for $X = 50$ the condition $\cos(2\theta) \geq 0.8/x^2$ allows for $\theta \leq 0.5 \arccos(\frac{8}{25}10^{-5})$, e.g. $\theta \leq 0.785238$ very close to $\pi/4 \sim 0.7853981635$.

4. Guaranteed resonance enclosures and exclosures

In [10, reply to comment], Rittby *et al.* listed a set of 44 approximate resonances ϵ_k^\pm of (1.1) that they computed numerically, along with a set of 40 approximate resonances claimed to be found numerically by Korsch *et al.* in [5, comment]; here, the superscript $+$ occurs for even k , whereas $-$ occurs for odd k . The differences in modulus between these two approximate resonance strings are smaller than $2 \cdot 10^{-3}$ up to ϵ_{15}^- and start to be larger than 10^{-2} from ϵ_{16}^+ on, getting as huge as 56.19 for ϵ_{40}^+ (figure 2).

We computed guaranteed enclosures for all 44 approximate resonances by Rittby *et al.* from [4] as well as exclosures for the approximate resonances ϵ_{16}^+ up to ϵ_{28}^+ by Korsch *et al.* from [5, comment]. In addition, we enclosed the two pairs of resonances discovered numerically in [14] that are visible by the complex scaling method.

All computed enclosures for resonances, except for one of these pairs, were performed with interval length $X = 50$, varying scaling angle θ as displayed in the tables, and corresponding guaranteed upper bound $A_{X,\theta}^0$ for $\alpha_{X,\theta}$ as in table 1 at the end of §3. The enclosure for one of the additional resonance pairs in [14] turned out to be by far more challenging than all other computations.

We employ the interval arithmetic-based software library VNODE developed by Nedialkov *et al.* (see [15]) where all operations are performed with complex ‘intervals’, i.e. rectangles $[z] = [x] + [y]i$, where $[x], [y] \subset \mathbb{R}$ are closed intervals or singletons (see [16, §2] for a more detailed description). In the following, we use notation of the form

$$7.439759^{70244416010}_{16958921987} := [7.43975916958921987, 7.43975970244416010]$$

for intervals containing the real and imaginary part of resonances. Further, we use the enumeration ϵ_k^\pm to indicate the resonance number k and parity \pm in the list of approximate resonances of Rittby *et al.* in [10, reply to comment, table I].

Note that the resonances coming from the boundary condition $y(0) = 0$ have parity ‘ $-$ ’, because the eigenfunctions of the corresponding eigenvalues of (2.4) are odd, whereas those coming from the boundary condition $y'(0) = 0$ have parity ‘ $+$ ’, because the eigenfunctions of the corresponding eigenvalues of (2.5) are even (see §2).

(a) Guaranteed enclosures for resonance approximations by Rittby *et al.*

First, we present the computed enclosures for the 44 resonances λ_k^\pm of problem (2.1) corresponding to the resonances ϵ_k^\pm listed in [10, reply to comment, table I].

Table 2 contains the enclosures for resonances $\lambda = e^{-2i\theta}z$ via enclosures for eigenvalues z of (2.2) restricted to $[0, \infty)$ with Dirichlet boundary condition $y(0) = 0$, i.e. eigenvalues of problem (2.4); table 3 contains the corresponding enclosures using eigenvalues z of (2.2) restricted to $[0, \infty)$ with Neumann boundary condition $y'(0) = 0$, i.e. for eigenvalues of problem (2.5). Table 4




Figure 2. Resonance approximations computed by Rittby *et al.* (circles), Korsch *et al.* (asterisks), Abramov *et al.* (squares and star) and analytic bound from theorem 2.1. (Online version in colour.)

contains the enclosures for the 44 resonances $\epsilon_k^\pm = (\lambda_k^\pm + 1.6)/2$ of the original problem (1.1) arising from the two sets of resonances λ_k^\pm of (2.2) displayed in tables 2 and 3.

The enclosing boxes for the resonances ϵ of (1.1) are obtained from the enclosing boxes for the eigenvalues z of (2.2) as follows. If $[u_1, u_2] + [v_1, v_2]i$ is an enclosing box in the z -plane, then the enclosing box $[x_1, x_2] + [y_1, y_2]i$ for a resonance $\lambda = e^{-2i\theta} z$ of (2.1) is the smallest axis-parallel box that contains the rotated box $e^{-2i\theta}([u_1, u_2] + [v_1, v_2]i)$. The corresponding enclosing box for a resonance $\epsilon = (\lambda + 1.6)/2$ of (1.1) is obtained from

$$\lambda \in [x_1, x_2] + [y_1, y_2]i \iff \epsilon \in \left[\frac{x_1 + 1.6}{2}, \frac{x_2 + 1.6}{2} \right] + \left[\frac{y_1}{2}, \frac{y_2}{2} \right]i.$$

The values of the 44 approximate resonances of (1.1) listed in [10, reply to comment, table I], which were computed by Rittby *et al.* in floating point arithmetic without error bounds, are displayed in the right column in table 4; they agree with our enclosures at least up to order 10^{-4} . Thus, our guaranteed enclosures prove that all values computed by Rittby *et al.* do indeed lie near true resonances.

(b) Guaranteed enclosures for resonance approximations by Korsch *et al.*

On the other hand, we applied our method to the numerical values of the resonance approximations of Korsch *et al.* numbered $16^+, 17^-, \dots, 27^-, 28^+$ in [10, reply to comment, table II]; note that the resonance approximations $29^-, \dots, 40^+$ therein can not be seen by the complex scaling method.

Using larger boxes around these numerical values, we found that in each case the interval-valued argument principle yields an interval $[c_1, c_2]$ with $[c_1, c_2] \cap \mathbb{N}_0 = \{0\}$, which proves that there are no eigenvalues in the considered box (see (1.2)). The box side lengths $l_k \in [0.1, 2]$ are listed in table 5. For every resonance approximation ϵ_k^\pm , the corresponding approximate value in the z -plane is denoted by z_k^\pm . The midpoint $M_k \in \mathbb{C}$ of the box with side length l_k in the z -plane is chosen such that

$$|\operatorname{Re}(M_k) - \operatorname{Re}(z_k^\pm)| < 0.05 \leq \frac{l_k}{2}, \quad |\operatorname{Im}(M_k) - \operatorname{Im}(z_k^\pm)| < 0.05 \leq \frac{l_k}{2}.$$

Table 2. Resonances for (2.2) from (2.4) on $[0, \infty)$ with $y(0) = 0$.

$[\lambda_1^-]$	1.241941 ⁹³⁹³²⁷⁹⁵⁷ ₈₆₀₁₈₈₂₉₇	-0.0001165 ¹⁴⁸⁸⁸³³³⁰⁸⁵⁸ ₉₂₂₃₂₈₈₆₄₀₃₃ i	$\theta = 0.5$
$[\lambda_3^-]$	3.569165 ⁸⁰⁶⁹⁹⁷³¹¹ ₆₇₃₀₁₂₀₇	-0.347501 ³²¹⁵⁵²⁵⁰² ₄₅₅₅₃₇₇₄₂ i	$\theta = 0.45$
$[\lambda_5^-]$	4.910972 ⁵⁶⁹⁰⁶⁸⁰² ₄₃₄₂₁₂₄₆₂	-2.223063 ¹⁰⁶⁰²⁹⁵²⁵ ₂₄₀₈₈₅₀₉₆ i	$\theta = 0.4$
$[\lambda_7^-]$	6.04865 ⁹⁰⁷³⁷⁵¹⁴⁰⁷ ₈₉₉₃₇₄₂₄₅₂	-4.974892 ³²⁰²⁹¹⁵⁴⁸ ₄₁₀₄₂₃₆₁₁ i	$\theta = 0.4$
$[\lambda_9^-]$	6.899910 ¹⁶¹⁹⁰⁰⁷⁶⁹ ₀₇₉₅₄₀₆₃₇	-8.366339 ²⁵⁸⁵¹⁶⁸⁹³ ₃₄₀₈₇₇₀₅₈ i	$\theta = 0.5$
$[\lambda_{11}^-]$	7.45761 ⁹⁰²⁰²⁸³⁴¹⁸ ₈₉₄₀₁₂₆₈₅₉	-12.309461 ²²¹⁶³³⁵³ ₃₀₁₇₉₀₁₄ i	$\theta = 0.55$
$[\lambda_{13}^-]$	7.723776 ⁸³⁵³²⁵⁴¹⁶ ₇₆₁₉₄₈₆₇₀	-16.751102 ²⁸¹¹³⁸⁴⁹ ₃₅₄₅₁₅₂₈ i	$\theta = 0.65$
$[\lambda_{15}^-]$	7.688281 ¹⁶⁷⁵⁶⁴⁸³⁴ ₀₉₈₆₉₆₅₉₃	-21.652520 ⁸⁴¹³⁵⁸⁷⁹ ₉₁₀₂₂₇₀₇ i	$\theta = 0.7$
$[\lambda_{17}^-]$	7.439759 ⁹⁹⁹³¹⁵⁰⁰ ₂₇₅₁₁₃₅₃	-26.931640 ⁴³⁴¹⁸⁰⁶ ₅₅₉₂₄₉₅ i	$\theta = 0.73$
$[\lambda_{19}^-]$	5.34207 ⁶³⁷⁷⁸⁶⁴⁴¹⁴ ₅₈₄₇₀₄₁₇₄₈	-30.84463 ⁰⁹³⁷⁵⁶⁷¹⁷ ₁₄₆₈₃₈₉₉₀ i	$\theta = 0.725$
$[\lambda_{21}^-]$	6.96683 ²²⁶¹⁷²²⁶⁵⁰ ₁₂₁₆₉₂₄₆₄₂	-33.21458 ⁸⁵⁵⁴⁶⁸⁹⁷⁸ ₉₅₉₉₄₈₇₈₆ i	$\theta = 0.735$
$[\lambda_{23}^-]$	6.915830 ⁶¹⁴⁸⁸³⁹⁶⁸ ₅₄₈₀₁₄₅₅₂	-37.987895 ⁶⁹⁵¹¹⁸⁶⁴ ₇₆₁₉₈₈₁₄ i	$\theta = 0.72$
$[\lambda_{25}^-]$	6.91656 ⁹⁴³⁴⁴⁸⁷⁵⁹² ₀₉₄₁₃₂₅₃₂₃	-41.75293 ⁰⁹⁶⁰⁶⁸⁵⁹⁴ ₉₄₅₃₈₄₈₂₉ i	$\theta = 0.725$
$[\lambda_{27}^-]$	7.263855 ⁶⁴²⁰¹¹⁰⁹⁴ ₅₀₉₃₀₅₃₉₇	-46.005638 ⁵⁰⁹⁹²¹⁷⁶ ₆₄₂₆₂₇₅₅ i	$\theta = 0.725$
$[\lambda_{29}^-]$	7.1713970 ⁹²⁶⁹⁸⁵⁷² ₂₀₂₇₈₄₁₈	-50.640608 ³⁰²⁵⁷¹¹³ ₄₂₇₆₄₀₀₉ i	$\theta = 0.73$
$[\lambda_{31}^-]$	7.41606 ⁸⁹⁵⁶¹¹⁹⁷⁶³ ₆₃₇₅₆₅₇₇₇₃	-55.41738 ⁷²⁹¹⁷²²⁷⁸ ₉₄₅₀₉₉₅₂₉ i	$\theta = 0.73$
$[\lambda_{33}^-]$	7.595038 ⁸⁷¹⁹¹⁶³⁰⁶ ₇₇₄₅₀₅₁₇₀	-60.2052079 ²³⁷³⁹⁷⁶ ₉₁₄₇₅₉₂ i	$\theta = 0.74$
$[\lambda_{35}^-]$	7.354328 ⁶³⁴⁹³⁴¹⁹⁸ ₅₄₉₉₁₉₇₈₅	-65.0217411 ²⁸⁵⁹¹⁹¹ ₉₄₄₉₆₆₁ i	$\theta = 0.745$
$[\lambda_{37}^-]$	5.26985 ⁵⁹⁸⁵⁵⁶⁵¹²⁵ ₁₉₄₆₁₃₈₈₃₈	-68.0334 ³⁸⁹⁰⁰⁶⁸³²¹ ₄₂₉₄₀₁₀₉₆₂ i	$\theta = 0.755$
$[\lambda_{39}^-]$	7.31756 ²⁰⁴⁰⁴¹³⁶⁰⁰ ₉₇₂₅₂₅₆₈₆	-71.126990 ⁰⁴⁴⁸⁸¹²⁴ ₁₆₈₀₀₈₃₆ i	$\theta = 0.75$
$[\lambda_{41}^-]$	7.6551306 ⁷⁸⁵⁷³⁴¹¹ ₁₄₉₀₁₇₆₀	-76.322137 ¹⁸⁸³⁰²⁰² ₂₅₁₉₇₃₈₁ i	$\theta = 0.75$
$[\lambda_{43}^-]$	7.54176 ²⁵⁰⁵⁰⁰⁴⁷¹³ ₁₄₁₈₇₉₈₈₀₆	-81.61068 ⁰⁴⁶⁵³¹⁸³² ₂₄₃₅₃₄₉₄₉ i	$\theta = 0.75$

The corresponding box in the ϵ -plane is scaled and rotated owing to the relation $\epsilon_k^\pm = (e^{-2i\theta} z_k^\pm + 1.6)/2$. The box has side length $l_k/2$ and is rotated clockwise by the angle 2θ around the midpoint $m_k := (e^{-2i\theta} M_k + 1.6)/2$. The minimal distance d_k of ϵ_k^\pm to the boundary of the rotated box satisfies $d_k > (l_k/4) - 0.025 \geq 0$ (figure 3).

Hence, our guaranteed enclosures prove that none of the numerical values of Korsch *et al.* numbered 16^+ , 17^- , \dots , 27^- , 28^+ lies near a true resonance of (1.1).

(c) Enclosures of resonance approximations by Abramov *et al.*

Finally, we considered the three pairs of additional resonances found in [14, p. 72], one pair near each of the points

$$\hat{\epsilon}_1 = 0.69 - 7.91i, \quad \hat{\epsilon}_2 = 1.26 - 8.51i, \quad \hat{\epsilon}_3 = 2.08 - 11.61i;$$

Table 3. Resonances for (2.2) from (2.5) on $[0, \infty)$ with $y'(0) = 0$.

$[\lambda_2^+]$	$2.654394_{092835094}^{176821622}$	$-0.030894_{64975797340}^{56577144434} i$	$\theta = 0.35$
$[\lambda_4^+]$	$4.248843_{771878984}^{813059968}$	$-1.1295890_{76541343}^{25361269} i$	$\theta = 0.5$
$[\lambda_6^+]$	$5.514431_{624202591}^{752453071}$	$-3.511012_{170826048}^{042575560} i$	$\theta = 0.55$
$[\lambda_8^+]$	$6.510869_{364918580}^{433786815}$	$-6.597282_{116776595}^{047908355} i$	$\theta = 0.7$
$[\lambda_{10}^+]$	$7.2155450_{18333105}^{98489665}$	$-10.272906_{38654053}^{30638393} i$	$\theta = 0.55$
$[\lambda_{12}^+]$	$7.6261961_{08709429}^{185861521}$	$-14.470753_{13697016}^{05981802} i$	$\theta = 0.6$
$[\lambda_{14}^+]$	$7.74690_{5966973172}^{6068847760}$	$-19.141473_{98214909}^{388027448} i$	$\theta = 0.75$
$[\lambda_{16}^+]$	$7.59265_{6102498221}^{7204389647}$	$-24.28598_{528682110}^{418492963} i$	$\theta = 0.7$
$[\lambda_{18}^+]$	$6.93961_{3775608598}^{4828882919}$	$-29.61377_{609379499}^{504052061} i$	$\theta = 0.73$
$[\lambda_{20}^+]$	$5.463690_{436278557}^{503147965}$	$-31.299089_{45643974}^{38957026} i$	$\theta = 0.72$
$[\lambda_{22}^+]$	$7.09171_{3513412909}^{5717195734}$	$-35.73909_{940568757}^{720190466} i$	$\theta = 0.7$
$[\lambda_{24}^+]$	$6.7766621_{07796643}^{74666058}$	$-39.913828_{10780846}^{04093896} i$	$\theta = 0.72$
$[\lambda_{26}^+]$	$7.1641875_{27769781}^{94122646}$	$-43.794599_{31759526}^{25124230} i$	$\theta = 0.725$
$[\lambda_{28}^+]$	$7.217834_{296771199}^{827593876}$	$-48.285288_{93378272}^{40295994} i$	$\theta = 0.725$
$[\lambda_{30}^+]$	$7.26422_{0853959247}^{117277859}$	$-53.03794_{509870568}^{483538696} i$	$\theta = 0.73$
$[\lambda_{32}^+]$	$7.532859_{772590297}^{837890213}$	$-57.809513_{70761888}^{64231886} i$	$\theta = 0.735$
$[\lambda_{34}^+]$	$7.563081_{137412970}^{906616847}$	$-62.59350_{947366904}^{883239546} i$	$\theta = 0.735$
$[\lambda_{36}^+]$	$6.577678_{639847483}^{703519130}$	$-67.9668690_{7952254}^{1585076} i$	$\theta = 0.75$
$[\lambda_{38}^+]$	$5.86009_{0527100523}^{5868261508}$	$-68.1042_{1052927093}^{0518810982} i$	$\theta = 0.75$
$[\lambda_{40}^+]$	$7.566457_{115996163}^{416558969}$	$-73.72887_{409019465}^{387699096} i$	$\theta = 0.745$
$[\lambda_{42}^+]$	$7.629471_{689847954}^{817191209}$	$-78.94320_{909751929}^{897017587} i$	$\theta = 0.75$
$[\lambda_{44}^+]$	$7.424570_{055864947}^{340663545}$	$-84.175934_{92887123}^{43734233} i$	$\theta = 0.755$

the corresponding values $\lambda = 2\epsilon - 1.6$ are

$$\hat{\lambda}_1 = -0.22 - 15.82i, \quad \hat{\lambda}_2 = 0.92 - 17.02i, \quad \hat{\lambda}_3 = 2.46 - 23.22i.$$

These new resonances were conjectured to exist not by means of complex scaling, but by exploiting the asymptotic properties of the solution of a differential equation with a rapidly decaying potential; numerical methods were developed to locate these resonances (see [14, §6]). In fact, the method of complex scaling does not allow one to see the first pair of resonances near $\hat{\lambda}_1$ because it has negative real part, but it does allow one to see the second and third pair near $\hat{\lambda}_2, \hat{\lambda}_3$.

We computed guaranteed enclosures for the two pairs of resonances near $\hat{\lambda}_2, \hat{\lambda}_3$; each of these pairs originates in one eigenvalue z of (2.4) with boundary condition $y(0) = 0$ with odd eigenfunction (denoted by superscript ‘-’) and one eigenvalue z of (2.5) with boundary condition

Table 4. Resonances for (1.1).

	guaranteed enclosures		numerical values by Rittby <i>et al.</i>
ϵ_1^-	1.4209709 ⁶⁹⁶⁶³⁹⁷⁸⁵ ₃₀₀₉₄₁₄₈₅	—0.0000582 ⁵⁷⁴⁴⁴¹⁶⁶⁵⁴²⁹⁰ ₉₆₁₆₄₄₃₂₀₁₆₅ i	1.420971 — 0.00005826663i
ϵ_2^+	2.1271970 ⁸⁸⁴¹⁰⁸¹¹⁰ ₄₆₄₁₇₅₄₇₀	—0.015447 ²⁸²⁸⁸⁵⁷²²¹⁷⁰ ₃₂₄₈₇₈₉₈₆₇₀₀ i	2.127197 — 0.01544732i
ϵ_3^-	2.584582 ⁹⁰³⁴⁹⁸⁶⁵⁵ ₈₃₆₅₀₆₀₃₅	—0.173750 ⁶⁶⁰⁷⁷⁶²⁵¹⁰ ₇₂₇₇₆₈₈₇₁₀ i	2.584583 — 0.1737507i
ϵ_4^+	2.92442 ⁹⁰⁶⁵²⁹⁹⁸⁴⁰ ₈₈₅₉₃₉₉₄₇₀	—0.5647949 ⁶⁷⁶⁸⁰⁶³⁴⁵ ₈₈₂₇₀₆₇₁₅ i	2.924422 — 0.564795i
ϵ_5^-	3.2554862 ⁸⁴⁵³⁴⁰¹⁰⁰ ₁₇₁₀₆₂₃₁₀	—1.111531 ⁵⁵³⁰¹⁴⁷⁶²⁵ ₆₂₀₄₄₂₅₄₈₀ i	3.255486 — 1.111531i
ϵ_6^+	3.5572158 ⁷⁶²²⁶⁵³⁵⁵ ₁₂₁₀₁₂₉₅₅	—1.7555060 ²¹²⁸⁷⁷⁸⁰⁰ ₈₅₄₁₃₀₂₄₀ i	3.557216 — 1.755506i
ϵ_7^-	3.824329 ⁵³⁶⁸⁷⁵⁷⁰³⁵ ₄₉₆₈₇₁₂₂₆₀	—2.487446 ¹⁶⁵¹⁴⁵⁷⁷⁴⁰ ₂₀₅₂₁₁₈₀₅₀ i	3.824330 — 2.487446i
ϵ_8^+	4.055434 ⁷¹⁶⁸⁹³⁴⁰⁷⁵ ₆₈₂₄₅₉₂₉₀₀	—3.2986410 ²³⁹⁵⁴¹⁷⁷⁵ ₅₈₃₈₈₂₉₇₅ i	4.055435 — 3.298641i
ϵ_9^-	4.2499550 ⁸⁰⁹⁵⁰³⁸⁴⁵ ₃₉₇₇₀₃₁₈₅	—4.183169 ²⁹²⁵⁸⁴⁴⁶⁵ ₇₀₄₃₈₅₂₉₀ i	4.249955 — 4.183170i
ϵ_{10}^+	4.4077725 ⁴⁹²⁴⁴⁸³²⁵ ₀₉₁₆₆₅₅₂₅	—5.1364531 ⁵³¹⁹¹⁹⁶⁵ ₉₃₂₇₀₂₆₅ i	4.407773 — 5.136453i
ϵ_{11}^-	4.528809 ⁵¹⁰¹⁴¹⁷⁰⁹⁰ ₄₇₀₀₆₃₄₂₉₅	—6.1547306 ¹⁰⁸¹⁶⁷⁶⁵ ₅₀₈₉₅₀₇₀ i	4.528809 — 6.154731i
ϵ_{12}^+	4.6130980 ⁹²⁹³⁰⁷⁶⁰⁵ ₅₄₃₅₄₇₁₄₅	—7.2353765 ²⁹⁹⁰⁹⁰¹⁰ ₆₈₄₈₅₀₈₀ i	4.613098 — 7.235377i
ϵ_{13}^-	4.661888 ⁴¹⁷⁶⁶²⁷⁰⁸⁰ ₂₈₀₉₇₄₃₃₅₀	—8.3755511 ⁴⁰⁵⁶⁹²⁴⁵ ₇₇₂₅₇₆₄₀ i	4.661888 — 8.375551i
ϵ_{14}^+	4.67345 ³⁰³⁴⁴²³⁸⁸⁰ ₂₉₈₃₄₈₆₅₈₆₀	—9.5707369 ⁴⁰¹³⁷²⁴⁰ ₉₁₀₇₄₅₄₅ i	4.673453 — 9.570737i
ϵ_{15}^-	4.6441405 ⁸³⁷⁸²⁴¹⁷⁰ ₄₉₃₄₈₂₉₆₅	—10.8262644 ²⁰⁶⁷⁹³⁹⁵ ₅₅₁₁₃₅₃₅ i	4.644141 — 10.82626i
ϵ_{16}^+	4.596328 ⁶⁰²¹⁹⁴⁸²³⁵ ₀₅₁₂₄₉₁₁₀₅	—12.142992 ⁰⁹²⁴⁶⁴⁸¹⁵ ₆₄₃₄₁₀₅₅₀ i	4.596328 — 12.14299i
ϵ_{17}^-	4.5198798 ⁹⁹⁹⁶⁵⁷⁵⁰⁰ ₆₃₇₅₅₆₇₆₅	—13.4658202 ¹⁷⁰⁹⁰³⁰⁰ ₇₉₆₂₄₇₅₀ i	4.519880 — 13.46582i
ϵ_{18}^+	4.26980 ⁷⁴¹⁴⁴⁴¹⁴⁵⁹⁵ ₆₈₈₇₈₀₄₂₉₉₀	—14.80688 ⁷⁵²⁰²⁶⁰³⁰⁵ ₈₀₄₆₈₉₇₄₉₅ i	4.269807 — 14.80689i
ϵ_{19}^-	3.47103 ⁸¹⁸⁸⁹³²²⁰⁷⁰ ₇₉₂₃₅₂₀₈₇₄₀	—15.422315 ⁴⁶⁸⁷⁸³⁵⁸⁵ ₇₃₄₁₉₄₉₅₀ i	3.471038 — 15.42232i
ϵ_{20}^+	3.5318492 ⁵¹⁵⁷³⁹⁸²⁵ ₁₈₁₃₉₂₇₈₅	—15.649544 ⁶⁹⁴⁷⁸⁵¹³⁰ ₇₂₈₂₁₉₈₇₀ i	3.531849 — 15.64954i
ϵ_{21}^-	4.28341 ⁶¹³⁰⁸⁶¹³²⁵⁰ ₅₆₀₈₄₆₂₃₂₁₀	—16.607294 ²⁷⁷³⁴⁴⁸⁹⁰ ₇₉₉₇₄₃₉₃₀ i	4.283416 — 16.60729i
ϵ_{22}^+	4.34585 ⁷⁸⁵⁸⁵⁹⁷⁸⁶⁷⁰ ₆₇₅₆₇₀₆₄₅₄₅	—17.86954 ⁸⁶⁰⁰⁹⁵²³³⁰ ₉₇₀₂₈₄₃₇₈₅ i	4.345857 — 17.86955i
ϵ_{23}^-	4.257915 ³⁰⁷⁴⁴¹⁹⁸⁴⁰ ₂₇₄₀₀₇₂₇₆₀	—18.9939478 ⁴⁷⁵⁵⁹³²⁰ ₈₀₉₉₄₀₇₀ i	4.257915 — 18.99395i
ϵ_{24}^+	4.1883310 ⁸⁷³³³⁰²⁹⁰ ₅₃₈₉₈₃₂₁₅	—19.9569140 ²⁰⁴⁶⁹⁴⁸⁰ ₅₃₉₀₄₂₃₀ i	4.188331 — 19.95691i
ϵ_{25}^-	4.25828 ⁴⁷¹⁷²⁴³⁷⁹⁶⁰ ₀₄₇₀₆₆₂₆₆₁₅	—20.87646 ⁵⁴⁸⁰³⁴²⁹⁷⁰ ₉₇₂₆₉₂₄₁₄₅ i	4.258283 — 20.87647i
ϵ_{26}^+	4.3820937 ⁹⁷⁰⁶¹³²³⁰ ₆₃₈₈₄₈₉₀₅	—21.8972996 ²⁵⁶²¹¹⁵⁰ ₅₈₇₉₇₆₃₀ i	4.382094 — 21.89730i
ϵ_{27}^-	4.431927 ⁸²¹⁰⁰⁵⁵⁴⁷⁰ ₇₅₄₆₅₂₆₉₈₅	—23.002819 ²⁵⁴⁹⁶⁰⁸⁸⁰ ₃₂₁₃₁₃₇₇₅ i	4.431928 — 23.00282i
ϵ_{28}^+	4.408917 ⁴¹³⁷⁹⁶⁹³⁸⁰ ₁₄₈₃₈₅₅₉₉₅	—24.142644 ²⁰¹⁴⁷⁹⁹⁷⁰ ₄₆₆₈₉₁₃₆₀ i	4.408918 — 24.14264i
ϵ_{29}^-	4.3856985 ⁴⁶³⁴⁹²⁸⁶⁰ ₁₀₁₃₉₂₀₉₀	—25.320304 ¹⁵¹²⁸⁵⁵⁶⁵⁰ ₂₁₃₈₂₀₀₄₅₀ i	4.385699 — 25.32030i
ϵ_{30}^+	4.432110 ⁵⁵⁸⁶³⁸⁹²⁹⁵ ₄₂₆₉₇₉₆₂₃₅	—26.518972 ⁴¹⁷⁶⁹³⁴⁸⁰ ₅₄₉₃₅₂₈₄₀ i	4.432110 — 26.51897i

(Continued.)

Table 4. (Continued.)

	guaranteed enclosures		numerical values by Rittby <i>et al.</i>
ϵ_{31}^-	4.50803 ⁴⁴⁷⁸⁰⁵⁹⁸⁸¹⁵ ₃₁₈₇₈₂₈₈₈₆₅	−27.70869 ³⁶⁴⁵⁸⁶¹³⁹⁰ ₄₇₂₅₄₉₇₆₄₅ i	4.508034 − 27.70869i
ϵ_{32}^+	4.56642 ⁹¹⁸⁹⁴⁵¹⁰⁶⁵ ₈₈₆₂₉₅₁₄₈₅	−28.9047568 ²¹¹⁵⁹⁴³⁰ ₅₃₈₀₉₄₄₀ i	4.566430 − 28.90476i
ϵ_{33}^+	4.597519 ⁴³⁵⁹⁵⁸¹⁵³⁰ ₃₈₇₂₅₂₅₈₅₀	−30.1026039 ⁶¹⁸⁶⁹⁸⁸⁰ ₉₅₇₃₇₉₆₀ i	4.597520 − 30.10260i
ϵ_{34}^+	4.581540 ⁹⁵³³⁰⁸⁴²³⁵ ₅₆₈₇₀₆₄₈₅₀	−31.296754 ⁴¹⁶¹⁹⁷⁷³⁰ ₇₃₆₈₃₄₅₂₀ i	4.581541 − 31.29675i
ϵ_{35}^+	4.477164 ³¹⁷⁴⁶⁷⁰⁹⁹⁰ ₂₇₄₉₅₉₈₉₂₅	−32.5108705 ⁶⁴²⁹⁵⁹⁵⁵ ₉₇₂₄₈₃₀₅ i	4.477164 − 32.51087i
ϵ_{36}^+	4.0888393 ⁵¹⁷⁵⁹⁵⁶⁵⁰ ₁₉₉₂₃₇₄₁₅	−33.9834345 ⁰⁷⁹²⁵³⁸⁰ ₃₉₇₆₁₂₇₀ i	4.088839 − 33.98343i
ϵ_{37}^+	3.43492 ⁷⁹⁹²⁷⁸²⁵⁶²⁵ ₅₉₇₃₀₆₉₄₁₉₀	−34.0167 ¹⁹⁴⁵⁰³⁴¹⁶⁰⁵ ₂₁₄₇₀₀₅₄₈₁₀ i	3.434927 − 34.01672i
ϵ_{38}^+	3.73004 ⁷⁹³⁴¹³⁰⁷⁵⁴⁰ ₅₂₆₃₅₅₀₂₆₁₅	−34.05210 ²⁵⁹⁴⁰⁵⁴⁹¹⁰ ₅₂₆₄₆₃₅₄₆₅ i	3.730047 − 34.05210i
ϵ_{39}^+	4.45878 ¹⁰²⁰²⁰⁶⁸⁰⁰⁰ ₀₉₈₆₂₆₂₈₄₃₀	−35.5634950 ²²⁴⁴⁰⁶²⁰ ₈₄₀₀₄₁₈₀ i	4.458781 − 35.56350i
ϵ_{40}^+	4.583226 ²⁰⁸²⁷⁹⁴⁸⁴⁵ ₀₅₇₉₉₈₀₈₁₅	−36.86441 ¹⁹³⁸⁴⁹⁵⁴⁸⁰ ₂₀₄₅₀₉₇₃₂₅ i	4.583226 − 36.86441i
ϵ_{41}^+	4.6275653 ³⁹²⁸⁶⁷⁰⁵⁵ ₀₇₄₅₀₈₈₀₀	−38.161068 ⁵⁹⁴¹⁵¹⁰¹⁰ ₆₂₅₉₈₆₉₀₅ i	4.627565 − 38.16107i
ϵ_{42}^+	4.614735 ⁹⁰⁸⁵⁹⁵⁶⁰⁴⁵ ₈₄₄₉₂₃₉₇₇₀	−39.471604 ⁴⁸⁵⁰⁸⁷⁹³⁵ ₅₄₈₇₅₉₆₄₅ i	4.614736 − 39.47160i
ϵ_{43}^+	4.57088 ¹²⁵²⁵⁰²³⁵⁶⁵ ₀₇₀₉₃₉₉₄₀₃₀	−40.80534 ⁰²³²⁶⁵⁹¹⁶⁰ ₁₂₁₇₆₇₄₇₅ i	4.570881 − 40.80534i
ϵ_{44}^+	4.512285 ¹⁷⁰³³¹⁷⁷²⁵ ₀₂₇₉₃₂₄₇₃₅	−42.087967 ²¹⁸⁶⁷¹¹⁶⁵ ₄₆₄₄₃₅₆₁₅ i	4.512285 − 42.08797i

Table 5. Excluded resonances.

	numerical values by Korsch <i>et al.</i>	chosen box side length l_k and angle θ	
$[\epsilon_{16}^+]$	4.589120 − 12.13151i	$l_{16} = 0.1$	$\theta = 0.7$
$[\epsilon_{17}^-]$	4.493625 − 13.49000i	$l_{17} = 0.1$	$\theta = 0.73$
$[\epsilon_{18}^+]$	4.362774 − 14.89927i	$l_{18} = 0.6$	$\theta = 0.73$
$[\epsilon_{19}^-]$	4.196770 − 16.35807i	$l_{19} = 1.0$	$\theta = 0.725$
$[\epsilon_{20}^+]$	3.995807 − 17.86525i	$l_{20} = 1.4$	$\theta = 0.72$
$[\epsilon_{21}^-]$	3.760076 − 19.41977i	$l_{21} = 1.5$	$\theta = 0.72$
$[\epsilon_{22}^+]$	3.489755 − 21.02065i	$l_{22} = 0.5$	$\theta = 0.725$
$[\epsilon_{23}^-]$	3.185024 − 22.66701i	$l_{23} = 1.4$	$\theta = 0.725$
$[\epsilon_{24}^+]$	2.846045 − 24.35801i	$l_{24} = 2.0$	$\theta = 0.725$
$[\epsilon_{25}^-]$	2.472984 − 26.09287i	$l_{25} = 2.0$	$\theta = 0.725$
$[\epsilon_{26}^+]$	2.065991 − 27.87087i	$l_{26} = 2.0$	$\theta = 0.72$
$[\epsilon_{27}^-]$	1.625219 − 29.69132i	$l_{27} = 2.0$	$\theta = 0.725$
$[\epsilon_{28}^+]$	1.150811 − 31.55357i	$l_{28} = 2.0$	$\theta = 0.725$

$y'(0) = 0$ with even eigenfunction (denoted by superscript '+'). The guaranteed enclosures we obtained for the four resonances $\hat{\lambda}_2^-, \hat{\lambda}_2^+, \hat{\lambda}_3^-, \hat{\lambda}_3^+$ are shown in table 6.

The computation of the resonance pair $\hat{\lambda}_3^-, \hat{\lambda}_3^+$ was performed in the same way as the enclosures described in §4*a*. Choosing $\theta = 0.735$, our provably correct computations showed that




Figure 3. Rotated box of side length $l_k/2$ excluding approximate resonance $\epsilon_k^\pm, k = 16^+, \dots, 28^+$, from Korsch *et al.* marked by asterisk. (Online version in colour.)

Table 6. Enclosures of resonance pairs near $\hat{\lambda}_2, \hat{\lambda}_3$ computed by Abramov *et al.* [14].

$[\hat{\lambda}_2^-]$	$0.91^{8411650805308}_{7362166176397}$	$-17.00^{0249282468423}_{1197218127467}i$	$\theta = 0.76$
$[\hat{\lambda}_2^+]$	$0.91^{8411650805308}_{7362166176397}$	$-17.00^{0249282468423}_{1197218127467}i$	$\theta = 0.76$
$[\hat{\lambda}_3^-]$	$2.55^{7790013021589}_{6817300474516}$	$-23.21^{200893060222}_{306897699860}i$	$\theta = 0.735$
$[\hat{\lambda}_3^+]$	$2.560^{188212981834}_{054480374655}$	$-23.208^{44496992380}_{57870390974}i$	$\theta = 0.735$

for each of the two boundary conditions there is only one resonance $\hat{\lambda}_3^-$ and $\hat{\lambda}_3^+$, respectively, in the disjoint boxes displayed in table 6. Moreover, they guarantee that in the larger λ -box $e^{-2i\theta}([23, 24] + [0.05, 1]i)$ containing these two boxes as well as the numerical value $\hat{\lambda}_3$ of Abramov *et al.* there is only one resonance for each of the two boundary conditions. Altogether, we thus proved that there is precisely one pair of disjoint resonances $\hat{\lambda}_3^- \neq \hat{\lambda}_3^+$ near the resonance approximation $\hat{\lambda}_3 = 2.46 - 23.22i$ of Abramov *et al.* and that this approximation has distance approximately $1 \cdot 10^{-1}$ to the true resonance pair $\hat{\lambda}_3^\pm$.

The computation of the resonance pair $\hat{\lambda}_2^-, \hat{\lambda}_2^+$ turned out to be much harder and computationally more expensive than all other enclosures and exclosures. To make it work, we had to use a slight modification of usual complex scaling, using stretching by some parameter $R > 0$ in addition to rotation of the variable by an angle $\theta \in [0, \pi/4)$. The potential $q_{\theta,R}$ and the eigenvalue parameter z in the spectral problem for the corresponding operator $H_{\theta,R}$ (compare (2.2), (2.3)) then become

$$q_{\theta,R}(x) := R^2 e^{2i\theta} (R^2 e^{2i\theta} x^2 - 1.6) e^{-R^2 e^{2i\theta} x^2 / 10}, \quad x \in \mathbb{R}, \quad z := R^2 e^{2i\theta} \lambda;$$

note that usual complex scaling corresponds to $R = 1$.

In order to apply Levinson's theorem, we needed to find suitable $X \geq 0, \theta \in [0, \pi/4)$ and $R > 0$ such that $\alpha_{X,\theta,R} := \int_X^\infty |q_{\theta,R}(x)| dx$ satisfies $\alpha_{X,\theta,R} < 1$. Proceeding as for usual complex scaling,




Figure 4. Excluded approximate resonances of Korsch *et al.* (asterisks) surrounded by respective excluding boxes, together with approximate resonances of Rittby *et al.* (circles), Abramov *et al.* (star and square) and analytic exclusion (grey-shaded) in the λ -plane. (a) Approximate resonances $0^+, \dots, 38^+$ of Rittby *et al.*, $0^+, \dots, 29^-$ of Korsch *et al.* and excluding boxes for approximate resonances $16^+, \dots, 28^+$ by Korsch *et al.* (b) Zoom into region marked by dashed line in (a) showing the approximate resonances $16^+, 17^-, 18^+$ with respective excluding boxes. (Online version in colour.)

instead of (3.3), we used

$$\alpha_{X,\theta,R} \leq \frac{10R^2}{\cos(2\theta)} e^{-R^2 \cos(2\theta)X^2/10} \left(X + \frac{1}{R^2} \frac{5}{\cos(2\theta)X} \right) =: A_{X,\theta,R}.$$

The main benefit of the additional stretching is that the upper bound $A_{X,\theta,R}$ decays exponentially fast in R . As for usual complex scaling, we then applied Taylor’s theorem with remainder in Lagrange form to obtain the rigorous computable upper bound $A_{X,\theta,R}^0 = 1.77 \cdot 10^{-17}$ for $X = 10, \theta = 0.76$ and $R = 10$.

With these parameters, we succeeded to enclose the resonances $\hat{\lambda}_2^-, \hat{\lambda}_2^+$ for the boundary condition $y(0) = 0$ and $y'(0) = 0$, respectively. The corresponding values in the z -plane are both in the box $1702.59_{49} + 5.4_3i$, hence

$$\hat{\lambda}_2^-, \hat{\lambda}_2^+ \in R^{-2} e^{-2i\theta} (1702.59_{49} + 5.4_3i) \subset 0.91_{7362166176397}^{8411650805308} - 17.00_{1197218127467}^{0249282468423}i.$$

Here, the first set is a box with midpoint $R^{-2} e^{-2i\theta} (1702.54 + 5.35i) \approx 0.918 - 17.001i$ and side length $R^{-2}10^{-1} = 1 \cdot 10^{-3}$, rotated clockwise by the angle $2\theta = 1.52$; the second set, which is the one displayed in table 6, is the smallest axis-parallel box containing this rotated box. Note

that these enclosures for $\hat{\lambda}_2^\pm$ differ in modulus by approximately $2 \cdot 10^{-2}$ from the value $\hat{\lambda}_2 = 0.92 - 17.02i$ calculated by Abramov *et al.* [14].

Hence, our guaranteed enclosures prove that not far from each of the two numerically computed values $\hat{\lambda}_2$ and $\hat{\lambda}_3$ of Abramov *et al.* there is indeed a pair of true resonances of (2.1); the distance is approximately $2 \cdot 10^{-2}$ for $\hat{\lambda}_2$ and approximately $1 \cdot 10^{-1}$ for $\hat{\lambda}_3$.

5. Conclusion

In this paper, we have presented a method which, for the first time, permits one to compute resonances in atomic physics with absolute certainty. At the same time, it allows one to detect with absolute certainty wrongly computed resonance approximations. The absolute reliability of our approach is based on a combination of interval arithmetic and the argument principle. To prove the efficiency of our method, we have established guaranteed enclosures for all numerical resonance approximations of Rittby *et al.* in [4,7] for problem (1.1) and guaranteed enclosures for the numerically computed values of Korsch *et al.* in [5] that are visible to complex scaling, thus definitely settling a dispute between these two groups of authors. The greatest challenge was to provably enclose two additional pairs of approximate resonances computed by Abramov *et al.* in [14] that were found neither by Rittby *et al.* nor by Korsch *et al.* Thus, we have proved the conjecture in [4,7] that the real parts of auto-ionizing resonances of certain atoms and molecules exhibit an oscillatory behaviour beyond a threshold and we have added new information on this threshold originating in the two new confirmed pairs of resonances.

Figure 4a displays all our results in the rectangle $0 \leq \text{Re}(\lambda) \leq 15$, $-70 \leq \text{Im}(\lambda) \leq 0$: in the top right corner of the λ -plane, the analytic exclusion from theorem 2.1 (grey-shaded), the enclosed approximate resonances $1^-, \dots, 38^+$ of Rittby *et al.* surrounded by circles, the additional ones by Abramov *et al.* as star and square, and the claimed approximate resonances $1^-, \dots, 29^-$ of Korsch *et al.* as asterisks; note that the resonances 0^+ , 29^- and $\hat{\lambda}_1$ to the left of the imaginary axis cannot be seen by the complex scaling method because of their negative real part. Around every disproved approximate resonances $16^+, \dots, 28^+$ of Korsch *et al.*, our excluding box is shown (grey-shaded). Figure 4b illustrates that for resonance 16^+ it was especially difficult to find a box that simultaneously excludes the computed value of Korsch *et al.* and does *not* contain the value computed by Rittby *et al.*

Acknowledgements. The authors gratefully acknowledge the support of the following funding bodies: Swiss National Science Foundation, SNF, grant no. 200020_146477 (S. Bögli, C. Tretter); EPSRC grant no. GR/R46410/01 (B.M. Brown); Leverhulme Trust grant no. RPG 167 (M. Marletta). C. Tretter thanks the Isaac Newton Institute of Mathematical Sciences, Cambridge, UK, where a first draft of this paper was written. All authors thank H. Siedentop for drawing their attention to this problem.

References

1. Jaulin L, Le Bars F. 2013 An interval approach for stability analysis: Application to sailboat robotics. *IEEE Trans. Robot.* **29**, 282–287.
2. Armellin R, Di Lizia P, Makino K, Berz M. 2012 Rigorous global optimization of impulsive planet-to-planet transfers in the patched-conics approximation. *Eng. Optim.* **44**, 133–155. (doi:10.1080/0305215X.2011.570757)
3. Berz M, Makino K, Kim Y-K. 2006 Long-term stability of the tevatron by verified global optimization. In *Nuclear instruments and methods in physics research section A: accelerators, spectrometers, detectors and associated equipment. Proc. 8th Intern. Comp. Accelerator Physics Conference ICAP 2004*, vol. 558, pp. 1–10.
4. Rittby M, Elander N, Brändas E. 1981 Weyl's theory and the complex-rotation method applied to phenomena associated with a continuous spectrum. *Phys. Rev. A* **24**, 1636–1639. (doi:10.1103/PhysRevA.24.1636)
5. Korsch HJ, Laurent H, Möhlenkamp R. 1982 Comment on: 'Weyl's theory and the complex-rotation method applied to phenomena associated with a continuous spectrum' [*Phys. Rev. A*

- (3) **24** (1981), no. 3, 1636–1639; by M Rittby, N Elander, E Brändas]. *Phys. Rev. A* **26**, 1802–1803. (doi:10.1103/PhysRevA.26.1802)
6. Moiseyev N, Certain PR, Weinhold F. 1978 Resonance properties of complex-rotated Hamiltonians. *Mol. Phys.* **36**, 1613–1630. (doi:10.1080/00268977800102631)
 7. Rittby M, Elander N, Brändas E. 1982 Weyl's theory and the method of complex rotation. A synthesis for a description of the continuous spectrum. *Mol. Phys.* **45**, 553–572. (doi:10.1080/00268978200100431)
 8. Engdahl E, Brändas E. 1988 Resonance regions determined by a projection-operator formulation. *Phys. Rev. A* **37**, 4145–4152. (doi:10.1103/PhysRevA.37.4145)
 9. Korsch HJ, Laurent H, Möhlenkamp R. 1982 Milne's differential equation and numerical solutions of the Schrödinger equation. II. Complex energy resonance states. *J. Phys. B* **15**, 1–15. (doi:10.1088/0022-3700/15/1/008)
 10. Rittby M, Elander N, Brändas E. 1982 Reply to 'Comment on 'Weyl's theory and the complex-rotation method applied to phenomena associated with a continuous spectrum'' [*Phys. Rev. A* (3) **26** (1982), no. 3, 1802–1803; by HJ Korsch, H Laurent, RM Öhlenkamp]. *Phys. Rev. A* **26**, 1804–1807. (doi:10.1103/10)
 11. Andersson N. 1993 A multiple-transition-point WKB investigation of complex energy resonances. *Int. J. Quant. Chem.* **46**, 375–381. (doi:10.1002/qua.560460305)
 12. Brown BM, Eastham MSP, McCormack DKR. 2000 Resonances and analytic continuation for exponentially decaying Sturm–Liouville potentials. *J. Comput. Appl. Math.* **116**, 181–193. (doi:10.1016/S0377-0427(99)00324-6)
 13. Hehenberger M, McIntosh HV, Brändas E. 1974 Weyl's theory applied to the Stark effect in the hydrogen atom. *Phys. Rev. A* **10**, 1494–1506. (doi:10.1103/PhysRevA.10.1494)
 14. Abramov AA, Aslanyan A, Davies EB. 2001 Bounds on complex eigenvalues and resonances. *J. Phys. A* **34**, 57–72. (doi:10.1088/0305-4470/34/1/304)
 15. Nedialkov NS, Jackson KR, Pryce JD. 1982 An effective high-order interval method for validating existence and uniqueness of the solution of an IVP for an ODE. *Reliab. Comput.* **7**, 449–465. (doi:10.1023/A:1014798618404)
 16. Brown BM, Langer M, Marletta M, Tretter C, Wagenhofer M. 2010 Eigenvalue enclosures and enclosures for non-self-adjoint problems in hydrodynamics. *LMS J. Comput. Math.* **13**, 65–81. (doi:10.1112/S1461157008000466)
 17. Simon B. 1978 Resonances and complex scaling: a rigorous overview. *Int. J. Quant. Chem.* **14**, 529–542. (doi:10.1002/qua.560140415)
 18. Siedentop H. 1987 On the localization of resonances. *Int. J. Quant. Chem.* **31**, 795–821. (doi:10.1002/qua.560310509)
 19. Harrell II, EM. 2007 Perturbation theory and atomic resonances since Schrödinger's time. In *Spectral theory and mathematical physics: a Festschrift in honor of Barry Simon's 60th birthday. Proc. Sympos. Pure Math.*, vol. 76, pp. 227–248. Providence, RI: American Mathematical Society.
 20. Edmunds DE, Evans WD. 1987 *Spectral theory and differential operators*. Oxford, UK: Oxford University Press.
 21. Aguilar J, Combes JM. 1971 A class of analytic perturbations for one-body Schrödinger Hamiltonians. *Commun. Math. Phys.* **22**, 269–279. (doi:10.1007/BF01877510)
 22. Cycon HL, Froese RG, Kirsch W, Simon B. 1987 *Schrödinger operators with application to quantum mechanics and global geometry*. Texts and Monographs in Physics, study edition. Berlin, Germany: Springer.
 23. Aslanyan A, Davies EB. 2000 Spectral instability for some Schrödinger operators. *Numer. Math.* **85**, 525–552. (doi:10.1007/PL00005391)
 24. Elander N, Rakityansky S. 2013 Resonances and their relations to spectral densities and scattering cross sections in the Schrödinger formulation. *Few-Body Syst.* **54**, 685–695. (doi:10.1007/s00601-012-0443-x)
 25. Brown BM, Langer M, Marletta M, Tretter C, Wagenhofer M. 2003 Eigenvalue bounds for the singular Sturm–Liouville problem with a complex potential. *J. Phys. A* **36**, 3773–3787. (doi:10.1088/0305-4470/36/13/310)
 26. Abramowitz M, Stegun IA. 1964 *Handbook of mathematical functions with formulas, graphs, and mathematical tables*. New York, NY: Dover Publications.