

Durham Research Online

Deposited in DRO:

13 October 2021

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Schuetz, Dirk (2021) 'Corrigendum to: A fast algorithm for calculating S-invariants.', *Glasgow Mathematical Journal* .

Further information on publisher's website:

<https://doi.org/10.1017/S001708952100032X>

Publisher's copyright statement:

This article has been published in a revised form in *Glasgow Mathematical Journal*, <https://doi.org/10.1017/S001708952100032X>. This version is published under a Creative Commons CC-BY-NC-ND. No commercial re-distribution or re-use allowed. Derivative works cannot be distributed. © The Author, 2021.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

**CORRIGENDUM TO: A FAST ALGORITHM FOR
CALCULATING S -INVARIANTS**

DIRK SCHÜTZ

Table 1 in [Sch21] claims to list all knots K with up to 15 crossings for which one entry of $s^{\mathbb{S}q^1}(K)$ differs from $s^{\mathbb{F}_2}(K)$. However, the table is incomplete. We list the missing knots in Table 1' below.

Knot	$s^{\mathbb{S}q^1}$	$s^{\mathbb{F}_2}$	$s^{\mathbb{F}_3}$
15n154386	(2, 2, 0, 0)	0	2
15n165952	(2, 2, 0, 0)	0	2
15n165966	(2, 2, 0, 0)	0	2
15n166064	(2, 2, 0, 0)	0	2
15n166244	(0, 0, -2, -2)	0	-2

TABLE 1'. Prime knots with non-standard $s^{\mathbb{S}q^1}$ missing from Table 1.

The original computation was done in batches of 10,000 knots. It appears that only the first 150,000 non-alternating 15-crossing knots were checked. A subsequent computation confirmed the results in Table 1, but also found the knots in Table 1' among the remaining 18,030 non-alternating 15-crossing knots.

REFERENCES

[Sch21] Dirk Schütz, *A fast algorithm for calculating S -invariants*, Glasg. Math. J. **63** (2021), no. 2, 378–399. MR 4244204

DEPARTMENT OF MATHEMATICAL SCIENCES, DURHAM UNIVERSITY

Email address: dirk.schuetz@durham.ac.uk